

TOXICITY ASSESSMENT OF ANABAENA SP. FOLLOWING EXPOSURE TO COPPER OXIDE NANOPARTICLES AND SODIUM CHLORIDE

KARIMI, R.¹ – NORASTEHNIA, A.^{2*} – ABBASPOUR, H.¹ – SAEIDISAR, S.¹ – NAEEMI, A. S.²

¹Faculty of Science, Damghan Branch, Islamic Azad University, Damghan, Iran

²Department of Biology, Faculty of Science, University of Guilan
Rasht P.C. 4193833697, Iran
(phone: +98-13-3333-3648; fax: +98-13-3333-3647)

*Corresponding author
e-mail: norasteh@guilan.ac.ir

(Received 10th Mar 2017; accepted 2nd Jun 2017)

Abstract. The widespread use of nanomaterials in recent years has caused rising pollution in surface and waste waters. It may be seriously affecting aquatic life such as cyanobacteria and thus, there is a compelling need to evaluate the potential effects of these substances in aquatic ecosystems. Copper oxide is one of the most common nanoparticles (CuO-NPs) used today. In this study, the responses of *Anabaena* sp. to CuO-NPs, salt stress and their combined effects (CuO-NPs+NaCl) were investigated. After 72 h exposure, MDA (malondialdehyde), proline, protein content, soluble sugar, total phenols and the activities of antioxidative enzymes such as peroxidase (POD), catalase (CAT), ascorbate peroxidase (APX), and polyphenol oxidase (PPO) were measured using spectrophotometry. Scanning electron microscopy (SEM) was used to image the results of exposure of cyanobacterial cells to nanoparticles. The effective concentrations (EC10, EC50, EC90), no observed effect concentration (NOEC) and percent inhibition (I %) was also calculated after 24, 48 and 72 hours. EC and NOEC data for CuO-NPs treatments were considerably higher than were those for NaCl treatments. CuO-NPs had a significant impact on the shape and morphology of *Anabaena* sp. cells and resulted in their swelling and enlargement. Modulations in enzymatic and non-enzymatic antioxidants were clearly evident in *Anabaena* sp. exposed to salt and CuO-NPs stress. Lipid peroxidation, measured in terms of MDA levels, increased with CuO-NPs and CuO-NPs+NaCl stress. Flavonoid and proline contents were also increased by CuO-NPs stress. Similarly, increases in POD and CAT activities were more pronounced in the CuO-NPs+NaCl treatments. On the other hand, APX activity enhancement was strongest in *Anabaena* sp. exposed to CuO-NPs.

Keywords: antioxidant activity, aquatic ecosystems pollution, cyanobacteria, CuO-NPs, salinity, two combinational stress

Introduction

Nanomaterials, products of nanotechnology, are an important part of numerous industrial and medical products and their use has increased dramatically in the past decade (Peralta-Videa et al., 2011). Being so widely used, it is likely that they will increasingly contaminate aquatic ecosystems (Klaine et al., 2008). Despite this obvious eventuality, the toxic ecological effects of nanomaterials have nevertheless not been easily evaluable; the mechanisms of toxicity of nanoparticles are not understood (Dhawan et al., 2009). The key physicochemical features of nanomaterials associated with toxicity which have been identified by toxicological researchers to date include factors such as aggregation, solubility, size, shape, elemental

composition, contact area, porosity, surface ionic charge, and hydrodynamic diameter (Griffitt et al., 2008; Johnston et al., 2010; Peralta-Videa et al., 2011).

Biological responses of organisms are very much dependent on the physicochemical properties of nanoparticles. For example, nanoparticles are highly mobile in water and thus they can easily pollute aquatic ecosystems (Oukarroum et al., 2012). Nanotoxicity, as one of the most important aspects of nanotechnology, is induced by nanomaterials whose safety and toxicity characteristics are not clearly defined; and therefore must be determined to ensure public safety. In addition, high concentrations of salts, especially sodium chloride, are among the most important abiotic factors which can affect the survival of living organisms, including aquatic organisms such as cyanobacteria (Denizet et al., 2011). The presence of toxic ions, mainly Na^+ , creates distress due to osmotic potential and their effect on the uptake of inorganic nutrients (Deniz et al., 2011). This leads directly to inhibitory effects on growth, oxidative stress and finally death. It has been shown that high salinity inhibits photosynthesis, resulting in a decrease in the carbon pool of freshwater cyanobacteria (Srivastava et al., 2008) and lowering of the synthesis of compatible solutes (Ferjani et al., 2003). Furthermore, although nano metals and metal-oxides (Haulik et al., 2014) and salinity have harmful effects on cyanobacteria and plants individually, their effect cannot be directly extrapolated when two or more combinational stresses are induced (Suzuki et al., 2014). Algal cells, in contrast to human and animal cells, possess a cellulose cell wall, which acts as a barrier to particle uptake, prohibiting uptake of nanoparticles larger than 20 nm in size. Nevertheless, nanoparticles are capable of inducing pore formation, which may result in uptake of larger-sized particles (Navarro et al., 2008). Cyanobacteria are major organisms in aquatic productivity and significantly contribute to the availability of nitrogen in agriculture (Singh, 1961) and the evolution of oxygen-dependent respiration. *Anabaena*, as one of the significant cyanobacteria, is not only widely distributed across a wide range of salinities (Srivastava et al., 2011), able to cope with high osmotic potential by enhanced osmolyte synthesis (Hagemann, 2011), but also a preferred candidate for dinitrogen fixation (Singh, 1961). On the other hand, phytoplanktons are used as an early and direct warning signal because of their rapid response to stresses (Del Arco et al., 2014). However, not many studies have previously been conducted on the combined effect of salt and CuO-NPs in cyanobacteria in general and *Anabaena* in particular, although their exposition to many kinds of stresses, individually, has been reported previously (Rai et al., 2013).

In the present study, the impacts of combined effect of CuO-NPs and salt stress were investigated on antioxidant activity and growth inhibition of *Anabaena* sp. collected from Guilan wetlands.

Materials and methods

Nanoparticles

CuO-NPs produced by US Research Nanomaterials Company, were used. The size of used CuO nanoparticles was 40 nm. A 2000 mg/l stock solution was used for preparing appropriate concentrations (Table 1 and Fig. 1).

Table 1. CuO nanoparticle characteristics

Purity	99%
Color	Black
Particle size	40nm
Specific area	20m ² /g~
Morphology	nearly spherical
Bulk Density	0.79g/cm ³
True Density	6.4g/m ³




Figure 1. SEM of CuO-NPs

Release of Cu²⁺ From CuO-NPs

Stocks of nanoparticles at concentrations of 5, 10, 20 and 25 ppm were prepared with distilled water and the particles dispersed by sonication (Misonix Sonicator 3000, Iran) for 60 minutes. The pH was adjusted to 8.5 using NaOH. After 24 hours, 10 mL of the shaken up solution was taken from each flask, centrifuged 20 minutes at 4500 rpm and filtered through a 150 nm filter membrane. The content of Cu²⁺ was measured by atomic absorption flame spectrometer (Varian Spectra AA 220FS Varian Spectra Co. Australia).

Isolation of *Anabaena* sp. and preparation of culture medium

Cyanobacteria *Anabaena* sp. were collected from Guilan wetlands and isolated by repeated plating on solid Zander culture medium (Miller, et al., 1978). The pH of the culture medium was adjusted to 6.8, and the culture temperature was maintained at 24 ± 2 °C with aphoto period of 12 hours light/12 hours darkness. Light intensity was 60 µmol photon m⁻²S⁻¹. Culture medium was sterilized by autoclaving at 121°C for 15 min and then refrigerated at 6°C.

Preparation of main treatments

The main stock of nanoparticles was prepared by 60 minutes ultrasonic treatment. Six treatments and one control with three replicates for each treatment and control were

prepared. The concentrations of nanoparticles used in the experiments were 0, 5, 10, 15, 20 and 25 mg/l. 4×10^4 cells from the main stock of *Anabaena* sp. were added to 40 ml of the treatments by dilution of primary algal stock. This step was performed under completely sterile conditions. The test tubes were then stored at $24 \pm 2^\circ\text{C}$ which was adjusted by a thermostat. All samples were maintained at constant light periodic condition (12 hours darkness and 12 hours light) for the 72 hours of the experiment.

Determination of growth rates in cyanobacteria

Growth inhibitory effects were studied according to the OECD201 method (Handy et al., 2012). At 24, 48 and 72 hours after the start of the experiment, aliquots of the solution were removed from the test tubes using Pasteur pipettes and the cells present were counted in a Neubauer chamber under an optical microscope. After counting and recording the data, the average number of cells in the top and bottom squares was calculated and then the number of cells was calculated using the following formula:

Cell density per ml

$$= \text{average cells counted in the large square} \times 10^4 \times \text{dilution factor}$$

In each experiment, percent inhibition values were calculated using spectrophotometric data compared with the growth in control systems. EC values were calculated using linear regression analysis of transformed chemical concentration as natural logarithm data versus percent inhibition.

Experimental design and stress application

The EC_{30} doses of single and combined salt and CuO-NPs treatments were determined using the plate colony count method (Rai and Raizada., 1985). A 0.052 M solution of NaCl was used for salt treatment. In addition, 12.58 mg/L of CuO-NPs was used. For the NaCl+CuO-NPs study, the two doses were applied simultaneously. Biochemical evaluations were determined similarly 72 h after treatment. All experiments were done in triplicate.

Antioxidant assays

Assay for total phenolics

Measurement of total phenol was performed by method of Gao et al. (2000), using Folin- Ciocalteu reagent and a standard of gallic acid. For this purpose, 100 μl of extract (2 mg/ ml) was poured into a glass tube. Then, 200 μl Folin solvent and 2 ml of distilled water were added to the extract. 1 ml of sodium carbonate (21%) was added after 9 minutes to the glass tube in darkness. Finally, the samples were stored at room temperature in the dark for an hour. Blanks were prepared with 100 ml of solvent, instead of the extract. Absorptions were read at 765 nm. The total phenolic content was determined as gallic acid equivalents (mg GAE/g extract).

Assay for total flavonoids

Total flavonoid content was determined using a spectrophotometer according to Arvouet-Grand *et al.* (1994). 1 ml of extract solution with 2 mg/ml concentration was mixed with 1 ml of 2% aluminum trichloride (AlCl_3) methanolic solution. Absorbance of

the reaction mixtures were measured at 415 nm against a blank after 10 min. Quercetin (QE) was employed as a standard reference and the total flavonoids content of the extracts was expressed as μg quercetin equivalents per gram of extract (μg QE/g extract).

Extraction and measurement of proline

Proline content in the samples was measured using the method of Bates et al. (1973). For this purpose, 1 g fresh biomass of Cyanobacteria was harvested and their cell walls were broken using 4 ml of 3% sulfosalicylic acid and sonication for 10 minutes (5 pulse seconds and 5 seconds rest). The resulting solution was centrifuged for 20 minutes at a speed of 13,000 rpm and immediately transferred to an ice bath. Finally, 4 ml of toluene was added to the reaction solution and absorbance was measured at 520 nm. Proline content in micrograms per gram fresh weight was calculated.

Total soluble protein

The measurement of total protein was conducted using 2 mg/ml extracts and Coomassie Brilliant Blue G-250 staining solution - in 95% ethanol and 85% ortho phosphoric acid (Bradford, 1976) using bovine serum albumin as a standard.

Lipid peroxidation content assay

Lipid peroxidation was measured in terms of the total content of 2-thiobarbituric acid (TBA)-reactive substances and expressed as equivalent of MDA (malondialdehyde), extinction coefficient 155 mM^{-1} , using the method of Heath and Packer (1968).

Estimation of enzymatic antioxidants

Measuring the activity of catalase (CAT)

The method of Aebi (1984) was used for investigation of catalase activities. For this purpose, 2.5 ml of 0.05 M phosphate buffer was mixed with 0.2 ml of 3% hydrogen peroxide and 0.2 ml enzyme extract. Absorbance was read at 240 nm for 2 minutes. The enzymatic activity of the enzyme was reported as units/mg of protein.

Measuring the activity of ascorbate peroxidase (APX)

APX activity was determined at an absorbance of 290 nm (extinction coefficient $2.8 \text{ mM}^{-1} \text{ cm}^{-1}$) for 1 min in 1 ml reaction mixture containing 50 mM potassium phosphate buffer (pH 7.0), 0.5 mM ascorbic acid, 0.1 mM H_2O_2 , and 200 μl of enzyme extract (Nakano and Asada., 1981)

Measuring the activity of polyphenol oxidase (PPO)

Activity of polyphenol oxidase was assayed according to the method of Grigori (1993). 3 ml reaction solutions containing 2.5 ml phosphate buffer 50 mM (pH =7.2), 0.2 ml Pyrogallol 0.02 M and 300 Micro liters of enzyme extract were used. Changes in absorption were read at 430 nm for 2 minutes. The enzyme activity was calculated based on units of enzyme in 1 gram of fresh weight with extinction coefficient of $0.001 \text{ mM}^{-1}/\text{cm}^{-1}$.

Measuring the activity of Guaiacol peroxidase (POD)

Kinetic activity of Guaiacol peroxidase was determined according to Kalir et al. (1984). First, 1 ml of the reaction solution containing 475 μ l Guaiacol 45 mM, 475 μ l hydrogen peroxide 100 mM and 50 μ l enzyme supernatant was prepared. Changes in absorption due to Guaiacol oxidation were read by spectrophotometer at 470 nm for 2 minutes. An extinction coefficient of $26.6 \text{ mM}^{-1}/\text{cm}^{-1}$ was used in the calculation of the enzyme activity. Peroxidase enzyme activity was calculated in Unit/gr FW. Each enzymatic unit is the amount of enzyme which increases 0.01% of the absorption at 470 nm per minute.

Electronic images

Scanning electron microscopy (SE-SEM, MIRA3 TESCAN) was used to investigate the effects of nanoparticles on the shape and size of cells in microscopic tissues of *Anabaena* sp. and to image the surfaces. Imaging of the surfaces of control and NP-treated cells was performed on cells exposed to the 10.93 mg/L NP concentration, which had affected 30% of the cells ($\text{EC}_{30} = 10.93 \text{ mg/L}$).

Statistical analysis

Each treatment was replicated three times; the results were reported as the average of three parallel determinations of the mixture of three replicated samples. All statistical analyses were carried out using SPSS 16.0. One Way ANOVA with Duncan's test was used to determine significant changes in the results ($p < 0.05$).

Results

Release of Cu^{2+}

Suspensions of 0, 5, 10, 20 and 25 mg/l Cu^{2+} nanoparticles caused the release of 0, 0.017, 0.116, 0.249 and 0.310 mg/l of Cu^{2+} respectively, as assayed by atomic absorption flame spectrometry. After 24 hours, less than 0.31 mg/l Cu^{2+} was detected in our experiments (Fig. 2).




Figure 2. Cu^{2+} release from N- CuO

Effects on growth rates

Anabaena growth in the presence of various concentrations of copper nanoparticles and salt is shown in Figure 3. The results indicate that cyanobacterial growth decreased by increasing the concentration of salt particles as compared with the control. The highest reduction occurred at 0.32 M NaCl and 25 mg/l CuO-NPs (Fig. 3). This was visually apparent, as seen by yellowing and fragmentation of filaments.




Figure 3. Growth rate of *Anabaena* sp. after exposure to different concentrations of CuO-NPs (A) and NaCl (B). The data represents the average of three replicates $\pm$ standard error (SE). Different letters indicate significant differences among treatments according to Duncan's test with $p < 0.05$.

The results of 72-hour toxicity tests, the 24, 48 and 72-hour levels of EC_{10} , EC_{30} , EC_{50} and EC_{90} for CuO-NPs and NaCl toxicity are summarized in Tables 2 and 3. The range of EC_{30} values varied from 177.82 mg L^{-1} (24 h) to 12.58 mg L^{-1} (72 h) for CuO-NPs tests and from 0.114 M (24 h) to 0.052 M (72 h) for NaCl.

Table 2. The values of EC_{10} , EC_{30} , EC_{50} , EC_{90} and NOEC for CuO-NPs

Time (hours)	EC_{10} (mg/l)	EC_{30} (mg/l)	EC_{50} (mg/l)	EC_{90} (mg/l)	NOEC (mg/l)
24	13.7	177.82	1047.12	79432.82	104.71
48	8.91	18.19	29.51	97.72	2.95
72	7.76	12.58	17.37	38.9	1.73

Table 3. The values of EC_{10} , EC_{30} , EC_{50} , EC_{90} and NOEC for NaCl

Time (hours)	EC_{10} (M)	EC_{30} (M)	EC_{50} (M)	EC_{90} (M)	NOEC (M)
24	0.035	0.114	0.248	1.07	0.024
48	0.03	0.085	0.168	0.928	0.016
72	0.022	0.052	0.095	0.398	0.095

The effects on non-enzymatic antioxidants

The content of protein and soluble sugars

Protein content in all CuO-treatments showed a significant decrease compared to the control (Table 4). The percentage decreases of protein, relative to the control value were

13, 44.8 and 41% in NaCl, CuO and NaCl+CuO-treated cultures, respectively. However cellular contents of soluble sugars after treatment with NaCl and CuO-NPs+NaCl increased significantly (*Table 4*).

Table 4. Content of protein, proline, soluble sugars, total phenol, flavonoids and MDA in *Anabaena* sp. under NaCl, CuO-NPs, NaCl+CuO-NPs treatments. The data represents the average of three replicates $\pm$ standard error (SE). Different letters indicate significant differences among treatments according to Duncan's test with $p < 0.05$.

Column	Protein (mg/g FW)	Proline (mg/l)	Soluble Sugars (mg/l)	Total Phenol (mg/l)	Flavonoids (mg/l)	MDA (nmol/g FW)
Control	2.94 $\pm$ 0.25 a	6.27 $\pm$ 0.25 bc	15.65 $\pm$ 0.45 b	13.5 $\pm$ 0.59 a	0.016 $\pm$ 0.0004 bc	6.25 $\pm$ 0.02 b
NaCl	2.6 $\pm$ 0.34 ab	5.26 $\pm$ 0.02 c	63.47 $\pm$ 3.23 a	13.33 $\pm$ 1.24 a	0.015 $\pm$ 0.0002 c	9.89 $\pm$ 0.22 a
CuO	1.62 $\pm$ 0.39 b	10.21 $\pm$ 0.51 a	18.05 $\pm$ 0.1 b	13.6 $\pm$ 0.15 a	0.024 $\pm$ 0.0014 a	7.29 $\pm$ 0.29 b
NaCl+CuO	1.75 $\pm$ 0.12 b	7.42 $\pm$ 0.44 b	65.96 $\pm$ 1.83 a	13.87 $\pm$ 0.90 a	0.017 $\pm$ 0.0002 b	8.81 $\pm$ 0.63 a

The effect on malondialdehyde

Lipid peroxidation was assayed by measuring malondialdehyde (Zhanga et al., 2013). Changes of malondialdehyde in cyanobacteria *Anabaena* sp. under stress from copper nanoparticles, salinity and both stresses at the same time is shown in *Table 4*. Malondialdehyde content significantly increased under CuO-NPs and CuO-NPs+NaCl compared to control.

Proline content

Proline content in treatments with NaCl and NaCl+CuO-NPs showed no significant changes. However, in the CuO-NPs treatment, proline content increased 64.1 % at the 10.93 mg/l level of CuO-NPs (*Table 4*).

Total phenols and total flavonoids

Phenolic content did not differ significantly between the three types of treatment compared and with the control group. Flavonoid content significantly increased 63.1% in CuO treatment compared to the control, while there was no significant difference between control, NaCl and NaCl+CuO treatments.

The effect on enzymatic antioxidants


In this study, the enzyme activity of POD in treatment with CuO-NPs and NaCl was found to be significantly increased. In contrast, polyphenol oxidase (PPO), another enzyme that is involved in neutralizing ROS compounds and assayed in this study, was significantly reduced in all treatments compared to control. CAT activity showed a different behavior, with a significant increase compared to control (~ 600 %, $p < 0.05$) in treatment with NaCl+CuO as well as (~ 400 %, $p < 0.05$) in treatment with CuO, while it remained unchanged in NaCl treatments; and finally, the other peroxide scavenging enzyme, APX, was also found to be induced under CuO-NPs stress (*Table 5*).

Table 5. Enzymatic antioxidant content in *Anabaena* under NaCl, CuO-NPs, NaCl+CuO-NPs treatments. The data represents the average of three replicates $\pm$ standard error (SE). Different letters indicate significant differences among treatments according to Duncan's test with $p < 0.05$.

Factors Treatments	POD (Unit/g FW)	PPO (Unit/g FW)	APX (Unit/g FW)	CAT (Unit/mg protein)
Control	0.008 $\pm$ 0.0006b	2200 $\pm$ 57.7a	0.446 $\pm$ 0.026c	0.0025 $\pm$ 0.0006c
NaCl	0.005 $\pm$ 0.0013a	1666.7 $\pm$ 33.3b	0.625 $\pm$ 0.026b	0.005 $\pm$ 0.0006c
CuO	0.009 $\pm$ 0.0012a	1666.7 $\pm$ 33.3b	0.834 $\pm$ 0.015a	0.0095 $\pm$ 0.001b
NaCl+ CuO	0.016 $\pm$ 0.0013a	1000 $\pm$ 9c	0.521 $\pm$ 0.039c	0.0152 $\pm$ 0.0019a

Electron microscopy

CuO-NPs attached to the surface of the cyanobacterial cells in direct contact with them is demonstrated clearly by SEM images in *Figure 4*. Generally, electron microscopic images showed that the nanoparticles bound to each other and, more relevantly, to the surface of the cyanobacterial cells, altering the appearance of cells through disruption of direct physical interactions between the cells and increasing the number of lysed cells (*Fig. 4, A-E*).






Figure 4. SEM images of *Anabaena* sp. after 72 hours exposure to CuO-NPs at concentration of 10.93mg/l. Figures of A and C show control samples and figures B, D and E show *Anabaena* cells which treated with CuO-NPs.

Discussion

As illustrated in Figure 2, small amounts of Cu^{2+} were released. This can be attributed to the aggregation of CuO-NPs and the slightly alkaline pH environment of the suspension (Wang et al., 2013). Cells cannot exclude Cu^{2+} and accumulate it in high quantity. Cu^{2+} becomes powerfully toxic for phytoplankton communities at concentrations between 0.1 and 1.0 mg/l (Deniz et al., 2011). The release of Cu^{2+} which is seen in our experiment is in this range too. Although algal cells are able to encase and detoxify Cu^{2+} but this can occur in restricted ranges (Toncheva et al., 2006). The decrease in the rate of cell division caused by metals has been primarily attributed to their binding to sulfhydryl groups which are important for regulating plant cell division (Visviki and Rachlin., 1991). Furthermore, The difference between effective concentrations ($\text{EC}_{10,30,50}$) in two treatments indicated that the degree of nanoparticle toxicity is also likely to depend on their nano structures and high surface to mass ratio as well as the nature of their constitutive element (Hernández Battez et al., 2010). This varies between algal species (Pendashte et al., 2013). The results of Shi (2011) have also indicated that CuO-NPs toxicity is three to four times higher than that of ionic Cu, because of the larger uptake of NPs-released Cu. Therefore, as it is shown in Figure 2, cell counting decreased during CuO-NPs treatment, compared to NaCl treatment.

As is shown in Table 4, percentages of proteins decreased in treated samples relative to control samples meaningfully. Rahman et al. (2011) observed similar findings in cyanobacteria under metal stress; Cu^{2+} treatment also resulted in reduction of protein content extensively. This decline may be due to production of ROS, which is known to damage protein, and therefore disturbs the cellular homeostasis. Similar observations were made, also, in *A. doliolum* exposed to cadmium and UV-B stressed (Bhargava et al., 2007). In contrast, adaptation to salt stress in cyanobacteria is generally accomplished by excretion of inorganic ions from the cells to balance osmotic potential; cells can also prevent denaturation of macromolecules via accumulation of some osmolytes, such as sucrose, trehalose, glucosylglycerol, glycine betaine, proline and/or glutamate (Allakhverdiev et al., 2005).

Lipid peroxidation is linked to the production of O_2^- . The presence of high amounts of transitional metals such as Cu (II) favors the enhanced generation of OH° from O_2^-

through the Fenton reaction (Luna et al., 1994). Thus, the increased levels of MDA suggest that Cu (II) ions stimulated free radical formation in *Anabaena* sp.

Increases in proline concentration under heavy metal and salt stresses have been reported in some higher plants and in the cyanobacteria *Spirulina*, *Anabaena* and *Cylindrospermum* (Rahnama and Ebrahimzadeh., 2004; Chris et al., 2006; Choudhary et al., 2007). It seems that an adaptive mechanism via increase in proline is activated for neutralization of accumulated CuO-NPs and the acidity. Proline's chelating ability to bind metal ions can also be a defense mechanism for survival (Deniz et al., 2011).

It has been recently shown that accumulation of secondary metabolites, such as phenylpropanoids, including flavonoids, which has scarcely been reported in cyanobacteria, could be one of the strategies for cyanobacterial organisms to protect against cellular damage (Singh et al., 2014). Flavonoids, phenolic acids, and tannins, as well as some derivatives of flavonoids, are also compounds which have many different biological activities including antioxidant activity under stress conditions (Agati et al., 2011), in addition to some medicinal activities (Fresco et al., 2006). Although only scarce reports are available on total phenols and total flavonoid content under salt and nano particle stress tolerance responses in cyanobacteria, positive correlations with the accumulation of polyphenolics, enhanced antioxidant activity and tolerance to stresses have been reported in plant species (Korkina, 2007). As a result, there is no additive response of non-enzymatic antioxidant factors in NaCl+CuO-NPs combination treatment compared to their responses to single stresses. This may be due to their antagonistic effects (Rai et al., 2013).

During oxidative stress, POD is a very important enzyme able to remove the dangerous radical products of superoxide and hydrogen peroxide (Beier et al., 1991). The greater activity of POD in treated compared to control plants can be attributed to stress caused by the treatments (Cosio and Dunand, 2010). These results suggest that NaCl and CuO-NPs individually as well as NaCl+CuO-NPs can be very important stressors impacting plant physiology. Reduction of PPO in both NaCl and CuO-NPs-treated cells was more than other treatments. In our study, CAT activity significantly increased compared to control in all treatments of CuO-NPs. Free Cu²⁺ can catalyze the formation of highly toxic ROS such as hydroxyl radicals (OH[•]) from superoxide anions (O₂^{•-}) or hydrogen peroxide (H₂O₂) (Melegaria et al., 2013). Therefore, increase in CAT activity of treated samples compared to control very probably is a response to substantial ROS production. APX induction may be attributed to the *apx* operon induction by H₂O₂ production (Vranova et al., 2002). Therefore, it seems that H₂O₂ can be detoxified by a combined effort of POD, CAT and APX in *Anabaena* sp. That enhanced levels of most antioxidant enzymes, CAT, POD and APX, were observed for *Anabaena* sp. in this study indicates the protective role of these enzymes against copper-induced oxidative stress (Table 5). Finally, it can be noted that, except for CAT, which shows more activity in NaCl+CuO-NPs treatments compared to NaCl and CuO-NPs treatments, there were no additive activities in other enzymatic antioxidants.

CuO-NPs often increased cell aggregation by forming bridges between the cells in the areas where the nanoparticles had accumulated. The CuO-NPs attached to the surface of the cyanobacterial cells and the direct contact is demonstrated clearly by the SEM images in Figure 4. Their external surfaces had begun to fracture compared with the surfaces of cells that were not in contact with these particles. The structure of the cell wall was probably affected by the metal ions (Fig. 4E). Similarly, others have found cellular ultrastructural changes in *Microcystis aeruginosa* which were exposed to Ag-

NPs (1 mg l^{-1}) (Duong et al., 2016). After 48 h exposure with Ag-NPs the structure of *Microcystis* was changed; the cells were shrunken and distorted. Other studies on the impact of nanoparticles on other species have yielded similar results (Ayatallahzadeh Shirazi et al., 2015).

Conclusion

The outcomes of this study suggest that single stresses of NaCl and CuO-NPs can have inhibitory effects on *Anabaena* sp. in the used ranges, inducing severe damages in high concentrations. However, stress induced by these factors in combination did not result in elevated inhibitory effects, more than those observed in single stresses. This may be related to interactions between released ions from NaCl and CuO-NPs.

Acknowledgements. We would like to thank Mehdi Rassa for the English Grammar revision.

REFERENCES

- [1] Aebi, H. (1984): Catalase in vitro. – *Methods in Enzymology* 105: 121-126.
- [2] Agati, G., Biricolti, S., Guidi, L., Ferrini, F., Fini, A., Tattini, M. (2011): The Biosynthesis of flavonoids is enhanced similarly by UV radiation and root zone salinity in *L. vulgare* leaves. – *Journal of Plant Physiology* 168: 204-212.
- [3] Allakhverdiev, S. I., Klimov, V. V., Hagemann, M. (2005): Cellular energization protects the photosynthetic machinery against salt-induced inactivation in *Synechococcus*. – *BBA-Bioenergetics* 1708: 201-208.
- [4] Arvouet-Grand, A., Vennat, B., Pourrat, A., Legret, P. (1994): Standardisation d'un extrait de propolis et identification des principaux constituants. – *Journal de Pharmacie de Belgique* 49: 462-468.
- [5] Ayatallahzadeh Shirazi, M., Shariati, F., Keshavarz, A. K., Ramezanzpour, Z. (2015): Toxic Effect of aluminum oxide nanoparticles on green micro-algae *Dunaliella salina*. – *International Journal of Environmental Research* 9(2): 585-594.
- [6] Bates, L. S., Wadern, R. P., Teare, I. D. (1973): Rapid estimation of free proline for water stress determination. – *Plant Soil* 39: 205-207.
- [7] Beier, K., Fahimi, H. D. (1991): Environmental pollution by common chemicals and peroxisome proliferation: efficient detection by cytochemistry and automatic image analysis. – *Progress in Histochemistry and Cytochemistry* 23: 150-163.
- [8] Bhargava, P., Atari, N., Srivastava, A. K., Rai, L. C. (2007): Cadmium mitigates ultraviolet-B stress in *Anabaena doliolum*: enzymatic and non-enzymatic antioxidants. – *Biologia Plantarum* 51: 546-550.
- [9] Bradford, M. M. (1976): A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein dye-binding. – *Analytical Biochemistry* 38: 248-252.
- [10] Choudhary, M., Jetley, U. K., Khan, M. A., Zutshi, S., Fatma, T. (2007): Effect of heavy metal stress on proline, malondialdehyde, and superoxide dismutase activity in the cyanobacterium *Spirulina platensis*-S5. – *Ecotoxicology and Environmental Safety* 66: 204-209.
- [11] Chris, A., Zeeshan, M., Abrahama, G., Prasad, S. M. (2006): Proline accumulation in *Cylindrospermum* sp. – *Environmental and Experimental Botany* 57: 154-159.
- [12] Cosio, C., Dunand, C. (2010): Transcriptome analysis of various flower and silique development stages indicates a set of class III peroxidase genes potentially involved in pod shattering in *Arabidopsis thaliana*. – *BMC Genomics* 11: 528.

- [13] Del Arco, A. I., Guerrero, F., Jiménez-Gómez, F., Parra, G. (2014): Shifts across trophic levels as early warning signals of copper sulfate impacts in plankton communities. – *Applied Ecology and Environmental Research* 12: 493-503.
- [14] Deniz, F., Saygideger, S. D., Karaman, S. (2011): Response to copper and sodium chloride excess in *Spirulina* sp. (Cyanobacteria). – *Bulletin of Environmental Contamination and Toxicology* 87:11–15.
- [15] Dhawan, A., Sharma, V., Parmar, D. (2009): Nanomaterials: a challenge for toxicologists. – *Nanotoxicology* 3: 1–9.
- [16] Duong, T. T., Le, T. S., Huong Tran, T. T., Nguyen, T. K., Ho, C. T. Dao, T. H., Le, T. P. Q., Nguyen, H. C., Dang, D. K., Le, T. T. H., Ha, P. T. (2016): Inhibition effect of engineered silver nanoparticles to bloom forming cyanobacteria. – *Advances in Natural Sciences: Nanoscience and Nanotechnology* 7: 1-7.
- [17] Ferjani, A., Mustardy, L., Sulpice, R., Marin, K., Suzuki, I., Hagemann, M. Murata, N. (2003): Glucosylglycerol, a compatible solute, sustains cell division under salt stress. – *Plant Physiology* 131: 1628-1637.
- [18] Fresco, P., Borges, F., Diniz, C., Marques, M. P. M. (2006): New insights on the anticancer properties of dietary polyphenols. – *Medicinal Research Reviews* 26: 747-766.
- [19] Gao, X., Ohlander, M., Jepsson, N., Bjork, L., Trajkovski, V. (2000): Changes in antioxidant effects and their relationship to phytonutrients in fruits of sea buckthorn (*Hippophae rhamnoides* L.) during maturation. – *Journal of the Agricultural and Food Chemistry* 48: 1485-1490.
- [20] Griffitt, R. J., Luo, J., Gao, J., Bonzongo, J. C., Barber, D. S. (2008): Effects of particle composition and species on toxicity of metallic nanomaterials in aquatic organisms. – *Environmental Toxicology and Chemistry* 27: 1972-1978.
- [21] Grigori, R. P. F., Bendal, D. S. (1966): The purification and some properties of the polyphenol oxidase from tea (*Camellia sinensis* L.). – *Biochemical Journal* 101(3): 569-581.
- [22] Hagemann, M. (2011): Molecular biology of cyanobacterial salt acclimation. – *FEMS Microbiology Reviews* 35: 87-123.
- [23] Handy, R. D., Van den Brinke, N., Chappell, M., Muhling, M., Behra, R., Dusinska, M., Simpson, P., Ahtinen, J., Jha, A. N., Seiter, J., Bednar, A., Kennedy, A., Fernandes, T. F., Riediker, M. (2012): Practical considerations for conducting ecotoxicity test methods with manufactured nanomaterials: what have we learnt so far? – *Ecotoxicology* 21(4): 933-972.
- [24] Haulik, B., Balla, S., Pálfi, O., Szekeres, L., Juri'ková, T., Sály, P., Bakonyi, G. (2014): Comparative ecotoxicity of the nano Ag, TiO₂ and ZnO to aquatic species assemblages. – *Applied Ecology and Environmental Research* 13(2): 325-338.
- [25] Heath, R. L., Packer, L. (1968): Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation. – *Archives in Biochemistry and Biophysics* 125: 189-198.
- [26] Hernández Battez, A., Viesca, J. L., González, R., Blanco, D., Asedegbega, E., Osorio, A. (2010): Friction reduction properties of a CuO-nanolubricant used as lubricant for a NiCrBSi coating. – *Wear* 268: 325-328.
- [27] Johnston, H. J., Hutchison, G. R., Christensen, F. M., Aschberger, K., Stone, V. (2010): The biological mechanisms and physicochemical characteristics responsible for driving fullerene toxicity. – *Journal of Toxicological Sciences* 114: 162-182.
- [28] Kalir, A., Ornri, G., Poljakoff-Mayber, A. (1984): Peroxidase and catalase activity in leaves of *Halimione portulacoides* exposed to salinity. – *Physiologia Plantarum* 62: 238-244.
- [29] Klaine, S. J., Alvarez, P. J. J., Batley, G. E., Fernandes, T. F., Handy, R. D., Lyon, D. Y., Mahendra, S., McLaughlin, M. J., Lead, J. R. (2008): Nanomaterials in the environment: behavior, fate, bioavailability, and effects. – *Environmental Toxicology and Chemistry* 27: 1825-1851.

- [30] Korkina, L. G. (2007): Phenylpropanoids as naturally occurring antioxidants from plant defence to human health. – *Cellular and Molecular Biology* 53: 15-25.
- [31] Luna, C. M., Gonzalez, V. S., Trippi, V. S. (1994): Oxidative damage caused by excess copper in oat leaves. – *Plant and Cell Physiology* 35: 11–15.
- [32] Melegaria, S. P., Perreault, F., Ribeiro C. R. H., Popovi, R., Gerson, M. W. (2013): Evaluation of toxicity and oxidative stress induced by copper oxide nanoparticles in the green alga *Chlamydomonas reinhardtii*. – *Aquatic Toxicology* 142-143: 431-440.
- [33] Miller, W. E. (1978). The *Selenastrum capricornutum* Printz algal assay bottle test: experimental design, application, and data interpretation protocol. – Corvallis, Or.: Corvallis Environmental Research Laboratory, Office of Research and Development, U.S. Environmental Protection Agency.
- [34] Nakano, Y., Asada, K. (1981): Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. – *Plant and Cell Physiology* 22: 867-880.
- [35] Navarro, E., Piccapietra, F., Wagner, B., Marconi, F., Kaegi, R., Odzak, N., Sigg, L., Behra, R. (2008): Toxicity of silver nanoparticles to *Chlamydomonas reinhardtii*. – *Environmental Science and Technology* 42(23): 8959-8964.
- [36] Oukarroum, A., Bras, S., Perreault, F., Popovic, R. (2012): Inhibitory effects of silver nanoparticles in two green algae, *Chlorella vulgaris* and *Dunaliella tertiolecta*. – *Ecotoxicology and Environmental Safety* 78: 80-85.
- [37] Pendashte, H., Shariati, F., Keshavarz, A., Ramzanpour, Z. (2013): Toxicity of zinc oxide nanoparticles to *Chlorella vulgaris* and *Scenedesmus dimorphus* algae species. – *World Journal of Fish and Marine Sciences* 5(5): 563-570.
- [38] Peralta-Videa, J. R., Zhao, L., Lopez-Moreno, M. L., de la Rosa, G., Hong, J., Gardea-Torresdey, J. L. (2011): Nanomaterials and the environment: a review for the biennium 2008–2010. – *Journal of Hazardous Materials* 186: 1-15.
- [39] Rahman, Md. A., Soumya, K. K., Tripathi, A., Sundaram, S., Singh, S., Gupta, A., (2011): Evaluation of sensitivity of cyanobacteria, *Nostoc muscorum* and *Synechococcus* PCC 7942 for heavy metals stress – a step toward biosensor. – *Toxicological and Environmental Chemistry* 93: 1982-1990.
- [40] Rahnama, H., Ebrahimzadeh, H. (2004): The effect of NaCl on proline accumulation in potato seedlings and cali. – *Acta Physiologiae Plantarum* 26: 263-270.
- [41] Rai, L. C., Raizada, M. (1985): Effect of nickel and silver ions on survival, growth, carbon fixation and nitrogenase activity of *Nostoc muscorum*: regulation of toxicity by EDTA and calcium. – *Journal of General and Applied Microbiology* 31: 329-337.
- [42] Rai, S., Singh, S., Srivastava, A. K., Rai, L. C. (2013): Salt and UV-B induced changes in *Anabaena* PCC 7120: physiological, proteomic and bioinformatic perspectives. – *Photosynthesis Research* 118: 105-114.
- [43] Shi, J., Abid, A. D., Kennedy, I. M., Hristova, K. R., Silk, W. K. (2011): To duckweeds (*Landoltia punctata*), nanoparticulate copper oxide is more inhibitory than the soluble copper in the bulk solution. – *Environmental Pollution* 159: 1277-1282.
- [44] Singh, R. N. (1961): Role of Blue-Green algae in the nitrogen economy of Indian agriculture. – New Delhi: Indian Council of Agricultural Research.
- [45] Singh, D. P., Prabha, R., Meena, K. K. Sharma, L., Sharma, A. K. (2014): Induced accumulation of polyphenolics and flavonoids in cyanobacteria under salt stress protects organisms through enhanced antioxidant activity. – *American Journal of Plant Sciences* 5: 726-735.
- [46] Srivastava, A. K., Bhargava, P., Thapar, R. Rai, L. C. (2008). Salinity-induced physiological and proteomic changes in *Anabaena doliolum*. – *Environmental and Experimental Botany* 64: 49-57.
- [47] Srivastava, A. K., Alexova, R., Jeon, Y. J., Kohli, G. S., Neilan, B. A. (2011): Assessment of salinity-induced photorespiratory glycolate metabolism in *Anabaena* sp. PCC 7120. – *Microbiology* 157: 911-917.

- [48] Suzuki, N., Rivero, R. M., Shulaev, V., Blumwald, E., Mittler, R. (2014): Abiotic and biotic stress combinations. – *New Phytologist* 203: 32-43.
- [49] Toncheva–Panova, T., Merakchiyska, M., Djingova, R., Ivanova, J., Sholeva, M., Paunova, S. (2006): Effect of Cu^{2+} on the red microalga *Rhodella reticulata*. – *General and Applied Plant Physiology, Special Issue*: 53-60.
- [50] Visviki, I., Rachlin, J. W. (1991): The toxic action and interactions of copper and cadmium to the marine alga *Dunaliella minuta* in both acute the chronic exposure. – *Archives of Environmental Contamination and Toxicology* 2: 271-275.
- [51] Vranova, E., Inze, D., Breusegem, F. V. (2002): Signal transduction during oxidative stress. – *Journal of Experimental Botany* 53: 1227-1236.
- [52] Wang, L., Wang, M., Peng, Ch., Pan, J. (2013): Toxic Effects of Nano-CuO, Micro-CuO and Cu^{2+} on *Chlorella* sp. – *Journal of Environmental Protection* 4: 86-91.
- [53] Zhanga, W., Xiong, B., Chena, L., Lin, K., Cui, X., Bi, H., Guo, M., Wang, W. (2013): Toxicity assessment of *Chlorella vulgaris* and *Chlorella protothecoides* following exposure to Pb(II). – *Environmental Toxicology and Pharmacology* 36: 51-57.