

MORPHOLOGICAL AND PHYSIOLOGICAL RESPONSES OF PURPLE CHRYSANTHEMUM (*ASTER SPHATHULIFOLIUS*) UNDER LONG-TERM STRESS OF CALCIUM CHLORIDE AS DEICING SALT

JU, J.-H.¹ – YEUM, K.-J.² – SON, H.-M.³ – YOON, Y.-H.^{1*}

¹Department of Green Technology Convergence, Konkuk University, 268 Chungwondaero, Chungju-si, Chungcheongbuk-do 27478, South Korea

²Division of Food Bioscience, Konkuk University, 268 Chungwondaero, Chungju-si, Chungcheongbuk-do 27478, South Korea

³Department of Green Technology Convergence, Graduate School of Natural Science, Konkuk University, 268 Chungwondaero, Chungju-si, Chungcheongbuk-do 27478, South Korea

*Corresponding author

e-mail: yonghan7204@kku.ac.kr; phone: +82-43-840-3538; fax: +82-43-851-4169

(Received 3rd Aug 2017; accepted 5th Dec 2017)

Abstract. Long-term research on the effect of deicers on groundcover plants can open up usage of groundcover plants for ornamental purposes on the roadside affected by deicer salt. The objective of this study was to investigate the long-term effect of CaCl₂ on morphological and physical responses of purple chrysanthemum (*Aster sphathulifolius*) grown in a greenhouse. Five different concentrations (0, 1, 2, 5, 10, and 15 g/L) of road deicer (CaCl₂ 74%) solutions (100 ml) were applied twice a week for a 5 month period. Survival rate, growth parameter, biomass, and physiological indices were measured. Increased CaCl₂ concentration resulted in decreased survival rate, especially at concentrations higher than 10 g/L. Exposure to increasing CaCl₂ concentrations resulted in dramatic decreases in growth index, number of leaves, leaf width, and leaf length at concentration higher than 2 g/L. Biomass was also negatively affected by increasing deicer salt stress, with shoot mass being reduced more than root fresh weight. Chlorophyll *b* content was decreased, while chlorophyll *a* and proline contents in leaves had a gradual increase when plants were exposed to increasing salt stress. Although a clear roadside negative effect did exist, there was no significant difference between plants under 1 g/L of CaCl₂ and control treatment for 5 months. Our results suggest that *Aster sphathulifolius* planting could be highly beneficial to the roadside or urban areas with mildly salt-affected soils.

Keywords: calcium chloride; ornamental groundcovers, road deicer; soil-plant continuum; tolerance of salt stress

Introduction

Deicing salt has been used for decades to melt snow and ice from the road to improve traffic safety in winter (Viskari and Kärenlampi, 2000). However, these salts are eventually displaced to roadside areas where they can negatively impact soils, vegetation, and water resources (ground or surface) (Devitt et al., 2014). A high amount of deicer has been used in South Korea due to its geographically distinctive winter season characterized by snowfall. On roads, CaCl₂ has been used as one of the most popular road deicer in South Korea (Shin et al., 2010), because it is more effective at low temperatures. Thus, at roadside areas that use CaCl₂, the Cl⁻ concentration is higher than the other sites. The use of deicing salts on roads has resulted in high concentrations of chloride (577 ~ 2,353 mg/kg) in urban roadside soil (Zhang et al., 2012). This is probably due to heavy volume of traffic at these sites (Baek et al., 2014). With

increasing of salts content, a large amount of ions enter the cells, thereby morphology and physiology of plants growing under salt stress are adversely affected (Cao et al., 2012). Even though CaCl_2 has on plant physiology include membrane permeability and reduction in Na^+ concentration within a certain limit (Amuthavalli et al., 2012). Therefore, ways to reduce salt damage should be presented including a planting guide to match species to a site. The harmful effects of deicing salt on roadside herbs and grasses depends on the sensitivity of plant species (Ali et al., 2012). Planting salt tolerant ornamental groundcover plants can be a very effective strategy to utilize deicer salt affected soils. Salt tolerance varies widely among plant species and genotype. Plants can adapt to salinity by tolerating or avoiding salt uptake. Some plants can achieve salt tolerance by osmotic adjustment (Mao et al., 2008). Many researchers have studied salt tolerance of herbaceous plants after they are exposed to high salts. Eom et al. (2007) have shown that salt tolerance of six groundcover species is different in response to NaCl treatment (0 ~ 400 mM). It can be grouped into three categories: highly sensitive to salt treatment (*Sedum acre*), intermediately sensitive (*Achemilla mollis*, *Nepeta* × *faassenii*, *Thymus praecox*, and *Phlox subulata*), and tolerant (*Solidago cutleri*). Ali et al. (2012) have reported that the ornamental plant *Anternanthera bettzickiana* can be characterized as a salt-tolerant glycophyte. Other studies have also reported salt tolerance of other ornamental herbaceous plants, including *Suaeda salsa* (Guan et al., 2011), *Foeniculum vulgare* (Semiz et al., 2012), *Eugenia myrtifolia* (Acosta-Motos et al., 2015), Aster perennials (Wu et al., 2016), *Salvia splendens* and *Ageratum houstonianum* (Jędrzejuk et al., 2016), and *Sedum* species, *Allium* species, and a mixture of turf grasses (Whittinghill and Rowe, 2011). Salt tolerance has also been demonstrated for woody plants such as *Ardisia japonica* (Lee et al., 2008), ornamental shrubs (Cassaniti et al., 2009), *Prosopis glandulosa* (Moore et al., 2010), *Atriplex nummularia* (Alharby et al., 2014), and *Rosa rubiginosa* (Hura et al., 2017). However, to the best of our knowledge, no study has reported the salt tolerance of *Aster spathulifolius* for a long-term. Also, the effect of calcium chloride deicer on ornamental herbaceous plants compared to NaCl remains to be investigated.

Asteraceae is one of the largest plant families with many important ornamental species (Wu et al., 2016). Seashore spatulate aster, specifically, *Aster spathulifolius* Maxim, is halophyte, which typically found on the coast areas of Korea. Because the demand for road deicers continues to increase, long-term research on the effect of deicers on groundcover plants is needed. Therefore, the objective of this study was to determine the effect of CaCl_2 on morphological and physiological responses of *Aster spathulifolius*.

Material and methods

Plant material and growth conditions

Seedlings of *Aster spathulifolius* were purchased from a commercial nursery (Sannea Botanical Garden, Chenonan, Chungnam, Korea). All plants with similar sizes used in experiments were 5-6 cm in height with 10-15 fully grown leaves. These seedlings were transferred to 12 cm-diameter pots filled with 0.5kg of artificial substrates (Wonjo-mix, NongKyung Inc., Korea) in March 2016. Pots were kept in the laboratory and watered every 2 days in the four weeks following transplanting. The experiment was conducted from April 2016 to October 2016 at KonKuk University, Chungju (latitude, 35°49'N; longitude, 127°08'E). Plants were grown in the greenhouse under

natural light. During the study period, air temperature and relative humidity were monitored with a thermo recorder (SK-1260, SATO, Japan). Photosynthetically active radiation (PAR) was measured with a digital light meter (Extech 401025, EXTECH, USA). Average air temperature, relative humidity, PAR were kept at 26.2°C, 62.7%, and 1,500 $\mu\text{mol}\cdot\text{m}^2/\text{s}$, respectively.

Treatments

The substrate used for this study was a commercially produced ridging (Nongkyung Floricultural materials, Co., Chungbuk, Korea) with pH 6.5. Prior to transplanting, each metal was thoroughly mixed with 0.5 kg of air-dried substrate. The mixture was then used to fill the pots. To evaluate the effect of CaCl_2 concentrations on *Aster spathulifolius*, a completely randomized block design with five treatments was adopted (3 replications, 3 seedlings per replication, total of 45 seedlings). Based on the results of earlier studies (Zhang et al., 2012; Thouvenot et al., 2012) on roadside soil concentration of chloride ions, CaCl_2 powder (74% of calcium chloride, Oriental Chemical Industries, Korea) was diluted in distilled water to obtain concentrations at 0 (Control), 1 (C1), 2 (C2), 5 (C5), 10 (C10), and 15 g/L (C15) corresponding to 0.5, 7, 14, 35, 70, and 105 mM, respectively. *Aster spathulifolius* seedlings were watered with 100 ml of CaCl_2 solution twice a week until the end of the experiment (for 5 months).

Measurements

The number of plants that were survived was recorded. This number was then used to estimate the survival rate (percentage). Plants that maintained 3 to 6 green or greenish and elongating leaves were scored as being alive (survived). Those with all leaves dried out were scored as dead. Survival rates were calculated with the following formula (Kanawapee et al., 2012): $\text{Survival rate} = (\text{survived plants} / \text{total plants}) \times 100 (\%)$.

The following growth parameters were observed: growth index, leaf number, leaf length, and leaf width. Growth index was measured once a month for each group of seedlings in May and September 2016 during peak growth period. Plant height (H) at the tallest point and width at the widest point in two directions (left to right and front to back: W1, W2) were also measured (Whittinghill and Rowe, 2011). Data of height and width were used to calculate growth index $[(W1 + W2)/2 + H]/2$ which is commonly used as an indicator of plant size (Hammond et al., 2007). Length of leaf, width of leaf, and leaf number on the stem were also measured for *Aster spathulifolius*. These growth parameters were partially based on the relative growth rates (RGR) of plants (Thouvenot et al., 2012): $\text{RGR} = (\ln L_2 - \ln L_1) / (T_2 - T_1)$, where L1 and L2 were total length at time 1 and time 2, respectively.

Each plant harvested was then divided into shoots (leaves and stem) and roots after 5 months of treatments. Fresh weights (FW) were measured after different organs were washed with distilled water. Shoots and roots were then dried in a drying oven (C-DF, Changshin Scientific Co., Korea) at 70 °C until they reached a constant weight in order to measure their respective dry weights (DW). To evaluate salinity tolerance, relative dry weight (RDW) was calculated as a ratio of average values for each accession of seedlings (Chen et al., 2013): $\text{RDW} = \text{DW (salt treatment)} / \text{DW (control)} \times 100 (\%)$, a trait commonly used to measure salinity tolerance.

Chlorophyll was extracted following the method outlined by Baruah et al. (2014). Briefly, collected leaves of experimental plants were washed properly and 100 mg of

fresh leaves from those plants was weighed and cut into small pieces with a razor. Chlorophyll pigment was then extracted by grinding these cut leaves with a mortar and pestle for 5 min in about 8 ml of 95 % (v/v) acetone. The extract was filtered with Whatman number 1 filter paper. The filtrate was transferred to a 100-ml volumetric flask. The volume of the filtrate was increased to 10 ml by addition of 95 % acetone. After that, optical density (OD) of the extract was measured using a spectrophotometer (Biochrom Libra S22, Biochrom, England) at wavelength of 645 nm and 663 nm using 95% acetone as a blank. The wavelengths chosen the maximum absorption wavelengths for total chlorophyll and chlorophyll a/b, respectively. The amount of chlorophyll a and b and total chlorophyll content of leaf tissues (in mg/g) were calculated using the following equations:

$$\text{Chlorophyll a} = [12.7(\text{OD}_{663 \text{ nm}}) - 2.69(\text{OD}_{645 \text{ nm}})] \times (V/1000W) \quad (\text{Eq.1})$$

$$\text{Chlorophyll b} = [22.9(\text{OD}_{645 \text{ nm}}) - 4.68(\text{OD}_{663 \text{ nm}})] \times (V/1000W) \quad (\text{Eq.2})$$

$$\text{Total chlorophyll} = [20.2(\text{OD}_{645 \text{ nm}}) + 8.02(\text{OD}_{663 \text{ nm}})] \times (V/1000W) \quad (\text{Eq.3})$$

Proline content was analyzed with the modified procedure of Kanawapee et al. (2012). Briefly, the third and fourth leaves from the apical shoot of three plants per treatment were frozen immediately in liquid nitrogen at harvest. Approximately 0.1 g of leaf was homogenized with 5 ml of 3% aqueous sulfosalicylic acid. Two ml of the extract was then reacted with 2 ml of acid ninhydrin and 2 ml of glacial acetic acid followed by boiling in a water bath at 100°C for 1 h. The reaction was stopped by placing the tubes on ice. The solution was then extracted with 4 ml of toluene and the absorbance of the toluene fraction was measured by spectrophotometry (Biochrom Libra S22, Biochrom, England) at wavelength of 520 nm. The amount of free proline was evaluated using a standard curve and expressed as µg/g tissue fresh weight.

Statistical analysis

Data are presented as mean ± standard deviation of nine replicates (N = 9). Statistical analysis was performed using SPSS 18.0 for Windows (SPSS Inc., Chicago, IL, USA) by one-way analysis of variance (ANOVA). Treatment means were separated with Duncan's Multiple Range Test ($P \leq 0.05$).

Results

The survival rate of *Aster sphathulifolius* showed no significant difference between treatments in the initial stage. However, at the end of 5 months under salt stress conditions, survival rates of plants were reduced sharply in the following order: CaCl₂ 0 (Control; 100%) > CaCl₂ 1 (C1; 100%) > CaCl₂ 2 (C2; 80%) > CaCl₂ 5 (C5; 70%) > CaCl₂ 10 (C10; 0%) > CaCl₂ 15 g/L (C15; 0%). The survival rate of *Aster sphathulifolius* were significantly decreased compared to those of control as well as 1 g/L of CaCl₂ treated plants. Increasing CaCl₂ concentration in the substrates resulted in decreasing survival rate, especially when its concentration was higher than 10 g/L (Figure 1).






Figure 1. Survival rates of *Aster spathulifolius* after treated with CaCl_2 at different concentrations (0, 1, 2, 5, 10, and 15 g/L as Control, C1, C2, C5, C10, and C15, respectively) for 5 months. Values are presented as means \pm SE of nine replications.


The growth index of *Aster spathulifolius* showed various degrees of growth retardation, although some differences existed between treatments with different CaCl_2 concentrations in June. However, the average growth indices of Control, C1, C2, C5, C10, and C15 were significantly decreased ($P \leq 0.05$) by 17.3, 15.2, 15.2, 11.8, 8.2, 6.7 cm, respectively in July, and plants grown in C10 and C15 treatment groups did not survive in September. The average growth indices of C1, C2, and C5 treatment groups were 20, 22, and 47%, respectively, compared to the control. For all treatments, the number of leaves showed a increase resulting in a range from 14.4 (C15) to 66.5% (Control) deicing salt stress in June, but then gradually decreased by 3.1 (Control) and 82.6% (C15) in response to salinity stress. The number of leaves of plants in C1, C2, and C5 treatment groups was significantly lower ($P \leq 0.05$) by 9, 18, and 57%, respectively, compared to that in the control at 5 months after treatment. However, there was no significant difference in the number of leaves between the Control and C1 treatment. Exposure to increasing CaCl_2 concentrations resulted in dramatic decreases in leaf width and leaf length with CaCl_2 at 1, 2, 5, and 10 $\text{g}\cdot\text{L}^{-1}$, leaf widths were 32, 33, 22, and 19 mm, respectively, while leaf lengths were 89, 92, 84, and 62 mm, respectively. There was no significant difference in leaf width or length between plants under Control and those with C1 treatment (Figure 2).




(A) : Growth index



(B): Number of leaves



(C) : Leaf length



(D) : Leaf width

Figure 2. Growth index(A), number of leaves(B), leaf width(C), and leaf length(D) of *Aster sphathulifolius* grown under CaCl_2 stress at various concentrations (0, 1, 2, 5, 10, and 15 g/L as Control, C1, C2, C5, C10, and C15, respectively) for 5 months. Values are presented as means \pm SE of nine replications.

As shown in *Figure 3*, exposure to increasing concentrations of calcium chloride resulted in dramatic decrease ($P \leq 0.05$) in biomass of *Aster sphathulifolius* compared to non-salt stress. Shoot showed more reduction than root fresh weight of *Aster*

spathulifolius. Average shoot fresh weights of plants in C1, C2, C5, C10, and C15 treatment groups were reduced by 3, 36, 56, 89, and 92%, respectively, while root fresh weights were reduced by 41, 50, 66, 83, and 88%, respectively in comparison with those in the Control (CaCl_2 0 g/L). Shoot dry weights of salt stressed plants in C1, C2, C5, C10, and C15 treatment groups were also reduced by 24, 31, 52, 75, and 81%, respectively, while dry weights were reduced by 39, 46, 59, 67 and 74%, respectively, compared to those of Control (Figure 4). Relative dry weight (RDW), a trait commonly used to measure salinity tolerance, was also substantially reduced by 76.4, 69.5, 48.5, 25.4, 19.4% relative to Control (data not referred).




Figure 3 Fresh(A) and dry weight(B) of *Aster spathulifolius* grown under CaCl_2 stress with at various concentrations (0, 1, 2, 5, 10, and 15 g/L as Control, C1, C2, C5, C10, and C15, respectively) measured after harvest. Values are presented as means \pm SE of nine replications (* P value \leq 0.05).

Salt stress for 5 months affected chlorophyll contents of *Aster spathulifolius*. Interestingly, salt stress resulted in significant increase of chlorophyll *a* content, whereas decrease of chlorophyll *b* content. The mean chlorophyll *a* content was increased from 8.2 to 14.97 mg/g FW. For chlorophyll *b*, the mean was decreased from 21.18 to 11.17 mg/g FW respectively. Total chlorophyll contents in C1, C2, C5, and

C10 treatments were decreased significantly by 98.6, 95.3, 92.9, and 89.5%, respectively, compared to those of Control. When *Aster sphathulifolius* plants were exposed to salt stress, the mean proline contents in the leaves were increased significantly from 23.0 in the control to 37.3 µg/g FW in C10 treatment (an increase of 62.2%). Salinity stress caused a significant increase in proline contents in C2, C5, and C10 treatment groups when compared with the control and C1 treatment group (Table 1).

Table 1. Differences of chlorophyll and proline contents of *Aster sphathulifolius* leaves grown under CaCl₂ stress at various concentrations (0, 1, 2, 5, 10, and 15 g/L as Control, C1, C2, C5, C10, and C15, respectively) measured after harvest.

Treatments	Chlorophyll contents a (mg/g FW)	Chlorophyll contents b (mg/g FW)	Total chlorophyll contents (mg/g FW)	Proline contents (µg/g FW)
Control	8.29 d ^z	21.18 a	29.46 a	23.00 c
C1	12.92 b	16.18 b	29.09 ab	23.33 c
C2	12.64 c	15.49 b	28.12 abc	26.00 bc
C5	12.88 b	14.56 b	27.43 bc	29.67 b
C10	14.96 a	11.47 c	26.44 c	37.33 a
C15	-	-	-	-

^z Different letters in the same column indicate significant difference according to Duncan's multiple range test at $P \leq 0.05$ (n=9).

Discussion

Plant survival and productivity is crucial for seedling establishment under saline conditions (Ponte et al., 2014). In this study, no significant correlation between plant survival and CaCl₂ concentrations were found initially, however, after treatment for 5 months, a significant negative correlation was observed between the survival rate and the dose of CaCl₂. Higher CaCl₂ concentrations were associated with lower survival rate for a long term exposure, although *Aster sphathulifolius* is a halophyte that has grown and evolved under saline conditions. Soil salinity can inhibit plant growth by a number of mechanisms including low external water potential, toxicity of absorbed Cl⁻ ions, inhibition of various enzymatic activities and different cellular processes, and interference with the uptake of essential nutrients (Taffoue et al., 2014). The fundamental mechanisms of salt tolerance in salt tolerant plants seem to be mostly dependent on their capacities to sequester toxic ions in vacuoles and accumulate compatible osmotic pressure in the cytoplasm as previously suggested (Munns, 2002). Salt tolerance of herbaceous perennial species including *Achemilla mollis*, *Nepeta × faassenii*, *Sedum acre*, *Tymus praecox*, *Phlox subulata*, and *Solidago cutleri*, to aqueous solution of sodium chloride at 0~400 mM over a 21 day period has been evaluated by measuring their growth. These plants were grouped into three tolerance categories : highly sensitive to salt treatment (*Sedum acre*), those with intermediate sensitivity (*A. mollis*, *N. × faassenii*, *T. praecox*, and *P. subulata*), and those with salt tolerance (*S.*

cutleri) (Eom et al., 2007). It has been reported that *Alternanthera bettzickiana* can grow even at a salinity level of 40 dS/m (Ali et al., 2012). In addition, it has been found that the critical level of species sensitive to chloride ions is 4 ~ 7 mg/g and that of species tolerant to chloride ions is 15 mg/g (Xu et al., 2000; White and Broadley, 2001). Considering this, it is generally accepted that *Aster sphathulifolius* could be able to grow at salt levels less than 1 g/L (Cl^- agent $0.094 \text{ g/L day}^{-1}$) for 5 months.

Biomass of seedlings is a trait commonly used to measure salinity tolerance (Chen et al., 2013). *Eupatorium greggii*, *Viguiera stenoloba*, and *Santolina chamaecyparissus* were the most salt-tolerance species with less reductions in shoot dry weight (Wu et al., 2016). It has been reported that *Alternanthera bettzickiana* plants at salinity level of 20 dS/m will produce 30.3% less biomass than controls. Further increasing salinity will lead to lower biomass in response to higher salt stress (Ali et al., 2012). Similarly, Semiz et al. (2012) have also reported that the biomass of *Foeniculum vulgare* is also affected negatively by chloride ions. Meanwhile, chloride ions (Cl^-) in soil is recognized as one key contributor to the decrease in the production of plants. The critical concentration of chloride ions is found to be 0.49 g/L. At this concentration, the biomass of plants is decreased by 10% (Dang et al., 2008). Considering the actual concentrations of chloride ions (Cl^-) in C1, C2, C5, C10, and C15 treatments are 0.047, 0.094, 0.236, 0.472, and 0.708 g/L day⁻¹, respectively, *Aster sphathulifolius* should be able to grow in roadside or urban soils contaminated by deicing salt because relatively low Cl^- concentrations are present in these field conditions.

The chlorophyll meter is a simple tool used to measure relative chlorophyll content or greenness. It is an efficient indicator of stress in plants (Netto et al., 2005). Changes in chlorophyll content of plants under salt stress depends on stress rate and plant species (Eom et al., 2007; Younes et al., 2016). Nadeem et al. (2006) have reported that salt stress can decrease chlorophyll pigments (a, b, and carotenoids contents) of plant. However, another study has shown that chlorophyll content cannot be used as an indicator of salt tolerance ability (Kanawapee et al., 2012). Although plant species can differ considerably in total amount of chlorophyll content under salt stress, results from this study suggest that chlorophyll content might be considered as an additional trait useful for screening salt tolerance.

Relative abundance of compatible solutes including proline is an important protective factor for plants under salt stress (Norastehnia et al., 2014; Abbas et al., 2014). Salinity stress has caused a significant increase in proline concentration in shoots of plants compared to that in the control (non-stressed seedlings) (Kamawapee et al., 2012). In response to salt stress, proline accumulation in plants has been implicated to play adaptive roles in osmoregulation and salt stress signaling (Szabados and Savaouré, 2009; García-Caparrós et al., 2016). Our results revealed that proline content in plants under salt stress were apparently higher than those in plants under non-salt stress condition. Plants showed significantly higher levels of proline content compared to plants without CaCl_2 stress. These results suggest that proline accumulation might be considered as an additional trait useful for screening plants for salt tolerance ability.

Results of this study showed that increasing CaCl_2 concentrations dramatically reduced survival rate and plant growth of *Aster sphathulifolius*. Biomass was also affected negatively by increasing salt stress of CaCl_2 , with shoots having more reduction in mass than root fresh weight. Calcium chloride salt stress significantly increased chlorophyll *a* and proline contents in leaves, but decreased chlorophyll *b*.

Conclusion

Salt tolerant ornamental groundcover plants can be a very effective strategy to utilizing deicer salt affected soils. The morphological and physiological responses of *Aster sphathulifolius* indicated that the one gram dosage of CaCl_2 for 5 months had no significant difference in survival rates and growth parameters as compared to those of control. However there was a significant negative effect exist by the cumulative dose of deicing CaCl_2 salt (>1 g/L). These results suggest that planting *Aster sphathulifolius* could be highly beneficial for sites with salinity soils such as roadside or urban areas since relatively low Cl^- concentrations are present in these fields. In the future, soil-chloride-plant continuum study warrants to determine the effect of deicers on diverse ecological characteristics of roadside groundcover plants.

Acknowledgements. This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (2015RIC1A2A01052633).

REFERENCE

- [1] Abbas, M.F., Jasim, A.M., Al-Taha, H.A. (2014): Effect of proline and salt stress on organogenesis, growth, proline and carbohydrate content of regenerated plantlets in *Citrus sinensis* (L.) Osbeck cv. local orange. – Advances in Agriculture and Botanic - International Journal of the Bioflux Society 6(2):186-198.
- [2] Acosta-Motos, J.R., Diaz-Vivancos, P., Alvarez, S., Fernandez-Garcia, N., Sanchez-Blanco, M.J., Hernandez, J.A. (2015): Physiological and biochemical mechanisms of the ornamental *Eugenia myrtifolia* L. plants for coping with NaCl stress. – Planta 242(4):829-846.
- [3] Alharby, H.F., Colmer, T.D., Barrett-Lennard, E.G. (2014): Salt accumulation and depletion in the root-zone of the halophyte *Atriplex nummularia* Lindl.: influence of salinity, leaf area and plant water use. – Plant Soil 382: 31-41.
- [4] Ali, A., Iqbal, N., Ali, F., Afzal, B. (2012): *Alternanthera bettzickiana* (Regel) G. Nicholson, a potential halophytic ornamental plant: Growth and physiological adaptations. – Flora 207(4): 318-321.
- [5] Amuthavalli, P., Anbu, D., Sivasankaramoorthy, S. (2012): Effect of calcium chloride on growth and biochemical constituents of cotton (*Gossypium hirsutum* L.) under salt stress. – Int. J. Res. Bot. 2:9-12.
- [6] Baek, M.J., Yoon, T.J., Kim, D.G., Lee, C.Y., Cho, K.J., Bae, Y.J. (2014): Effects of road deicer runoff on benthic macroinvertebrate communities in Korean freshwaters with toxicity tests of calcium chloride (CaCl_2). – Water Air Soil Pollut. 225: 1961-1966.
- [7] Baruah, P., Saikia, R.R., Baruah, P.P., Deka, S. (2014): Effects of crude oil contamination on the chlorophyll content and morpho-anatomy of *Cyperus brevifolius* (Rottb.) Hassk. – Environ. Sci. Pollut. Res. 21: 12530-12538.
- [8] Cao, Z., Zhou, X., Bai, C. (2012): Protection and treatment of landscape plant secondary disasters (deicing agent damage) caused by extreme weather. – Journal of Landscape Research 4:61-64.
- [9] Cassaniti, C., Leonardi, C., Flowers, T. J. (2009): The effect of sodium chloride on ornamental shrubs. – Scientia Horticulturae 122(4): 586-593.
- [10] Chen, L., Ren, J., Shi, H., Chen, X., Zhang, M., Pan, Y., Fan, J., Nevo, E., Sun, D., Fu, J., Peng, J. (2013): Physiological and molecular responses to salt stress in wild emmer and cultivated wheat. – Plant Mol. Biol. Rep. 31: 1212-1219.

- [11] Dang, Y.P., Dalal, R.C., Mayer, D.G., McDonald, M., Routely, R., Schwenke, G.D., Buck, S.R., Daniells, I.G., Singh, D.K., Manning, W., Ferguson, N. (2008): High subsoil chloride concentrations reduce soil water extraction and crop yield on Vertisols in northeastern Australia. – *Crop Pasture Sci.* 59(4): 321-330.
- [12] Devitt, D.A., Wright, L., Landau, F., Apodaca, L. (2014): Deicing salts; Assessing distribution, ion accumulation in plants and the response of plants to different loading rates and salt mixtures. – *Environment and Natural Resources Research* 4(1): 73.
- [13] Eom, S.H., Setter, T.L., Di Tomasso, A., Weston, L. A. (2007): Differential growth response to salt stress among selected ornamentals. – *Journal of Plant Nutrition.* 30: 1109-1126.
- [14] García-Caparrós, P., Llanderal, A., Pestana, M., Correia, P. J., Lao, M. T. (2016): Tolerance mechanisms of three potted ornamental plants grown under moderate salinity. – *Scientia Horticulturae* 201:84-91.
- [15] Guan, B., Yu, N., Chen, X., Xie, W., Lu, Z. (2011): Effects of salt stress and nitrogen application on growth and ion accumulation of *Suaeda salsa* plants. – In: *Remote Sens. Environ. Transport. Engin. 2011 International Conference.* 8268-8272.
- [16] Hammond, H.E., Norcini, J.G., Wilson, S.B., Schoellhorn, R.K., Miller, D.L. (2007): Growth, flowering, and survival of fire wheel *Gaillardia pulchella* Foug. based on seed source and growing location. – *Native Plants J.* 8(1): 25-39.
- [17] Hura, T., Szewczyk-Taranek, B., Hura, K., Nowak, K., Pawlowska, B. (2017): Physiological Responses of *Rosa rubiginosa* to saline environment. – *Water, Air, and Soil Pollution* 228(2): 81.
- [18] Jędrzejuk, A., Łukaszewska, A., Pacholczak, A. (2016): Effects of CaCl₂ solutions to alleviate drought stress effects in potted ornamentals *Salvia splendens* and *Ageratum houstonianum*. – *Acta Agrobotanica* 69(3): 1-11.
- [19] Kanawapee, N., Sanitchon, J., Lontom, W., Threerakulpisut, P. (2012): Evaluation of salt tolerance at the seedling stage in rice genotypes by growth performance, ion accumulation, proline and chlorophyll content. – *Plant Soil* 358: 235-249.
- [20] Lee, J.J., Han, S.G., Kang, C.H., Cho, H.K., Jeong, J.S., Lee, K.B. (2008): Adaptability of saline-tolerant plants selected indoors at reclaimed land. – *Plant Soil* 358: 235-249. *Proc. 2008 Fall Symposium Korean J. Plant Resour.* : 111.
- [21] Mao, H.P., Iwanaga, F., Yamanaka, N., Yamamoto, F. (2008): Growth, photosynthesis, and ion distribution in hydroponically cultured *Populus alba* L. cuttings grown under various salinity concentrations. – *Landscape Ecol. Eng.* 4: 75-82.
- [22] Moore, G.W., Barre, D.A., Owens, M. K. (2010): Changes in soil chloride following shrub removal and subsequent regrowth. – *Geoderma* 158: 148-155.
- [23] Munns, R. (2002): Comparative physiology of salt and water stress. – *Plant Cell Environ.* 25: 239-250.
- [24] Nadeem, S.M., Zahir, Z.A., Naveed, M., Arshad, M., Shahzad, S.M. (2006): Variation in growth and ion uptake of maize due to inoculation with plant growth promoting rhizobacteria under salt stress. – *Soil and Environment* 25: 78-84.
- [25] Netto, A.T., Campostrini, E., Oliveira, J.G., Bressan-Smith, R.E. (2005): Photosynthetic pigments, nitrogen, chlorophyll a fluorescence and SPAD-502 readings in coffee leaves. – *Sci. Hortic.* 104: 199-209.
- [26] Norastehnia, A., Niazazari, M., Sarmad, J., Rassa, M.. (2014): Effects of chloride salinity on non-enzymatic antioxidant activity, proline and malondialdehyde content in three flue-cured cultivars of tobacco. – *J. Plant Develop.* 21: 75-82.
- [27] Ponte, L.F.A., Silva, A.L.C., Carvalho, F.E.L., Maia, J.M., Voigt, E.L. Silveira, J.A.G. (2014): Salt-induced delay in cotyledonary globulin mobilization is abolished by induction of proteases and leaf growth sink strength at late seedling establishment in cashew. – *Journal of Plant Physiology* 171(15): 1362-1371.

- [28] Semiz, G.D., Unlukara, A., Yurtseven, E., Suarez, D.L., Telci, I. (2012): Salinity impact on yield, water use, mineral and essential oil content of fennel (*Foeniculum vulgare* Mill.). – J. Agri. Sci. 18: 177-186.
- [29] Shin, S.P., Yang, C.H., Ha, D.S., Baek, B.K., Cha, J.S. (2010): Study of assumption of functional adequate amount for the additional deicer save. – Korean Society of Road Engineers 12(4): 51-54.
- [30] Szabados, L., Savouré, A. (2009): Proline: a multifunctional amino acid. – Trends Plant Sci. 15(2): 89-87.
- [31] Taffouo, V.D., Sondi, S.E., Meguekam, L.T., Nouck, A.E., Wamba, O.F., Youmbi, E. (2014): Changes in growth and nutrient uptake in response to foliar application of sodium and calcium chloride in cowpea cultivars (*Vigna unguiculata* L. Walp). – Afr. J. Biotechnol. 13(47): 4382-4389.
- [32] Thouvenot, L., Haury, J., Thiébaud, G. (2012): Responses of two invasive macrophyte species to salt. – Hydrobiologia 686: 213-223.
- [33] Viskari, E.L., Kärenlampi, L. (2000): Roadside Scots pine as an indicator of deicing salt use – a comparative study from two consecutive winters. – Water, Air, and Soil Pollution 122: 405-419.
- [34] White, P.J., Broadey, M.R. (2001): Chloride in soils and its uptake and movement within the plant: a review. – Ann. Bot. 88(6): 967-988.
- [35] Whittinghill, L.J., Row, D.B. (2011): Salt tolerance of common green roof and green wall plants. – Urban Ecosyst. 14: 783-794.
- [36] Wu, S., Sun, Y., Niu, G., James, A., Raul, C. (2016): Response of 10 Aster species to saline water irrigation. – HortScience 51(2): 197-201.
- [37] Xu, G., Magen, H., Tarchitzky, J., Kafkafi, U. (2000): Advances in chloride nutrition of plants. – Adv. Agron. 68: 97-150.
- [38] Younes, K.A., Raouf, S.S., Mohammad, S., Morteza, B. (2016): Effect of zinc and bio fertilizers on antioxidant enzymes activity, chlorophyll content, soluble sugars and proline in Triticale under salinity condition.
- [39] Zhang, Y., Li, F., Sun, T., Wang, J. (2012): Effect of deicing salts on urban soils and the health of roadside pines (*Pinus tabulaeformis*) in Northeast China. – Applied Mechanics and Materials 178-181: 353-356.