

A REVIEW OF ASTRAGALUS SPECIES AS FOODSTUFFS, DIETARY SUPPLEMENTS, A TRADITIONAL CHINESE MEDICINE AND A PART OF MODERN PHARMACEUTICAL SCIENCE

SHAHRAJABIAN, M. H.^{1,2#} – SUN, W.^{1,2#} – CHENG, Q.^{1,2*}

¹*Biotechnology Research Institute, Chinese Academy of Agricultural Sciences, Beijing 100081, China*

²*Nitrogen Fixation Laboratory, Qi Institute, Building C4, No. 555, Chuangye County, Jiaxing 314000, Zhejiang, China*

[#]*These authors equally contributed to this paper.*

**Corresponding author
e-mail: chengqi@caas.cn*

(Received 12th Jun 2019; accepted 28th Aug 2019)

Abstract. Astragalus is a common Traditional Chinese Medicinal plant and is a widely used herbal product in China and other countries. Saponins, polysaccharides, amino acids, flavonoids, organic acid, glycosides, alkaloid, and trace elements, are the major classes of chemical compounds occurring in the species of Astragalus genus. In Traditional Chinese Medicine, Astragalus is considered to be effective in the treatment of diabetes, mellitus, nephritis, leukemia, uterine cancer, besides its tonic agent and diuretic effects. Some uses of Astragalus are in the treatment of kidney and urinary problems, digestion, liver problems, female reproductive system problems, muscular, skin problems, cardiovascular and blood related issues, immune and lymphatic system, nervous system, respiratory system, and for some specific diseases. It helps protect the body against various types of stress such as physical and emotional stress. Astragalus root has anti-aging properties, and also help in the prevention of bone loss. *Astragali radix*, the root of *Astragalus membranaceus* Bunge, has been reported to exert hepatoprotective effects, anti-oxidative effects, antiviral activity, anti-oxidative effects, anti-hypertensive effects, and immunostimulant properties; it has also been reported to strengthen superficial resistance, drainage action and new tissue growth. More clinical studies are necessary to discover the effects of Astraglaus.

Keywords: *health benefits, traditional Asian medicine, Yin-Yang balance, Silk Road, herbal medicine*

Introduction

Chinese herbs have been used as traditional medicine immune booster for human being for thousands of years in China (Yin et al., 2009; Shahrajabian et al., 2018). More than 3 million tons of herb medicines were produced in China, and their medicinal parts were consumed in traditional Chinese medicine (TCM) clinic (Soleymani and Shahrajabian, 2012, 2018; Zhang et al., 2016; Ogbaji et al., 2018). In traditional Chinese medicine, some herbals have been used for anti-aging since ancient times (Liu et al., 2017). *Astragalus membranaceus* as one of the most important Qi tonifying adaptogenic herbs in traditional Chinese medicine has a long history of medicinal use (Yang et al., 2010; Zhong et al., 2012; Liu et al., 2017). In traditional Chinese medicine, which laid a lot of emphasis on Qi (vital energy) and Yin-Yang balance (negative and positive equilibrium). Chinese herbs have used as traditional medicine immune booster for human being for thousands of years in China and many parts of Asia (Shahrajabian

et al., 2019a, b, c, 2020; Sun et al., 2020). The goal of this review is survey on some important ancient and modern pharmaceutical sciences of astragalus.

Materials and methods

All relevant papers in the English language of researchers from different countries were collected. The keywords of astragalus, traditional Chinese medicine, traditional Asian medicine, modern pharmaceutical science, health benefits and western medicine were searched in Google Scholar, Scopus, Research Gate and PubMed.

Results and discussion

Astragalus in traditional Chinese medicine and other parts of the world

Astragalus is considered as benefiting Qi and helping to pass water (Li et al., 2011). It has been used as therapy for Wei Zheng, a term for skeletal muscle fatigue and wasting (Zhou and Mei, 2014). The dried root of *A. membranaceus*, first documented in Shennong Bencao Jing (Shennong's Classic of Materia Medica, 200-300 AD), is one of the most popular health promoting herbal medicines commonly used in China for more than 2000 years. In modern Chinese medicine, it is used in Fu zheng therapy as an immune stimulant (Ionkova et al., 1997). Also known as Huang Qi (Chinese), Milk-Vetch (English), Hwanggi (Korean), and Ogi (Japanese) (Chou et al., 2007; Li et al., 2019). It is sold in dietary supplements in tea or capsule form in the USA, and in the tea, beverages, soup, and trail mix (gorp) in Asia (Song et al., 2008; Zhang et al., 2011). Chinese milk vetch (*Astragalus sinicus* L.) is also a traditional leguminous green manure which plays an important role in maintaining paddy soil fertility and in the popularizing of the double-rice farming system in southern China; it is ploughed into soil at full blooming stage and serves as an alternative to chemical nitrogen fertilizer in the region (Zhu et al., 2012). *Astragalus membranaceus* was originally described in the Shennong's Classic of Meteria medica, the earliest complete Pharmacopoeia of China written from Warring States Period to Han Dynasty (Hei et al., 2005; Auyeung et al., 2016). It is valued for its ability to strengthen the primary energy of the body which we know as the immune system, as well as the metabolic, respiratory and eliminative functions (Liu et al., 2017). This fact is being increasingly substantiated by pharmacological studies showing that it can increase telomerase activity, and has antioxidant, anti-inflammatory, immunoregulatory, anticancer, antitumor, antioxidant, hypolipidemic, antihyperglycemic, hepatoprotective, expectorant, immunomodulatory activity, and diuretic effects (Anon, 2003; Ma et al., 2011; Zhao et al., 2011). *Astragalus membranaceus* (Fisch.) Bunge. has been widely used an anti-osteoporosis herb is traditional Chinese medicine for many years (Du et al., 2004; Wong et al., 2007; Xi et al., 2008; Jiao et al., 2014). In Traditional Chinese Medicine, *Astragalus membranaceus* is a major component in a prescription to treat chronic phlegmatic disorders and general gastrointestinal disturbances including stomach ulcer, chronic diarrhea and intestinal inflammation (Kim et al., 2008; Yang et al., 2014). Other researchers have reported the values of Astraglaus s roots in traditional Chinese medicine with the function of strengthening exterior and promoting health for thousands of years (Ma et al., 2017; Zhao et al., 2011). Traditional Chinese herbs are generally applied in the form of multi-herb formulas in medical treatments and as dietary supplements (Takagi and Ishii, 1967; Li et al., 2011). Lu et al. (2013) reported that *Radix Astragali* is the root

of *Astragalus membranaceus* Bunge, and as a famous traditional Chinese medicine (TCM), it has been used to improve muscle wasting-related disorders for a long history. They have also introduced *Astragalus* polysaccharide (APS) as an important bioactive and a therapeutic agent in the management of muscle wasting. *Astragalus trojanus* Stev. is an endemic plant mostly found in eastern and central Anatolia (1300-3500 m), central Aegean region and slopes of Toros mountain (1300-2300 m) in Turkey (Nartop et al., 2015). This genus is widely distributed throughout the temperate and arid regions of the world, and is principally located in Asia (1500 species), North American (500 species) and South American (150 species), and Europe (120 species), but also on mountains in Africa. However, the centre of origin and biodiversity of *Astragalus* plants is Eurasia, especially the mountainous parts of South-Western and South-Central Asia (Lysiuk and Darmohray, 2016). Iran alone, being the richest centre of *Astragalus* habitation, shelters more than 850 species, 527 of which are endemic in the flora of Iran (Ranjbar and Karamian, 2002; Aslanipour et al., 2017; Ghasemian-Yadegari et al., 2017). Aslanipour et al. (2017) stated that the crude drugs prepared from *Astragalus* roots are used for treating some illnesses such as leukemia, respiratory infections and diabetes in Iranian folk medicine. In South Eastern Turkey (Anatolia district), the aqueous extract of the roots of different *Astragalus* spp. are traditionally used against leukemia and for its wound-healing properties (Yesilada et al., 2005; Nalbantasy et al., 2012; Napolitano et al., 2013). In the district of Anatolia, located in South Eastern Turkey, an aqueous extract of the roots of *Astragalus* is traditionally used against leukemia and for its wound healing properties (Bedir et al., 2001). *Astragalus corniculatus* Bieb. (Fabaceae) is distributed in Southeastern Romania, South Ukraine and Moldova (Tutin et al., 1972). *Astragalus Tragacantha* L. (Fabaceae) is a western Mediterranean perennial cushion-like plant species well-adapted to drought that grows even in the trace metal and metalloid polluted soils (Salducci et al., 2019). It has been reported that *Astragalus* is an adaptogen and is usually used in combination with other herbs, such as, ginseng, Echinacea, and glossy privet. *Astragalus* is primarily used in American medicine to potentiate the function of the immune system and in cardiovascular disease. In traditional Chinese medicine, it is used for influenza and the common cold (McKenna et al., 2002). Nishiyama et al. (1995) reported that in traditional oriental medicine, it is conventional to combine different herbs in order to achieve a variety of treatment purposes simultaneously, or to enhance a single effect without causing severe side effects. Erect milkvetch (*Astragalus adsurgens* Pall.), also as a palatable forage, are also widely used in returning farmland to grassland, it has an important role in restoring the degraded ecosystems and could be an effective and applicable to improve soil nutrients and prevent further soil degradation and erosion, because it grows rapidly, and was characterized by barren-tolerance, wide adaptability and strong resistance (Wang and Wang, 2013).

Bioactive phytochemicals, medicinal uses and potential health benefits of Astragalus in traditional and modern medicine industry

Constituents of the dried roots of *Astragalus* spp. *Radix Astragali* provide significant protection against heart, brain, kidney, intestine, liver and lung injury in various models of oxidative stress-related disease (Hong et al., 1992; Shahzad et al., 2016). Zhang et al. (2007) stated that *Astragalus* is an important traditional Chinese medicine (TCM), and now widely used an immune modulator, especially to support immune health for various chronic degenerative diseases. Recently, *Astragalus radix* was proved efficacious to be an adjunctive therapy medicine for cancers (Wang et al., 2003). In the

Bulgarian traditional medicine, *Astragalus glycyphyllos* is used an antihypertensive, diuretic and anti-inflammatory remedy (Nikolov, 2006). Major classes of compounds of *Astragalus* species (Table 1) are polysaccharides, saponins and isoflavonoids, alkaloid, choline, betaine, folic acid, organic acid, various amino acids, mucoitin, gum, cellulose and fourteen trace elements, including selenium, zinc, and iron, which are essential micronutrients for man and animals (Bedir et al., 2000; Block and Mead, 2003; Yin et al., 2006; Lu et al., 2016). *Astragalus* genera are the richest source of cycloartanes, the unique triterpenoids with a characteristics 9,19-cyclopropane (Nartop et al., 2015). According to the systematic review by a Chinese scientists on the chemical constituents of the plants (*genus Astragalus* L.) more than 140 cycloartane-type triterpene glycosides, 60 flavonoids and 18 different polysaccharides have been identified so far (Li et al., 2014). Under high soil moisture and wet conditions, *Astragalus* is susceptible to root rot caused by fungi, which is the main constraint to cultivation. Land must be well drained for *Astragalus*. Loose soil and raised beds can be used to control soil moisture (Shannon et al., 2014). The chemical structures and chain conformations of polysaccharides play a vital role in their biological activities; however, polysaccharides belong to a structurally diverse class of macromolecules (Jin et al., 2014). Cycloastragenol (CA) is the main aglycon of many cycloartane-type glycosides which only found in *Astragalus* genus, extends T cell proliferation by increasing telomerase activity which helps the delay the onset of cellular aging (Valenzuela et al., 2009). *Astragalus* root also contains a series of cycloartane triterpene glycosides, including astragalosides I-VIII, acetylastragaloside, isoastragaloside I and III, astramembrannin II, cycloastragenol, cyclosieversigenis, soyasaponin I, soyasapogenol B, and lupeol (Ko and Chik, 2009). Among these, *Astragalus membranaceus* has a high content of astragaloside IV, which is commonly used as a qualitative marker.

Table 1. *Astragalus membranaceus* main compounds

Compound	Effect
Flavonoids	Plant metabolites with antioxidant effects; give plants yellow color
Isoflavones	Polyphenolic compounds that are classified as phytoestrogens; formononetin is one of the prominent ones
Lectin	Carbohydrate-binding proteins
Polysaccharides	Carbohydrates whose molecules contain linked monosaccharides; starch, inulin, cellulose
Saponins	Amorphous glycosides of terpenes and steroids that can form emulsions and can foam (soap-like); Astragaloside IV is one the most-studied saponins in Huang Qi

On the basis of traditional Chinese medicine view of cancer, causes are endogenous causes and exogenous causes. Endogenous causes are the seven emotional states (anger, grief, fear, worry, over joy, shock and melancholy) can be seen as the way that stress, worry, over work, and emotional grief can suppress the immune system and allow predispositions for cancer growth to take hold. So, while it can seem simplistic to attribute cancer to normal emotions such as sadness, worry, fear, etc., the TCM view is that when these emotions are excessive, prolonged or unresolved, they can cause disease. The concept of Jing in TCM can be likened to the role of genetics in cancer, which is an important factor indeed. Exogenous causes consists of six exogenous causes for all illness, including cancer, are climatic factors of wind, cold, dampness, dryness,

summer heat and fire. And, other miscellaneous causes are environmental causes, dietary causes, and drugs. The TCM concepts of yin/yang balance, the need for calmness of mind, absence of strife, the practice of health promotion through movement, all support modern ideas on the role of psychological, neurological and immunological health in cancer prevention (Shahrajabian et al., 2019d, e, f, g, h). Liu et al. (2011) reported that in many parts of the world, especially in China and Germany, the combined use of herbal treatment and conventional cancer treatment is far more widespread than in America. They have also mentioned that 66.44% of cancer patients in China combined the use of herbal medicine with Western treatment. A prospective, controlled study conducted in Israel (Yaal-Hahoshen et al., 2011) found significant improvement in anemia and neutropenia in breast cancer patients who were given an herbal mixture containing Huang Qi and other Chinese herbs. Rios and Waterman (1997) reported that cycloastragenol (CAG) is a secondary metabolite isolated from *Radix Astragali*, present in all known *Astragalus spp.*, CAG (9,19-cycloanostane-3,6,16,25-tetrol,20,24-epoxy-(3 β , 6 α ,16 β ,20R,24S); is both a triterpene aglycone and the most common genuine aglycone in the bioactive triterpenoid saponins called astragalosides. *Astragalus polysaccharides* (APS) are one of the main efficacious principles of *Radix Astragali* (*Astragalus membranaceus*), which is reported to have anti-oxidant, anti-diabetic, anti-hypertensive, and immunomodulatory activities (Wu and Chen, 2004; Wu et al., 2005). It has been noted that the crude polysaccharide extract of *A. membranaceus* was mainly composed of carbohydrates with small amount of proteins (Cho and Leung, 2007). It has been demonstrated that the main components of the ethylacetate extract of *Astragalus* were isoflavonoids such as calycosin-7-O- β -D-glycoside, formononetin-7-O- β -D-glycoside and (6R, 10R)-9,10-dimethoxypterocarpan-3-O- β -D-glycoside, and these glycosides and other their aglycones were proved to exhibit strong antioxidant activity (Zhang et al., 2007). Li et al. (2010) also mentioned that the dried root of *Astragalus* contains 2',4'-dihydroxy-5,6-dimethoxyisoflavone, kumatakenin, choline, betaine, polysaccharides, saponins, glucuronic acid, sucrose, amino acids, traces of folic acid and astraisoflavanin. So many other scientists also revealed that *Astragalus membranaceus* has a notable functional role in various pharmacopoeias as a herbal immunomodulator and an anti-diabetic drug (Wei et al., 2011; Agyeman et al., 2013). Its roots have been used in many state-approved Chinese Herbal formulas for the treatment of diabetes (Jai et al., 2003; Wei et al., 2011). Some experiments have showed that *Astragalus* exhibits immunomodulating and immunorestorative effects both in vitro and in vivo (Guo et al., 2005), and have shown preliminary promise against the experimental coccidial infection when used in conjunction with vaccine (Cho and Leung, 2007). Song et al. (2017) reported that *Astragalus* extract mixture HT042 is a combination of three standardized herbal extracts from *Astragalus membranaceus* root, *Elutherococcus senticosus* stem, and *Phlomis umbrosa* root, and it has been developed to promote height growth in children with short stature. Sun et al. (2012) revealed that *Astragalus membranaceus* is a popular traditional Chinese medicine, commonly used in Chinese herb prescription to treat liver disease, and the extract prepared from the roots of *Astragalus membranaceus* and *Paeonia lactiflora* demonstrated better hepatoprotective activity than the herbs used individually. Ko and Chik (2009) demonstrated that root extract of *Astragalus membranaceus* administrated orally and locally can protect rats against hapten-induced colitis through attenuation of TNF- α and IL-1 β and up-regulating of IL-10. Shen et al. (2008) indicated that *Astragalus membranaceus* has a potential role in treating allergic

asthma. Zhang et al. (2009) *Astragalus membranaceus* and its effective components are effective in reducing fasting blood glucose and albuminuria levels, in reversing the glomerular hyperfiltration state, and in ameliorating the pathological changes of early diabetic nephropathy in rat models. Ko and Chik (2009) demonstrates that both oral and locally administered *Astragalus membranaceus* possess protective effects against experimental colitis through differential modulation of colonic cytokines. Yang et al. (2013) found that *Astragalus membranaceus* polysaccharide (AMP) has antitumor activity in vivo at least partly via improving immune responses of host organism, and seems to be safe and effective for the use of anti-tumor therapy. Lv et al. (2017) suggest that *Astragalus polysaccharide* (APS) which is a bioactive extract of *Astragalus membranaceus* may represent a natural therapeutic approach for treating inflammatory bowel disease, such as ulcerative colitis. Yan et al. (2010) found that administration of *Astragalus mongholicus* polysaccharides could significantly increase serum and liver antioxidant enzyme activities in mice and decrease peroxidative lipid levels. Jalsrai et al. (2010) found that doses of *Astragalus mongholicus* extract which did not interfere with locomotor activity and situational anxiety appear to be useful in the treatment of convulsive disorders. Kim et al. (2016) highlighted the ability of *Astragalus membranaceus* to facilitate sperm development and semen quality. Tian et al. (2016) reported that *Astragalus* may be beneficial as an adjuvant therapy in the treatment of type 2 diabetes. Zhou et al. (2018) demonstrated that the extract from *Astragalus membranaceus* with water extraction-ethanol supernatant method inhibit cell growth and induce apoptosis in cultured breast cancer cells. This effect astragalus extract to suppress breast cancer cells growth was associated with its ability to inhibit PI3K/Akt/mTOR activity. Maresca et al. (2017) concluded that the 50% hydroalcoholic extract of *Astragali radix* is a valuable candidate for the adjuvant treatment of articular disease. Liu et al. (2017) concluded that the appropriate dose of *Astragalus* depends on several factors, such as the user's age, health status, and several other conditions. They have also found that natural products are not always necessarily safe, and dosages can be important. Sheng et al. (2005) found that one of the most important biological role of saponins is modulating the cellular oxidant antioxidant balance.

Conclusions

Traditional Chinese medicinal materials have been used for thousands of years and are believed to be abundant, safe, and inexpensive. *Astragalus* has been used in Chinese traditional medicine as an immunity booster for almost 2000 years. The genus *Astragalus* is a member of the Fabaceae of Legume family and it is native to northern China and Mongolia. It is widely distributed in China, Siberia, and northern Korea. Some actions of *Astragalus membranaceus* are anti-viral, anti-bacterial, immune system enhancing, immune stimulant, anti-infective some viruses, adaptogen, cardio-tonic, diuretic, hypotensive, anti-oxidant, immunomodulator, hypoglycaemic, circulatory stimulant, vasodilator, anti-fatigue, anti-cancer and hepatoprotective. Chemical constituents are polysaccharides, triterpenoid saponins (*Astragalosides*), flavonoids, choline, phytosterols, volatile oils, amino acids (*Asparagine*, *Gamma-aminobutyric acid*, *Canavanine*), aglycones, coumarins, *astrapterocarpan*, *betaine*, calcium, copper, isoflavonoids, rich in potassium and magnesium. *Astragalus membranaceus* classically prescribed in TCM in combination with other Chinese medicinal herbs as a dried root, powdered or as a decoction, with the combination depending on the desired therapeutic

effect and the specific TCM diagnosis. Polysaccharides in *Astragalus* intensify phagocytosis in reticuloendothelial systems, stimulate pituitary-adrenal cortical activity, and restore depleted red blood cell formations in bone marrow. *Astragalus* uses are in kidneys problems, strengthens the kidneys, incontinence and frequent urination, urinary tract infection, gastric ulcers, decreased appetite, chronic diarrhea, strengthens the spleen, poor digestion, liver problems, viral hepatitis, strengthen the liver, postpartum fever, uterine bleeding, topical adjuvant therapy for chronic viral cervicitis, vaginitis, edema, lupus, rheumatoid arthritis, myasthenia gravis, strengthens and builds bone marrow, excessive sweating, night sweats, slow healing wounds, increasing white blood cell count, leukopenia, ischemic heart disease, angina pectoris, recovery from severe loss of blood, diabetes, anemia, high blood pressure, heart palpitations, congestive heart failure, strengthens the blood, increases interferon production, impaired immunity, chronic viral infections, general debility, increases energy, HIV/AIDS, cancer, myeloc encephalomyelitis (chronic fatigue syndrome), improves sleep quality, upper respiratory infection, common cold, chemotherapy, radiation therapy, flu, combats coxsackie B myocarditis, ameliorate side effects of drugs, appropriate herb for weak and elderly. Isoflavonoids, such as calycosin-7-O- β -D-glucoside, ononin, astraisoflavan-7-O- β -D-glucoside, calycosin and formononetin, are principle bioactive compounds found in *Radix astragali*-based drugs or foods. In summary, *Astragalus* is an ancient herb for modern medicine which can promote good health and as drugs to treat diseases. Further clinical researches are necessary to uncover various substances and their effects on *astragalus* which contribute to public health.

REFERENCES

- [1] Agyemang, K., Han, L., Liu, E., Zhang, Y., Wang, T., Gao, X. (2013): Recent advances in *Astragalus membranaceus* anti-diabetic research: pharmacological effects of its phytochemical constituents. – Evid-Based Compl Alt Med Article ID 654643.
- [2] Anon (2003): *Astragalus membranaceus*. Monograph. – Altern Med Rev 8: 72-77.
- [3] Aslanipour, B., Gulcemal, D., Nalbantsoy, A., Yusufoglu, H., Bedir, E. (2017): Cycloartane-type glycosides from *Astragalus brachycalyx* FISCHER and their effects on cytokine release and hemolysis. – Phytochem Lett 21: 66-73.
- [4] Auyeung, K. K., Han, Q. B., Ko, J. K. (2016): *Astragalus membranaceus*: a review of its protection against inflammation and gastrointestinal cancers. – Am J Chin Med. 44: 1-22.
- [5] Bedir, E., Calis, I., Piacente, S., Pizza, C., Khan, I. A. (2000): New flavonol glycoside from the aerial parts of *Astragalus vulneraria*. – Chem. Pharm. Bull. 48(12): 1994-1995.
- [6] Bedir, E., Calis, I., Dunbar, C., Sharan, R., Buolamwini, J. K., Khan, I. A. (2001): Two novel cycloartane-type triterpene glycosides from the roots of *Astragalus prusianus*. – Tetrahedron 57: 5961-5966.
- [7] Block, K. I., Mead, M. N. (2003): Immune system effects of Echinacea, ginseng, and astragalus: a review. – Integr Cancer Ther. 2(3): 247-267.
- [8] Cho, W. C., Leung, K. N. (2007): In vitro and in vivo immunomodulating and immunorestorative effects of *Astragalus membranaceus*. – J Ethnopharmacol 113: 132-141.
- [9] Chou, M.-X., Wei, X.-Y., Chen, D.-S., Zhou, J.-C. (2007): A novel nodule-enhanced gene encoding a putative universal stress protein from *Astragalus sinicus*. – J. Plant Physiol 164: 764-772.
- [10] Du, S. H., Meng, Z. (2004): The treatment of *Astragalus membranaceus* (Fisch.) Bunge. for nephroasthenia syndrome and headache. – J Trad. Chi. Med. 45: 250.

- [11] Ghasemian-Yadegari, J., Nazemiyeh, H., Hamedeyazadan, S., Fathiazad, F. (2017): Secondary metabolites from the roots of *Astragalus maximus*. – Res J Pharmacog 4(2): 31-38.
- [12] Guo, F. C., Kwakkel, R. P., Williams, C. B., Suo, X., Li, W. K., Verstegen, M. W. (2005): Coccidiosis immunization: effects of mushroom and herb polysaccharides on immune responses of chickens infected with *Eimeria tenella*. – Avian Dis 49: 70-73.
- [13] Hei, Z. Q., Huang, H. Q., Zhang, J. J., Chen, B. X., Li, X. Y. (2005): Protective effect of *Astragalus membranaceus* on intestinal mucosa reperfusion injury after hemorrhagic shock in rats. – World J Gastroenterol 11: 4986-4991.
- [14] Hong, C. Y., Ku, J., Wu, P. (1992): *Astragalus membranaceus* stimulates human sperm motility in vitro. Am J Chin Med. 20: 289-294. – Trop J Pharm Res 15(9): 1897-1901.
- [15] Ionkova, I., Kartnig, T., Alfermann, W. (1997): Cycloartane saponin production in hairy root cultures *Astragalus Mongholicus*. – Phytochemistry 45(8): 1597-1600.
- [16] Jalsrai, A., Grecksch, G., Becker, A. (2010): Evaluation of the effects of *Astragalus mongholicus* Bunge saponin extract on central nervous system functions. – J Ethnopharmacol 131: 544-549.
- [17] Jiao, J., Wei, F.-Y., Gai, Q.-Y., Wang, W., Luo, M., Fu, Y.-J., Ma, W. (2003): A pilot-scale homogenization-assisted negative pressure cavitation extraction of *Astragalus* polysaccharides. – Int J Biol Macromol 67: 189-194.
- [18] Jin, M., Zhao, K., Huang, Q., Shang, P. (2014): Structural features and biological activities of the polysaccharides from *Astragalus membranaceus*. – Int J Biol Macromol 64: 257-266.
- [19] Kim, S. W., Fan, M. Z., Applegate, T. J. (2008): Nonruminant nutrition symposium on natural probiotics for health of young animals and poultry: mechanisms and application. – J. Anim. Sci 86: E138-E139.
- [20] Kim, W., Kim, D. R., Chang, M. S., Park, S. K. (2016): *Astragalus membranaceus* augment sperm parameters in male mice associated with cAMP-responsive element modulator and activator of CREM in testis. – J Tradit Complement Med 6: 294-298.
- [21] Ko, J. K., Chik, C. W. (2009): The protective action of *radix Astragalus membranaceus* against hapten-induced colitis through modulation of cytokines. – Cytokine 47: 85-90.
- [22] Li, C.-Y., Chen, H.-Y., Liu, W.-P., Rui, W. (2019): Multi-fingerprint profiling combined with chemometric methods for investigating the quality of *Astragalus* polysaccharides. – Int. J. Biol. Macromol. 123: 766-774.
- [23] Li, H., Nie, D., Wang, C., Fang, J., Li, D. (2016): Anti-osteoporosis activity of *Astragalus membranaceus*Bunge extract in experimental rats. – Trop J Pharm Res 15(9): 1897-1901.
- [24] Li, M., Xu, Y., Yang, W., Li, J., Xu, X., Zhang, X., Chen, F., Li, D. (2011): In vitro synergistic anti-oxidant activities of solvent-extracted fractions from *Astragalus membranaceus* and *Glycyrrhiza uralensis*. – LWT-Food Sci Technol 44: 1745-1751.
- [25] Li, R., Chen, W.-C., Wang, W.-P., Tian, W.-Y., Zhang, X.-G. (2010): Antioxidant activity of *Astragalus* polysaccharides and antitumour activity of the polysaccharides and siRNA. – Carbohydr Polym 82: 240-244.
- [26] Li, X., Qu, L., Dong, Y., Han, L., Liu, E., Fang, S., Zhang, Y., Wang, T. A. (2014): A review of recent research progress on the *Astragalus* genus. – Molecules 19(11): 18850-18880.
- [27] Liu, J., Li, X., Liu, J., Ma, L., Li, X., Fonnebo, V. (2011): Traditional Chinese medicine in cancer care: a review of case reports published in Chinese literature. – Forsch Komplementmed 18(5): 257-263.
- [28] Liu, P., Zhao, H., Luo, Y. (2017): Anti-aging implications of *Astragalus membranaceus* (Huangqi): A well-known Chinese tonic. – Aging Dis 8(6): 868-886.
- [29] Lu, L., Wang, D.-T., Shi, Y., Yin, Y., Wei, L.-B., Zou, Y.-C., Huang, B., Zhao, Y., Wang, M., Wan, H., Li, C. J., Diao, J.-X. (2013): *Astragalus* polysaccharide improves muscle atrophy from dexamethasone-and peroxide-induced injury in vitro. – Int. J. Biol. Macromol 61: 7-16.

- [30] Lu, L., Huang, Y.-F., Chen, D.-X., Wang, M., Zou, Y.-C., Wan, H., Wei, L.-B. (2016): *Astragalus* polysaccharides decrease muscle wasting through Akt/mTOR, ubiquitin proteasome and autophagy signalling in 5/6 nephrectomised rats. – *J Ethnopharmacol* 186: 125-135.
- [31] Lv, J., Zhang, Y., Tian, Z., Liu, F., Shi, Y., Liu, Y., Xia, P. (2017): *Astragalus* polysaccharides protect against dextran sulphate sodium-induced colitis by inhibiting NF- κ B activation. – *Int J Biol Macromol* 98: 723-729.
- [32] Lysiuk, R., Darmohray, R. (2016): Pharmacology and ethnomedicine of the genus *Astragalus*. – *Int. J. Pharmacol. Phytochem* 3: 46-53.
- [33] Ma, J., Qiao, Z., Xiang, X. (2011): Aqueous extract of *Astragalus mongholicus* ameliorates high cholesterol diet induced oxidative injury in experimental rat models. – *J. Med Plant Res* 5: 855-858.
- [34] Ma, Y., Liu, C., Qu, D., Chen, Y., Huang, M., Liu, Y. (2017): Antibacterial evaluation of silver nanoparticles synthesized polysaccharides from *Astragalus membranaceus* roots. – *Biomed Pharmacother* 89: 351-357.
- [35] Maresca, M., Micheli, L., Cinci, L., Bilia, A. R., Ghelardin, C., Di, L., Mannelli, C. (2017): Pain relieving and protective effects of *Astragalus hydroalcoholic* extract in rat arthritis models. – *J Pharm Pharmacol* 1858-1870.
- [36] McKenna, D. J., Hughes, K., Jones, K. (2002): *Astragalus*. – *Altern Ther Health Med* 8(6): 34-40.
- [37] Nalbantsoy, A., Nesil, T., Yilmaz-Dilsiz, O., Aksu, G., Khan, S., Bedir, E. (2012): Evaluation of the immunomodulatory properties in mice and in vitro anti-inflammatory activity of cycloartane type saponins from *Astragalus species*. – *J. Ethnopharmacol* 139: 574-581.
- [38] Napolitano, A., Akay, S., Mari, A., Bedir, E., Pizza, C., Piacente, S. (2013): An analytical approach based on ESI-MS, LC-MS and PCA for the quali-quantitative analysis of cycloartane derivatives in *Astragalus spp.* – *J Pharmaceut Biomed* 85: 46-54.
- [39] Nartop, P., Gurel, A., Akgun, I. H., Bedir, E. (2015): Astragaloside IV and cycloastragenol production capacity of *Astragalus trojanus Calli.* – *Rec. Nat. Prod.* 9(1): 49-61.
- [40] Nikolov, S. (2006): *Astragalus glycyphyllos*. – In: Nikolov, S. (ed.), *Specializirana enciklopedia na lechebnite rastenija*. BAS, Sofia, pp. 317-318.
- [41] Nishiyama, N., Wang, Y. L., Saito, H. (1995): Beneficial effects of S-113m, a novel herbal prescription, on learning impairment model in mice. – *Biol Pharm Bull* 18: 1498-1503.
- [42] Ogbaji, P. O., Li, J., Xue, X., Shahrajabian, M. H., Egrinya, E. A. (2018): Impact of bio-fertilizer or nutrient solution on Spinach (*Spinacea Oleracea*) growth and yield in some province soils of P. R. China. – *Cercetari Agronomice in Moldova* 2(174): 43-52.
- [43] Ranjbar, M., Karamian, R. (2002): *Astragalus* sect. *Astragalus* (Fabaceae) in Iran: complementary notes with a key to the species. – *Nordic. J. Bot.* 2: 177-181.
- [44] Rios, J. L., Waterman, P. G. (1997): A review of the pharmacology and toxicology of *Astragalus*. – *Phytother. Res.* 11: 411-418.
- [45] Salducci, M.-D., Folzer, H., Issartel, J., Rabier, J., Masotti, V., Prudent, P., Affre, L., Hardion, L., Tatoni, T., Laffont-Schwob, I. (2019): How can a rare protected plant cope with the metal and metalloid soil pollution resulting from part industrial activities? Phytometabolites, antioxidant activities and root symbiosis involved in the metal tolerance of *Astragalus tragacantha*. – *Chemosphere* 217: 887-896.
- [46] Shahrajabian, M. H., Sun, W., Cheng, Q. (2018): A review of Goji berry (*Lycium barbarum*) in traditional Chinese medicine as a promising organic superfood and superfruit in modern industry. – *Academia Journal of Medicinal Plants* 6(12): 437-445.
- [47] Shahrajabian, M. H., Sun, W., Cheng, Q. (2019a): The influence of traditional Iranian and Chinese medicine on western and Islamic countries. – *Asian J. Med. Bio. Res* 5(2): 94-99.

- [48] Shahrajabian, M. H., Sun, W., Cheng, Q. (2019b): Modern pharmacological actions of Longan fruits and their usages in traditional herbals remedies. – J. Med. Plants. Stud 7(4): 179-185.
- [49] Shahrajabian, M. H., Khoshkharam, M., Zandi, P., Sun, W., Cheng, Q. (2019c): Jujube, a super-fruit in traditional Chinese medicine, heading for modern pharmacological science. – J. Med. Plants. Stud 7(4): 173-178.
- [50] Shahrajabian, M. H., Sun, W., Cheng, Q. (2019d): The power of natural Chinese medicine, ginger and ginseng root in an organic life. – Middle East J. Sci. Res 27(1): 64-71.
- [51] Shahrajabian, M. H., Sun, W., Cheng, Q. (2019e): Clinical aspects and health benefits of ginger (*Zingiber officinale*) in both traditional Chinese medicine and modern industry. – Acta Agriculturae Scandinavica, Section B-Soil & Plant Science 1-11.
- [52] Shahrajabian, M. H., Sun, W., Cheng, Q. (2019f): Traditional Chinese medicine and agriculture; organic life and sustainability for future. – GSC Biol. Pharm. Sci 7(1): 091-095.
- [53] Shahrajabian, M. H., Sun, W., Cheng, Q. (2019g): A review of ginseng species in different regions as a multipurpose herb in traditional Chinese medicine, modern herbology and pharmacological science. – J Med. Plants Res 13(10): 213-226.
- [54] Shahrajabian, M. H., Khoshkharam, M., Sun, W., Cheng, Q. (2019h): The effect of pretreatment factors on seed germination and seedling growth of anise (*Pimpinella anisum* L.). – Middle East J. Sci 5(1): 86-93.
- [55] Shahrajabian, M. H., Khoshkharam, M., Zandi, P., Sun, W., Cheng, Q. (2020): The influence of temperatures on germination and seedling growth of pyrethrum (*Tanacetum cinerariifolium*) under drought stress. – International Journal of Advanced Biological and Biomedical Research 8(1): 29-39 (in press).
- [56] Shahzad, M., Shabbir, A., Wojcikowski, K., Wohlmuth, H., Gobe, G. C. (2016): The Antioxidant Effects of Radix Astragali (*Astragalus membranaceus* and Related Species) in Protecting Tissues from Injury and Disease. – Curr Drug Targets 17: 1331-1340.
- [57] Shannon, D. A., Wang, M., Kemppainen, B., Mitchell, C. C., Salmasi, S. Z. (2014): Adaptation of *Astragalus membranaceus* varieties to Southeastern United States: growth, root development and Astragaloside IV content. – J. Med. Plants. Stud 2(3): 80-91.
- [58] Shen, H.-H., Wang, K., Li, W., Ying, Y.-H., Gao, G. X., Li, X.-B., Huang, H.-Q. (2008): *Astragalus membranaceus* prevents airway hyperreactivity in mice related to Th2 response inhibition. – J Ethnopharmacol 116: 363-369.
- [59] Sheng, B. W., Cheng, H. F., Zhao, J., He, D. L., Nan, X. Y. (2005): *Astragalus membranaceus* reduces free radical-mediated injury to renal tubules in rabbits receiving high-energy shock waves. – Chinese Medic J (England) 118: 43-49.
- [60] Soleymani, A., Shahrajabian, M. H. (2012): Response of different cultivars of fennel (*Foeniculum vulgare*) to irrigation and planting dates in Isfahan, Iran. – Res Crops 13(2): 656-660.
- [61] Soleymani, A., Shahrajabian, M. H. (2018): Changes in germination and seedling growth of different cultivars of cumin to drought stress. – Cercetari Agronomice in Moldova 1(173): 91-100.
- [62] Song, J., Lee, D., Min, B., Bae, J.-S., Chang, G. Y., Kim, H. (2017): Safety evaluation of *Astragalus* extract mixture HT042 and its constituent herbs in Sprague-Dawley rats. – Phytomedicine 32: 59-67.
- [63] Song, J. Z., Yiu, H., Qiao, C. F., Han, Q. B., Xu, H. X. (2008): Chemical comparison and classification of Radix Astragali by determination of isoflavonoids and astragalosides. – J. Pharm. Biomed. Anal. 47: 399-406.
- [64] Sun, W.-Y., Wang, L., Liu, H., Li, X., Wei, W. (2012): A standardized extract from *Paeonia lactiflora* and *Astragalus membranaceus* attenuates liver fibrosis induced by porcine serum in rats. – Int J Mol Med 29: 491-498.

- [65] Sun, W., Shahrajabian, M. H., Qiman, H. (2020): Soybean seeds treated with single walled carbon nanotubes (SwCNTs) showed enhanced drought tolerance during germination. – *International Journal of Advanced Biological and Biomedical Research* 8(1): 9-16 (in press).
- [66] Takagi, K., Ishii, Y. (1967): Peptic ulcer inhibiting properties of a new fraction from licorice root (FM100). 1. Experimental peptic ulcer and general pharmacology. – *Arzneimittel-Forschung* 17(12): 1544-1547.
- [67] Tian, H., Lu, J., He, H., Zhang, L., Dong, Y., Yao, H., Feng, W., Wang, S. (2016): The effect of *Astragalus* as an adjuvant treatment in type 2 diabetes mellitus: A (preliminary) meta-analysis. – *J Ethnopharmacol* 191: 206-215.
- [68] Toshkova, R. A., Krasteva, I. N., Wesselinova, D. W., Nikolov, S. D. (2007): Influence of purified saponin mixture from *Astragalus corniculatus* Bieb. On phagocytic cells in Graffi-tumor bearing hamsters. – *J Ethnopharmacol* 109: 394-399.
- [69] Tutin, T. G., Heywood, V. H., Burges, N. A., Mooze, D. M., Valeutine, D. H., Walters, S. M., Webb, D. A. (1972): *Flora Europea*. Vol. 2. – Cambridge University Press, Cambridge, pp. 108-124.
- [70] Valenzuela, H. F., Fuller, T., Edwards, J., Finger, D., Molgora, B. (2009): Cycloastragenol extends T cell proliferation by increasing telomerase activity. – *J. Immunol* 182: 90-30.
- [71] Wang, Y. P., Li, X. Y., Song, C. Q., Hu, Z. B. (2002): Effect of astragaloside IV on T, B lymphocyte proliferation and peritoneal macrophage function in mice. – *Acta Pharmacol. Sin.* 23: 263.
- [72] Wang, Z.-B., Wang, Q.-Y. (2013): Cultivating erect milkvetch (*Astragalus adsurgens* Pall.) (Leguminosae) improved soil properties in loess hilly and gullies in China. – *J. Integr. Agric* 12(9): 1652-1658.
- [73] Wei, D. X., Yu, N. Z., Ya, O. Z. (2011): Traditional Chinese medicines in treatment of patients with type 2 diabetes mellitus. – Evidence-Based Complementary and Alternative Medicine Article ID 726723.
- [74] Wong, R. W., Rabie, B., Benedus, M., Hagg, U. (2007): The effects of *Rhizoma Curculiginis* and *Astragalus membranaceus* (Fisch.) Bunge. extracts on bones. – *J Chi. Med.* 2: 13-14.
- [75] Wu, F., Chen, X. (2004): A review of pharmacological study on *Astragalus membranaceus* (Fisch) Bge. – *Zhong Yao Cai* 27: 232-234.
- [76] Wu, Y., Ou-Yang, J.-P., Wu, K., Wang, Y., Zhou, Y.-F., Wen, C.-Y. (2005): Hypoglycemic effect of *Astragalus* polysaccharide and its effect on PTP1B¹. – *Acta Pharm Sin* 26(3): 345-352.
- [77] Xi, M., Hai, C., Tang, H., Fang, K., Liang, X. (2008): Antioxidant and antiglycation properties of total saponins extracted from traditional Chinese medicine used to treat diabetes mellitus. – *Phytother. Res.* 22(2): 228-237.
- [78] Yaal-Hahoshen, N., Maimon, Y., Siegelmann-Danieli, N., Lev-Ari, S., Ron, I. G., Sperber, F., Samuels, N., Shoham, J., Merimsky, O. (2011): A prospective, controlled study of the botanical compound mixture LCS101 for chemotherapy-induced haematological complications in breast cancer. – *Oncologist* 16(9): 1197-1201.
- [79] Yan, H., Xie, Y., Sun, S., Sun, X., Ren, F., Shi, Wang, S., Zhang, W., Li, X., Zhang, J. (2010): Chemical analysis of *Astragalus mongholicus* polysaccharides and antioxidant activity of the polysaccharides. – *Carbohydrate Polymers* 82: 636-640.
- [80] Yang, B., Xiao, B., Sun, T. (2013): Antitumor and immunomodulatory activity of *Astragalus membranaceus* polysaccharides in H22 tumor-bearing mice. – *Int J. Biol Macromol* 62: 287-290.
- [81] Yang, M., Lin, H.-B., Gong, S., Chen, P.-Y., Geng, L.-L., Zeng, Y.-M., Li, D.-Y. (2014): Effect of *Astragalus* polysaccharides on expression of TNF- α , IL-1 β and NFATc4 in a rat model of experimental colitis. – *Cytokine* 70: 81-86.

- [82] Yang, X., Huang, S., Chen, J., Song, N., Wang, L., Zhang, Z., Deng, G., Zheng, H., Zhu, X.-Q., Lu, F. (2010): Evaluation of the adjuvant properties of *Astragalus membranaceus* and *Scutellaria baicalensis* GEORGI in the immune protection induced by UV-attenuated *Toxoplasma gondii* in mouse models. – Vaccine 28: 737-743.
- [83] Yesilada, E., Bedir, E., Calis, I., Takaishi, Y., Ohmoto, Y. (2005): Effects of triterpene saponins from *Astragalus* species on in vitro cytokine release. – J. Ethnopharmacol 96: 71-77.
- [84] Yin, G., Jeney, G., Racz, T., Xu, P., Jun, X., Jeney, Z. (2006): Effect of two Chinese herbs (*Astragalus radix* and *Scutellaria radix*) on non-specific immune response of tilapia *Oreochromis niloticus*. – Aquaculture 253: 39-47.
- [85] Yin, G., Ardo, L., Thompson, K. D., Adams, A., Jeney, Z., Jeney, G. (2009): Chinese herbs (*Astragalus radix* and *Ganoderma lucidum*) enhance immune response of carp, *Cyprinus carpio*, and protection against *Aeromonas hydrophila*. – Fish Shellfish Immunol 26: 140-145.
- [86] Zhang, J., Xie, X., Li, C., Fun, P. (2009): Systematic review of the renal protective effect of *Astragalus membranaceus* (root) on diabetic nephropathy in animal models. – J. Ethnopharmacol 126: 189-196.
- [87] Zhang, L. J., Liu, H. K., Hsiao, P. C., Kuo, L. M., Lee, I. J., Wu, T. S., Chiou, W. F., Kuo, Y. H. (2011): New isoflavonoid glycosides and related constituents from *Astragali Radix* (*Astragalus membranaceus*) and their inhibitory activity on nitric oxide production. – J. Agric. Food Chem. 59: 1131-1137.
- [88] Zhang, X., Sun, Y. G., Cheng, M. C., Wang, Y. Q., Xiao, H. B., Liang, X. M. (2007): Simultaneous quantification of three isoflavonoid glycosides in rabbit plasma after oral administration of *Astragalus mongholicus* extract by high-performance liquid chromatography coupled with electrospray. – Anal Chim Acta 602: 252-258.
- [89] Zhang, Y., Li, X., Ruan, J., Wang, T., Dong, Y., Hao, J., Liu, E., Han, L., Gao, X., Wang, T. (2016): Oleanane type saponins from the stems of *Astragalus membranaceus* (Fisch.) Bge. Var. *mongholicus* (Bge.) Hsiao. – Fitoterapia 109: 99-105.
- [90] Zhao, L. H., Ma, Z. X., Zhu, J., Yu, X. H., Weng, D. P. (2011): Characterization of polysaccharide from *Astragalu radix* as the macrophage stimulator. – Cell Immunol 271(2): 329-334.
- [91] Zhong, R. Z., Yu, M., Liu, H. W., Sun, H. X., Cao, Y., Zhou, D. W. (2012): Effects of dietary *Astragalus* polysaccharide and *Astragalus membranaceus* root supplementation on growth performance, rumen fermentation, immune responses, and antioxidant status of lambs. – Anim Feed Sci Tech 174: 60-67.
- [92] Zhou, K., Mei, X. Y. (2014): Study on preventive and therapeutic effects of *Astagali radix* on denervated tibial muscle atrophy in rats. – Zhongguo Zhong Yao Za Zhi 39: 1083-1087.
- [93] Zhou, R., Chen, H., Chen, J., Chen, X., Wen, Y., Xu, L. (2018): Extract from *Astragalus membranaceus* inhibit breast cancer cells proliferation via PI3K/AKT/mTOR signalling pathway. – Complement Altern Med 18: 83.
- [94] Zhu, B., Yi, L.-X., Hu, Y.-G., Zeng, Z.-H., Tang, H.-M., Yang, G.-L., Xiao, X.-P. (2012): Effects of Chinese Milk Vetch (*Astragalus sinicus* L.) residue incorporation on CH₄ and N₂O emission from a double-rice paddy soil. – J. Integr. Agric 11(9): 1537-1544.