

RESPONSES OF SULFUR AND PHOSPHORUS DOSES ON THE YIELD AND QUALITY OF FENUGREEK (*Trigonella foenum-graecum* L.)

ÖZYAZICI, G.

Department of Field Crops, Faculty of Agriculture, Siirt University, Siirt, Turkey
e-mail: gulenozyazici@siirt.edu.tr; phone: +90-533-221-8208

(Received 24th Feb 2020; accepted 9th Jul 2020)

Abstract. This study was conducted to determine the effects of sulfur and phosphorus doses on the yield and quality of fenugreek in Siirt, Turkey, in the 2016-2017 and 2017-2018 growing seasons. Field trials were designed in a factorial complete block design with three replications at the experimental fields of the Agricultural Faculty of Siirt University. In the study, plant height (cm), first pod height (cm), the number of pods (pod plant⁻¹), the number of seeds in the pod (seed pod⁻¹), pod length (cm), thousand-seed weight (g), seed yield (kg ha⁻¹), protein content (%) and trigonelline content (%) were determined. All of the growth and yield parameters, except for plant height and pod length, were significantly affected by sulfur fertilization. All the parameters were affected by phosphorus fertilization. The highest seed yield (2224 kg ha⁻¹) and trigonelline content (1.26%) were obtained under 30 kg S ha⁻¹ and 90 kg P ha⁻¹ applications.

Keywords: *Trigonella foenum-graecum*, fertilizer, medicinal plants, seed yield, trigonelline

Introduction

Nowadays, due to the many side effects of synthetic and chemical drugs, trending towards alternative medicine or complementary medicine is gradually increasing. Medicinal and aromatic plants used for this purpose can be found in Turkey among many other countries, mainly by gathering from the natural flora. Therefore, medicinal plants of the desired quantity and quality cannot be obtained. This situation brings up the production of medicinal and aromatic plants with increasing the usage area.

Fenugreek (*Trigonella foenum-graecum* L.) which is grown for its seeds, fresh shoots, and leaves and is an important multi-purpose plant, one of the oldest known to have medicinal and aromatic properties. It has Mediterranean and Asian origin, and is an annual plant belonging to the Leguminosae family (Baldemir and İlgün, 2015; Bienkowski et al., 2016), containing about 50 species, and 45 of these species are found in the natural flora of Turkey (Davis, 1982; Beyzi et al., 2010). India is the world's largest producer of fenugreek, and it is also grown in South Asia, the Middle East, Far East, China, Iran, Pakistan, Turkey, the Mediterranean region, Europe, North Africa, Australia, Canada, the USA, and Argentina (Basu et al., 2019).

Fenugreek is a plant that has economic value in food, feed, medicine, and cosmetics. Fenugreek seeds are used in traditional medicine in the treatment of diabetes and cancer, as a natural dyestuff in the cosmetics industry, in animal feeding, as a coating material of some products in the food industry, and in making spices (Boeker, 1963; Baytop, 1984; Arslan et al., 1989; Hornok, 1992; Akgül, 1993; Küçük and Gürbüz, 1999; Soylu et al., 2000; Kızıl and Arslan, 2003; Tunçtürk et al., 2011; Abd Elhamid et al., 2016). It is also used as a green manure plant since it is a legume plant, and it has been reported to play a role in improving the physical and chemical properties of the soil (Abdelgani et al., 1999).

Phosphorus (P), which fulfills many functions related to plant growth, development, and metabolism and is essential for young tissues and cannot be renewed, is the most crucial macronutrient element after nitrogen. It is also called the key to life because it regulates many metabolic activities of plant life. If sufficient P is given to the plant, it causes rapid growth and increases plant resistance. In the case of P deficiency, plants stop and slow down the growth of the above-ground organs and accelerate the root growth. P, which also plays an essential role in increasing the yield of legumes by increasing biological activities such as nodulation, nitrogen fixation, nutrient uptake in the soil and in the rhizosphere environment, alleviates adverse effects of drought on physiological parameters in plants (Kacar and Katkat, 2007; Turan and Horuz, 2012; Yadav et al., 2014; Singh and Singh, 2016; Gezgin, 2018).

Generally, the properties of soils in Turkey affect the uptake of soil phosphorus by plants negatively (high pH, high lime content, low organic matter content, high clay content). It is very important to apply appropriate amounts of P to legume plants or to increase the availability of soil phosphorus, according to the results of soil analysis (Gezgin, 2018). Legumes' benefiting from soil phosphorus also differs according to its species and varieties (Gökmen Yılmaz et al., 2017).

Elemental sulfur (S) is a natural material and can be applied to increase the availability of plant nutrients and to reduce deficiencies in calcareous and alkaline soils (Manesh et al., 2013). Sulfur has vital importance in the activation of the process of photosynthesis, carbohydrate metabolism, and certain enzyme systems in plants; it can increase plants' seed and oil yields and protein contents. It is considered to be an essential nutrient in the vitamin and amino acids synthesis of legume plants. Sulfur is found in cysteine, cystine, and methionine, among the amino acids, and in the composition of proteins. It is necessary for chlorophyll formation (Lal et al., 2015). It accelerates root growth and nodule formation (Tonguç et al., 2017; Bolat and Kara, 2017). It has been reported that the amount of nitrogen fixed by legumes increases with the treatment of sulfur, and as a result of this, soil fertility improves (Mohamed El-Sayed Ali, 2018).

The reduced use of fossil fuels in the world and in Turkey and the widespread use of sulfur-free fertilizers have caused sulfur deficiency to emerge as a factor limiting the yield in plant production. Sulfur affects nitrogen utilization efficiency, and adverse effects of sulfur deficiency are observed on the growth of plants, chlorophyll amount, photosynthesis capacities, yield, and yield parameters (Tonguç et al., 2017). Sulfur positively affects not only the above-ground organs of plants but also the root growth, just like phosphorus. As a matter of fact, in the soybean plant given sulfur, the number and weight of side roots, the number and weight of nodules in which nitrogen fixation occurred were reported to increase significantly (Zhao et al., 2008). In beans, plant weight, the number of branches, the number of pods, the number of grains per plant, thousand-seed weight, harvest index, and yield were indicated to significantly increase in comparison with unfertilized plants (Tonguç et al., 2017).

This study was conducted to determine the effects of sulfur and phosphorus doses on the yield and quality characteristics of fenugreek (*T. foenum-graecum* L.).

Materials and methods

The study was carried out between 2016-2018 under the ecological conditions of Siirt province located in the Southeastern Anatolia Region of Turkey, which has a

semi-arid climate. Siirt is located at 37° 58' 7.37" N and 41° 51' 3.87" E coordinates with 894 m altitude (Figure 1, Figure 2).



Figure 1. Location of the study area in Turkey



Figure 2. A photo of the research area

In the second year of the study, less precipitation (522.8 mm) occurred in comparison with the first year (574.2 mm), while the long-term annual precipitation was 634.1 mm. While the highest precipitation occurred in March and April during the 2016-2017 vegetation period, the highest precipitation was recorded in May during the 2017-2018 vegetation period (Table 1). The mean temperature during the study years and the long-term mean temperature (38 years) were 11.2 °C, 13.8 °C, and 11.4 °C, respectively Table 1 (Anonymous, 2018).

Table 1. Climate characteristics of trial area

Climate parameters	Research years	Months								
		Nov	Dec	Jan	Feb	Mar	Apr	May	June	Mean/Sum
Average temperature (°C)	2016-2017	10.4	3.3	3.0	2.7	9.6	14.0	19.5	26.9	11.2
	2017-2018	11.2	8.0	5.7	8.2	13.7	16.8	19.8	27.4	13.8
	Long term*	10.3	4.9	3.0	4.5	8.8	14.3	19.5	26.2	11.4
Monthly precipitation (mm)	2016-2017	55.4	116.6	46.4	29.2	119.2	132.8	74.6	0.0	574.2
	2017-2018	86.0	47.4	56.4	74.2	47.6	61.6	139.6	10.0	522.8
	Long term*	85.1	91.1	82.2	96.6	108.7	96.3	64.3	9.8	634.1

*: 1980-2018

In the study, some physical and chemical analysis results of the soils taken before establishing the field trial were presented in *Table 2*. In the first year (2016-2017), the trial area soils were loamy textured, and they were clay-loam textured in the second year (2017-2018); both trial area soils were slightly alkaline, salt-free, their lime content was "medium calcareous," the organic matter content was "low," and the available potassium (K) content was "sufficient". The available P content of the soils in the first year was determined to be "very little," and the available P content of the soils in the second year was determined to be "low" (*Table 2*).

Table 2. Some physical and chemical properties of the study area soils (0-20 cm)*

Properties	Value	
	2016-2017	2017-2018
Clay, %	47.56	34.16
Silt, %	12.11	26.00
Sand, %	40.33	39.84
pH	7.72	7.53
Electrical conductivity (EC), mS cm ⁻¹	0.363	0.150
Lime (CaCO ₃), %	12.0	8.2
Organic matter, %	1.31	1.78
Available phosphorus, kg P ₂ O ₅ ha ⁻¹	24	49
Available potassium, kg K ₂ O ha ⁻¹	1430	1250

*: Analyses were carried out in Siirt University, Science and Technology Application and Research Center Laboratory

As the plant material in the study, fenugreek (*T. foenum-graecum* L.) seeds, belonging to the "Konya population" and seeded locally in the Central Anatolia Region of Turkey, were used.

In this study, the field trial was established as three replications in randomized blocks according to the factorial trial design. In the study, 4 different sulfur doses (S₀= 0, S₁₀= 10, S₂₀= 20, and S₃₀= 30 kg S ha⁻¹) and 4 different phosphorus doses (P₀= 0, P₃₀= 30, P₆₀= 60, and P₉₀= 90 kg P₂O₅ ha⁻¹) constituted the subject of the study. Elemental sulfur was used as the source of sulfur fertilizer, and triple superphosphate (43-44% P₂O₅) was used as the source of phosphorus fertilizer. According to the

research subjects, both fertilizer forms were mixed by applying them to the soil before seeding.

The seeding process was performed manually on 14 November 2016 in the first year and on 17 November 2017 in the second year on the grooves opened with the help of a marker. In the study, row distance was 30 cm and parcel distance was 100 cm. Each parcel constituted from four rows, while length and width of the parcels were 3 meter by 1.2 meter each with a total area of 3.6 m² per parcel. Sowing norm was 30 kg ha⁻¹, and sowing was done manually in rows opened with the help of a marker. Weed control was performed mechanically by hand several times in both years. At the harvest, two border rows and 50 cm from each side were excluded to eliminate border effects.

Plant height, first pod height, pod length, the number of pods per plant, and the number of seeds per pod were determined in 10 plants randomly selected in each parcel before harvest. The harvest was carried in the entire plot area, excluding borders in the first week of July in both years. The harvested plants were dried in the shade for 3-4 days, and seed yields were calculated per hectare. Crude protein determination in seeds was performed by the Kjeldahl nitrogen determination method, and trigonelline analysis was performed by the HPLC method. In trigonelline analysis was used Agilent TC-C18 (ODS 25 cm * 4.6 mm) column. Column temperature was maintained at 27 °C and the flow rate of the mobile phase was kept at 1 ml per min. The changes in absorbance at wavelength 210 nm were recorded with UV detector. The peak area was calibrated to trigonellin content with a standard.

The data were analyzed by JMP statistical software. A homogeneity test was applied to the data obtained from the study. According to the results of the homogeneity test, they were subjected to combined variance analysis (ANOVA) according to the factorial trial design in randomized blocks. According to the F-test results, differences between the groups were determined by the LSD multiple comparison test. In the study, the correlation coefficients of the pairwise relationships between the examined properties were calculated (Yurtsever, 1984; Düzgüneş et al., 1987).

Results and discussion

Plant height

In the study, the effect of S doses on the plant height of fenugreek was statistically insignificant, and according to S doses, plant height varied between 62.4-65.2 cm as the mean of P doses and years. The effect of phosphorus doses on plant height was determined to be statistically significant at the $p < 0.01$ level. This difference occurred between the P₀ and P₃₀ doses of phosphorus and its P₆₀ and P₉₀ doses. In this study, in which plant height increased according to P doses, the highest plant height was determined in P₆₀ (65.2 cm) and P₉₀ (66.3 cm) P doses. When the interaction of SxP was examined, the highest plant height was determined in the S₀P₉₀ application as 70.1 cm, but no statistically significant difference was found between it and S₀P₆₀, S₁₀P₆₀, S₁₀P₉₀, S₃₀P₀, and S₃₀P₉₀ applications. The lowest plant height was measured in unfertilized parcels (control). SxP interaction was determined to be statistically significant ($p < 0.01$) (Table 3). When the studies on this subject were reviewed, increasing P applications were reported to increase plant height, similarly to the results of our study (Halesh et al., 2000; Khiriya et al., 2001; Nehara et al., 2006; Meena et al., 2012; Mehta et al., 2012; Verma et al., 2014; Srivastava et al., 2015; Ahmad, 2017; Basu et al., 2019).

Table 3. Means of yield components at different sulfur and phosphorus doses in fenugreek

S doses	P doses	Plant height (cm)			First pod height (cm)			Pod length (cm)		
		2017	2018	Mean ¹	2017	2018	Mean ¹	2017	2018	Mean ¹
S ₀	P ₀	59.8	57.8	58.8 e	36.4	46.2	41.3	14.4	13.3	13.8
	P ₃₀	63.8	61.4	62.6 cde	44.5	50.1	47.3	14.5	14.6	14.6
	P ₆₀	69.9	68.8	69.4 ab	43.5	48.9	46.2	14.8	15.3	15.0
	P ₉₀	69.0	71.1	70.1 a	42.3	52.7	47.5	15.4	16.4	15.9
S ₀ Mean		65.6	64.8	65.2	41.7	49.5	45.6 A	14.8	14.9	14.8
S ₁₀	P ₀	60.7	56.9	58.8 e	41.3	38.2	39.8	14.1	14.5	14.3
	P ₃₀	60.8	62.4	61.6 de	40.3	42.6	41.4	14.7	15.4	15.1
	P ₆₀	65.1	66.5	65.8 a-d	43.1	42.7	42.9	15.2	15.7	15.4
	P ₉₀	66.3	68.53	67.4 abc	40.4	44.7	42.5	17.7	16.3	17.0
S ₁₀ Mean		63.2	63.6	63.4	41.3	42.1	41.7 B	15.5	15.5	15.5
S ₂₀	P ₀	61.7	64.5	63.1 cde	39.9	36.1	38.1	14.2	14.2	14.2
	P ₃₀	62.6	60.1	61.4 de	39.1	40.0	39.6	15.9	15.8	15.8
	P ₆₀	61.7	65.7	63.7 cde	43.1	43.2	43.1	15.2	15.2	15.2
	P ₉₀	62.6	60.1	61.4 de	40.9	42.4	41.7	16.2	16.3	16.3
S ₂₀ Mean		62.2	62.6	62.4	40.7	40.4	40.6 B	15.4	15.4	15.4
S ₃₀	P ₀	64.4	67.1	65.8 a-d	35.2	35.5	35.4	14.2	14.5	14.4
	P ₃₀	64.1	64.3	64.2 b-d	43.4	40.3	41.8	15.3	15.5	15.4
	P ₆₀	61.3	62.5	61.9 de	46.3	48.3	47.3	15.9	16.1	16.0
	P ₉₀	66.4	66.1	66.3 a-d	41.8	44.6	43.2	15.2	15.6	15.4
S ₃₀ Mean		64.1	65.0	64.5	41.7	42.2	41.9 B	15.2	15.4	15.3
Phosphorus mean										
P ₀		61.7	61.6	61.6 b	38.2	39.0	38.6 b	14.3	14.2	14.2 c
P ₃₀		62.8	62.1	62.5 b	41.8	43.3	42.6 a	15.1	15.3	15.2 b
P ₆₀		64.5	65.9	65.2 a	43.9	45.8	44.9 a	15.3	15.6	15.4 b
P ₉₀		66.1	66.5	66.3 a	41.3	46.1	43.7 a	16.1	16.2	16.1 a
Means		63.8	64.0		41.3	43.5		15.2	15.3	
CV (%)		7.2			10.8			6.5		
Year (Y)		ns			*			ns		
Sulfur (S)		ns			**			ns		
Phosphorus (P)		**			**			**		
SxP		**			ns			ns		
SxPxY		ns			ns			ns		

S₀= Control, S₁₀= 10 kg ha⁻¹ sulfur, S₂₀= 20 kg ha⁻¹ sulfur, S₃₀= 30 kg ha⁻¹ sulfur, P₀= Control, P₃₀= 30 kg ha⁻¹ phosphorus, P₆₀= 60 kg ha⁻¹ phosphorus, P₉₀= 90 kg ha⁻¹ phosphorus, ¹: The difference between the means indicated by the same letter in the same column and group is not significant, CV: Coefficient of variation, ns: Not significant, *: p<0.05, **: p<0.01

First pod height

In the fenugreek plant, S and P doses had a statistically significant effect on the first pod height at the p<0.01 level. The highest first pod height was measured in plants in parcels not treated with S, and no statistically significant difference was found between the other S doses. In other words, increased S doses did not affect the first pod height (Table 3). Tunçtürk et al. (2011) reported that the highest first pod height was obtained in the dose of 40 kg S ha⁻¹ and that contrary to the results of our study, the first pod height increased with increasing S doses. It is thought that seeding time and genotype differences may be effective in this difference in the literature. As a matter of fact, Babagil (2010) stated that the first pod height is a property that is significantly affected by genotype and environmental factors. When the results of phosphorus doses were examined, as the mean of years and S doses, the lowest first pod height (38.6 cm) was

measured in parcels not treated with phosphorus fertilizer, while the P₃₀, P₆₀, and P₉₀ doses of phosphorus were statistically in the same group and yielded the highest values (Table 3). In the studies conducted on fenugreek, the highest first pod height was determined in 40 kg P ha⁻¹ by Khiriya and Singh (2003) and in 30 kg P ha⁻¹ doses by Tunçtürk (2011), Nehara et al. (2006) reported that increased phosphorus doses increased yield properties.

Pod length

While the effects of S doses on pod length were statistically insignificant in the fenugreek plant, the effects of P doses were significant at the $p < 0.01$ level. The pod length varied between 14.8-15.5 cm according to sulfur doses. The pod length was observed to increase in parallel with the increasing P doses. The highest pod length was measured to be 16.1 cm at P₉₀ dose, and the lowest value was measured in the control parcel (14.2 cm) (Table 3). Similarly to our results, Khiriya et al. (2001), Khiriya and Singh (2003), Bhunia et al. (2006), and Meena et al. (2012) reported that increased P doses increased the pod length.

Number of pods

The statistical analysis results demonstrated that S and P doses and SxP interaction had significant effects on the number of pods per plant at the $p < 0.01$ level. In the study, the number of pods was determined to increase in the fenugreek plant depending on the increase in S and P doses. In both fertilizer applications, the highest values were determined at the highest fertilizer doses (Table 4). In the studies conducted by Kumar (2011) on pea and by Nawange et al. (2011) on chickpeas, it was reported that increasing phosphorus and sulfur applications increased the number of pods.

The number of pods per plant was reported to vary between 7.7-8.5 by Tunçtürk (2011), between 30.6-33.0 by Meena et al. (2012), between 46.73-50.72 by Lal et al. (2015), between 6.4-8.6 by Srivastava et al. (2015), and between 11.6-23.1 by Mitoo et al. (2018). The reason for the number of pods per plant obtained in this study to be higher than these values in the literature may be differences in genotype, soil, climate, and cultural practices.

When the SxP interaction was examined, the highest number of pods was obtained from S₃₀P₆₀ and S₃₀P₉₀ applications, while the lowest number of pods was obtained from the S₀P₀ application (Table 4).

Number of seeds per pod

When sulfur applications were examined, it was observed that the number of seeds per pod increased up to S₂₀ dose (15.59 seeds) depending on the increase in S doses, and after this dose, it was observed that it decreased statistically significantly. This difference between sulfur doses was statistically significant at the $p < 0.01$ level. In terms of phosphorus doses, the number of seeds per pod increased with increasing P doses. Although the highest value was obtained to be 15.85 seeds at P₉₀ dose, the difference between P₉₀ and P₆₀ doses (15.62 seeds) was statistically insignificant. This difference between phosphorus doses was determined to be statistically significant at the $p < 0.01$ level. According to the sulfur x phosphorus interaction, the lowest number of seeds per pod was in S₀P₀ interaction (12.40), and the highest values were determined in S₂₀P₉₀, S₂₀P₃₀, S₂₀P₆₀, S₃₀P₆₀, S₃₀P₉₀, S₀P₉₀, S₁₀P₉₀, S₁₀P₆₀, and S₀P₆₀ interactions. SxP

interaction was determined to be very important in terms of the number of seeds per pod ($p < 0.01$) (Table 4). Srivastava et al. (2015) reported that the number of seeds per pod varied between 13.20-16.60 and that the application of phosphorus increased the number of seeds per pod in the fenugreek plant. Mitoo et al. (2018) stated that the number of seeds per pod varied between 11.67-15.07 and increased with increasing phosphorus and sulfur applications.

Table 4. Means of yield components at different sulfur and phosphorus doses in fenugreek

S doses	P doses	Number of pods (pods/plant)			Number of seeds in pod (seeds/pod)			Thousand-seed weight (g)		
		2017	2018	Mean ¹	2017	2018	Mean ¹	2017	2018	Mean ¹
S ₀	P ₀	37.33	36.30	36.82 g	12.89	11.90	12.40 d	12.24	13.10	12.67
	P ₃₀	45.70	42.57	44.13 c-f	14.57	15.07	14.82 bc	13.23	13.20	13.21
	P ₆₀	43.70	45.67	44.68 cde	15.46	15.63	15.55 ab	13.07	13.42	13.25
	P ₉₀	45.60	49.30	47.45 bc	15.76	16.13	15.95 a	13.52	13.30	13.41
S ₀ Mean		43.08	43.68	43.27 B	14.67	14.68	14.68 C	13.02	13.25	13.14 B
S ₁₀	P ₀	39.33	39.73	39.53 fg	14.07	14.87	14.47 c	12.58	13.54	13.06
	P ₃₀	39.93	43.13	41.53 d-g	14.73	14.77	14.75 c	13.73	13.89	13.81
	P ₆₀	45.57	45.73	45.65 cd	15.40	15.73	15.57 ab	14.27	13.62	13.95
	P ₉₀	44.90	48.30	46.60 bcd	15.70	15.73	15.72 a	13.42	14.12	13.77
S ₁₀ Mean		42.43	43.97	43.33 B	14.98	15.28	15.13 B	13.50	13.79	13.65 AB
S ₂₀	P ₀	39.70	38.47	39.08 fg	14.63	14.60	14.62 c	12.64	13.21	12.93
	P ₃₀	43.57	45.00	44.28 cde	16.17	15.87	16.02 a	13.38	13.22	13.30
	P ₆₀	48.27	46.90	47.58 bc	15.63	15.63	15.63 a	13.93	13.72	13.82
	P ₉₀	51.90	50.03	50.97 b	16.00	16.20	16.10 a	14.60	15.91	15.26
S ₂₀ Mean		45.86	45.10	45.48 B	15.61	15.58	15.59 A	13.64	14.02	13.83 A
S ₃₀	P ₀	37.93	38.93	38.43 g	14.23	14.57	14.40 c	12.67	13.18	12.93
	P ₃₀	39.13	41.93	40.53 efg	14.83	14.80	14.82 bc	12.88	13.32	13.10
	P ₆₀	56.33	63.20	59.77 a	15.70	15.73	15.72 a	14.03	14.00	14.02
	P ₉₀	56.77	55.97	56.37 a	15.63	15.67	15.65 a	15.24	15.47	15.36
S ₃₀ Mean		47.54	50.01	48.78 A	15.10	15.19	15.15 B	13.70	13.99	13.84 A
Phosphorus mean										
P ₀		38.58	38.61	38.59 c	13.95	13.98	13.97 c	12.54	13.31	12.90 c
P ₃₀		42.08	43.16	42.62 b	15.08	15.12	15.10 b	13.83	14.20	13.36 bc
P ₆₀		48.47	50.38	49.42 a	15.55	15.68	15.62 a	13.26	13.41	13.76 b
P ₉₀		49.80	50.90	50.34 a	15.78	15.93	15.85 a	13.69	14.70	14.45 a
Means		44.73	45.76		15.09	15.18		13.47	13.76	
CV (%)		8.9			4.6			6.6		
Year (Y)		ns			ns			ns		
Sulfur (S)		**			**			*		
Phosphorus (P)		**			**			**		
SxP		**			**			ns		
SxPxY		ns			ns			ns		

S₀= Control, S₁₀= 10 kg ha⁻¹ sulfur, S₂₀= 20 kg ha⁻¹ sulfur, S₃₀= 30 kg ha⁻¹ sulfur, P₀= Control, P₃₀= 30 kg ha⁻¹ phosphorus, P₆₀= 60 kg ha⁻¹ phosphorus, P₉₀= 90 kg ha⁻¹ phosphorus, ¹: The difference between the means indicated by the same letter in the same column and group is not significant, CV: Coefficient of variation, ns: Not significant, *: $p < 0.05$, **: $p < 0.01$

Thousand-seed weight

The results of the statistical analysis showed that S doses had a significant effect on the thousand-seed weight at the $p < 0.05$ level and P doses at the $p < 0.01$ level. When Table 4 was examined, the highest thousand-seed weight among S doses was

determined at S₂₀ (13.83 g) and S₃₀ (13.84 g) doses. However, no statistically significant difference was determined between them and S₁₀ (13.65 g) S dose (*Table 4*). Ramkishor and Kumawat (2015) stated that different doses of S applied to the fenugreek plant had significant effects on the thousand-seed weight, and that the application of 40 and 60 kg S per hectare increased the thousand-seed weight compared to control subjects. Srivastava et al. (2015) obtained the highest thousand-seed weight to be 13.75 and 13.76 g at the 40 and 80 kg ha⁻¹ P doses, respectively. Tunçtürk et al. (2011) stated that the thousand-seed weight was significantly affected by S application in the first year and that the highest thousand-seed weight was obtained to be 18.8 g at dose of 40 kg ha⁻¹. The increased P doses increased the thousand-seed weight in the fenugreek plant. While the lowest thousand-seed weight was determined at P₀ dose (12.90 g), the highest thousand-seed weight was determined at P₉₀ dose (14.45 g) (*Table 4*). Tunçtürk (2011) reported that P doses increased the thousand-seed weight, and the highest thousand-seed weight was obtained from 60 and 90 kg ha⁻¹ P (17.8 g and 18.0 g) applications. The highest thousand-seed weight was determined by Sammuria and Yadav (2008) from 40 and 60 kg ha⁻¹ P application (11.62 g and 11.74 g, respectively), and by Mitoo et al. (2018) to be 9.98 g from 40 kg ha⁻¹ P application, which was the highest dose.

Seed yield

The variance analysis results showed that S and P doses and SxP interaction had significant effects on seed yield at the $p < 0.01$ level (*Table 5*). When *Table 5* was examined, it was determined that the highest seed yield was obtained to be 2224 kg ha⁻¹ from the S₃₀P₉₀ application, and the lowest seed yield was obtained to be 1038 kg ha⁻¹ from the S₀P₀ application. As P and S doses increased, seed yield was determined to increase in fenugreek. In both fertilizer applications, the highest seed yields were determined at the highest fertilizer doses. As the mean of years and P doses, the highest fenugreek seed yield with the sulfur application was obtained to be 1822 kg/ha (S₃₀), and as the mean of years and S doses, the highest fenugreek seed yield in phosphorus fertilizer applications was obtained to be 1810 kg ha⁻¹ (P₉₀) (*Table 5*). Many researchers stated that the phosphorus element is necessary for energy transfer in plants and that the yield, and above ground and root development of the plant are adversely affected in P deficiency or excess (Kacar and Katkat, 2007; Singh and Singh, 2016). Tunçtürk et al. (2011), Lal et al. (2015), Ramkishor and Kumawat (2015), Singh Manohar et al. (2017), and Verma et al. (2017) reported that increasing sulfur doses increase seed yield, and Halesh et al. (2000), Tunçtürk (2011), and Meena et al. (2012) reported that increasing phosphorus doses increase seed yield in the fenugreek plant.

Protein content

In the study, the effects of S and P doses on the protein content of fenugreek seed were found to be statistically significant ($p < 0.01$). In terms of sulfur doses, as the mean of years and P doses, the highest protein ratio was determined at S₃₀ (26.0%) dose and the lowest at S₀ (25.3%) dose. In terms of phosphorus doses, increasing P doses increased protein content in fenugreek, but statistically, P doses, except for P₀, were in the same group (*Table 5*). Baldaneeya Nitesh (2018) and Tunçtürk et al. (2011) stated that they determined the highest protein content at a dose of 40 kg ha⁻¹ S. Mehta et al. (2012) reported that although increasing P doses increased protein content, there was no

difference between 20 and 40 kg ha⁻¹ P doses. Tunçtürk (2011) also reported similar results.

Table 5. Means of yield, protein and trigonelline ratio at different sulfur and phosphorus doses in fenugreek

S doses	P doses	Seed yield (kg ha ⁻¹)			Protein ratio (%)			Trigonelline ratio (%)		
		2017	2018	Mean*	2017	2018	Mean*	2017	2018	Mean*
S ₀	P ₀	1005	1071	1038 i	24.6	25.1	24.9	0.78	0.83	0.81 i
	P ₃₀	1121	1241	1182 h	24.9	25.5	25.2	0.88	0.91	0.90 hi
	P ₆₀	1323	1432	1378 fg	25.2	25.6	25.4	0.91	0.94	0.93 gh
	P ₉₀	1520	1527	1523 de	25.5	25.8	25.6	0.93	0.95	0.94 efg
S ₀ Mean		1242	1318	1280 D	25.1	25.5	25.3 C	0.89	0.88	0.89 D
S ₁₀	P ₀	1199	1246	1223 h	25.6	24.9	25.3	0.84	0.89	0.87 h
	P ₃₀	1270	1327	1299 gh	25.8	25.2	25.5	0.87	0.89	0.88 h
	P ₆₀	1462	1416	1439 ef	25.7	25.5	25.6	0.92	0.94	0.93 gh
	P ₉₀	1501	1559	1530 de	25.8	25.7	25.8	0.94	0.96	0.95 ef
S ₁₀ Mean		1358	1387	1373 C	25.7	25.4	25.5 B	0.91	0.89	0.90 C
S ₂₀	P ₀	1234	1283	1259 gh	25.7	25.3	25.5	0.98	0.99	0.99 e
	P ₃₀	1329	1374	1351 fg	25.9	25.6	25.8	1.09	1.12	1.11 c
	P ₆₀	1661	1584	1622 cd	26.0	25.8	25.9	1.10	1.14	1.12 c
	P ₉₀	1902	2027	1964 b	25.7	25.6	25.9	1.19	1.22	1.21 ab
S ₂₀ Mean		1531	1567	1549 B	25.8	25.7	25.8 B	1.09	1.12	1.11 B
S ₃₀	P ₀	1428	1430	1429 ef	25.7	26.1	25.9	1.04	1.06	1.05 d
	P ₃₀	1692	1687	1690 c	26.0	26.2	26.1	1.16	1.19	1.18 b
	P ₆₀	1898	1989	1944 b	26.0	26.4	26.2	1.20	1.23	1.22 ab
	P ₉₀	2043	2405	2224 a	25.3	26.5	25.9	1.25	1.26	1.26 a
S ₃₀ Mean		1765	1879	1822 A	25.8	26.3	26.0 A	1.17	1.19	1.18 A
Phosphorus mean										
P ₀		1217	1258	1237 d	25.4	25.4	25.4 b	0.91	0.94	0.93 c
P ₃₀		1353	1408	1380 c	25.7	25.6	25.6 a	1.00	1.03	1.02 b
P ₆₀		1586	1605	1596 b	25.8	25.8	25.8 a	1.03	1.06	1.05 b
P ₉₀		1741	1879	1810 a	25.6	26.0	25.8 a	1.08	1.10	1.09 a
Means		1474	1538		25.6	25.7		1.02	1.04	
CV (%)		7.0			1.6			4.1		
Year (Y)		**			ns			ns		
Sulfur (S)		**			**			**		
Phosphorus (P)		**			**			**		
SxP		**			ns			**		
SxPxY		ns			ns			ns		

S₀= Control, S₁₀= 10 kg ha⁻¹ sulfur, S₂₀= 20 kg ha⁻¹ sulfur, S₃₀= 30 kg ha⁻¹ sulfur, P₀= Control, P₃₀= 30 kg ha⁻¹ phosphorus, P₆₀= 60 kg ha⁻¹ phosphorus, P₉₀= 90 kg ha⁻¹ phosphorus, *: The difference between the means indicated by the same letter in the same column and group is not significant, CV: Coefficient of variation, ns: Not significant, **: p<0.01

Trigonelline content

The effects of S and P doses and SxP interaction on the trigonelline content of fenugreek seeds were found to be significant at the p<0.01 level (Table 5). When Table 5 was examined, the highest trigonelline content was determined in S₃₀P₉₀ (1.26%), but there was no statistically significant difference between S₃₀P₆₀ (1.22%) and S₂₀P₉₀ (1.21%). On the other hand, as both S and P doses increased, trigonelline content was observed to increase (Table 5). Kan et al. (2007) stated that applied different phosphorous fertilizer sources did not change the trigonelline content, that trigonelline

content varied between 0.86-1.26%, and that ecological factors might be effective on the trigonelline content. Dar et al. (2015) reported that a 40 kg P ha⁻¹ application increased the trigonelline content. Mutlu (2011) reported in the study conducted in fenugreek of different origins that the trigonelline content varied between 0.66-1.40%. The trigonelline content obtained in this study was higher than the values reported by Akgül (1993) (0.36%), Mehrafarin et al. (2010) (0.20-0.36%), and Mathur and Yadav (2011) (0.27%).

Relationships between the investigated properties

The simple correlation coefficients showing linear relationships between the seed yield per plant and the investigated properties were presented in Table 6. The correlation analysis revealed that the seed yield was positively and significantly correlated with the number of pods per plant ($r= 0.703^{**}$), pod length ($r= 0.334^{**}$), the number of seeds per pod ($r= 0.509^{**}$), and thousand-seed weight ($r= 0.571^{**}$). Increases that occurred in these properties caused significant increases in the plant seed yield. The highest correlation coefficients in terms of seed yield were determined in the relationships between the number of pods per plant, the number of seeds per pod, and thousand-seed weight. Parchin et al. (2019) stated that there was a positive relationship between the number of pods per plant and seed yield.

Table 6. Correlation coefficients related to pairwise relations between the seed yield and other properties

	2	3	4	5	6	7
1. Plant height	0.274**	0.094	0.218*	0.243*	0.104	0.154
2. First pod height	-	0.280**	0.080	0.068	0.110	0.078
3. Number of pods per plant ⁻¹		-	0.391**	0.478**	0.436**	0.703**
4. Pod length			-	0.343**	0.321**	0.334**
5. Number of seeds per pod ⁻¹				-	0.299**	0.509**
6. Thousand-seed weight					-	0.571**
7. Seed yield						-

*: Significant at the $p<0.05$ level, **: Significant at the $p<0.01$ level

A positive and significant relationship was determined between plant height and the first pod height. Positive and significant relationships were detected between the number of pods per plant and pod length and between the number of seeds per pod and thousand-seed weight. Positive and significant relationships were found between the thousand-seed weight and the number of pods per plant and between pod length and the number of seeds per pod. The number of pods per plant was stated to be the most important factor affecting seed yield in many studies (Şehirali, 1980; Pooran-Chand, 1999; Amini et al., 2002; Kumar Singh et al., 2019, Singh et al., 2019), and finding the number of pods per plant as the most leading factor in this study supports the other literature on this subject.

Conclusions

This study, which was carried out under the ecological conditions of Siirt province located in the Southeastern Anatolia Region of Turkey, which has a semi-arid climate, revealed the importance of sulfur and phosphorus fertilizers when an evaluation was

made in terms of yield and some quality criteria. Increased sulfur and phosphorus doses positively affected all of the investigated properties. When an evaluation was made in terms of yield and yield components, 30 kg S and 90 kg P doses could be stated to be suitable for per hectare. However, it was revealed that studies involving further increasing doses of both elements should be performed due to the linear increase determined in yield and quality properties in parallel to the increasing sulfur and phosphorus doses.

REFERENCES

- [1] Abdelgani, M. E., Elsheikh, E. A. E., Mukhtar, N. O. (1999): The effect of Rhizobium inoculation and chemical fertilization on seed quality of fenugreek. – Food Chemistry 64: 289-293.
- [2] Abd Elhamid, E. M., Sadak, M. S., Tawfik, M. M. (2016): Physiological response of fenugreek plant to the application of proline under different water regimes. – Research Journal of Pharmaceutical, Biological and Chemical Sciences 7(3): 580-594.
- [3] Ahmad, T. S. (2017): Study the effect of phosphorus fertilizer and seaweeds extracts on the growth, yield, and oil percentage of fenugreek *Trigonella foenum-graecum* L. – Tikrit Journal for Agricultural Sciences 17(2): 87-93.
- [4] Akgül, A. (1993): Spices science and technology. – Food Technology Association Publications, No.15. Ankara, Turkey. (In Turkish).
- [5] Amini, A., Ghannadha, M., Abd-Mishani, C. (2002): Genetic diversity and correlation between different traits in common bean (*Phaseolus vulgaris* L.). – Iranian J. of Agricultural Sci. 33(4): 605-615.
- [6] Anonymous (2018): Data from the General Directorate of Meteorology.
- [7] Arslan, N., Tekeli, S., Gençtan, T. (1989): Seed yields of fenugreek (*Trigonella foenum-graecum* L.) populations from different regions. – VIII. Proceedings Book of Herbal Medicine Raw Materials Meeting Proceedings 2: 93-97.
- [8] Babagil, G. E. (2010): A study on yield and yield components of some chickpea (*Cicer arietinum* L.) varieties in Muş ecological conditions. – Journal of Tekirdag Agricultural Faculty 7(3): 181-186.
- [9] Baldaneeya Nitesh, S. (2018): Response of fenugreek (*Trigonella foenum-graecum* L.) to phosphorus and sulphur. – Junagadh Agricultural University, MSc. Thesis.
- [10] Baldemir, A., İlgin, S. (2015): Usage areas of fenugreek in the past and today: *Trigonella foenum-graecum* L. – Lokman Hekim Journal 5(1): 1-4.
- [11] Basu, S. K., Cetzal-Ix, W., Zandi, P., Mozdzeń, K., Siavoshi, M. (2019): Rich potential for fenugreek (*Trigonella foenum-graecum* L.) cultivation in Iran. – The 2nd International Conference on Medicinal Plants, Organic Farming, Natural and Medicinal Materials, 13-14 February.
- [12] Baytop, T. (1984): Treatment with Plants. – İstanbul University, Faculty of Pharmacy Publications. No: 3255. (In Turkish).
- [13] Bhunia, S. R., Chauhan, R. P. S., Yadav, B. S., Bhati, A. S. (2006): Effect of phosphorus, irrigation and *Rhizobium* on productivity, water use and nutrient uptake in fenugreek (*Trigonella foenum-graecum* L.). – Indian Journal of Agronomy 51(3): 239-241.
- [14] Boeker, P. (1963): Forage Crops. – Ege University, Faculty of Agriculture. No: 82. Translation: Demir, I. pp. 97-98.
- [15] Bolat, İ., Kara, Ö. (2007): Plant nutrients: sources, functions, deficiencies and redundancy. – Journal of Bartın Faculty of Forestry 19(1): 218-228. (In Turkish).
- [16] Dar, T. A., Uddin, M., Masroor Khan, M. A., Akbar A., Rasool Mir, S., Varshney, L. (2015): Effect of Co-60 gamma irradiated chitosan and phosphorus fertilizer on growth,

- yield and trigonelline content of *Trigonella foenum-graecum* L. – Journal of Radiation Research and Applied Sciences 8: 446-458.
- [17] Davis, P. H. (1982): Flora of Turkey and The East Aegean Islands. – Edinburgh Univ. Press 3: 465-482.
- [18] Düzgüneş, O., Kesici, T., Kavuncu, O., Gürbüz, F. (1987): Research and Experimental Methods. – Ank. Univ. Agr. Fac. Pub. Lecture book. No. 295. (In Turkish).
- [19] Gezgin, S. (2018): Plant nutrition and fertilization in edible legumes. – Edible Legumes Workshop, 10-12 Mayıs 2017, Adana/Turkey, pp. 181-195. (In Turkish).
- [20] Gökmen Yılmaz, F., Harmankaya, M., Gezgin, S., Dursun, N., Hamurcu, M., Hakkı, E. E. (2017): Phosphorus efficiency of mainly grown beans in Turkey. – IPNC Congress, 21-24.08.2017, Kopenhagen/Danimarka.
- [21] Halesh, D. P., Gowda, M. C., Forooqi, A. A., Vasundhara, M., Srinivasappa, K. N. (2000): Influence of nitrogen and phosphorus on growth, yield and nutrient content of fenugreek (*Trigonella foenum - graecum* L.). Spices and aromatic plants: challenges and opportunities in the new century. – Contributory papers. Centennial conference on spices and aromatic plants, Calicut, Kerala, India, 20-23 September, pp. 191-194.
- [22] Hornok, L. (1992): The cultivation of medicinal plants. – In: Hornok, L. (ed.) Cultivation and processing of medicinal plants. Budapest, pp. 289-290.
- [23] Kacar, B., Katkat, A. V. (2007): Plant nutrition. – Nobel Publication Distribution. (In Turkish).
- [24] Kan, Y., Kartal, M., Abuataker, M. (2007): The effect of organic and inorganic fertilizer on some quality characters of fenugreek seeds (*Trigonella foenum graecum* L.). – Selcuk University Journal of Agriculture 21(41): 118-122.
- [25] Khiriya, K. D., Sheoran, R. S., Singh, B. S. (2001): Growth analysis of fenugreek (*Trigonella foenum- graecum* L.) under various levels of farmyard manure and phosphorus. – Journal of Spices and Aromatic Crops 10(2): 105-110.
- [26] Khiriya, K. D., Singh, B. S. (2003): Effect of phosphorus and farmyard manure on yield, yield attributes and nitrogen, phosphorus and potassium uptake of fenugreek (*Trigonella foenum-graecum* L.). – Indian Journal of Agronomy 48(1): 62-65.
- [27] Kızıl, S., Arslan, N. (2003): Investigation of the effects on yield and yield components of different sowing rates in some fenugreek (*Trigonella foenum-graecum* L.) lines. – Journal of Agriculture Science 9(4): 395-401.
- [28] Küçük, M., Gürbüz, B. (1999): A research on oil content and fatty acid composition of some fenugreek (*Trigonella foenum graecum* L.) lines. – Journal of Food 24(2): 99-101.
- [29] Kumar, J. (2011): Effect of phosphorus and sulphur application on performance of vegetable pea (*Pisum sativum* L.) cv. pant matar-2. – Legume Research-An International Journal 34(4): 292-295.
- [30] Kumar Singh, A., Singh, D. R., Singh, A., Kumar Maurya, J., Pandey, V. P., Sriom. (2019): Studies on character association and path analysis of yield with important yield contributing traits in fenugreek (*Trigonella foenum-graecum* L.). – Journal of Pharmacognosy and Phytochemistry 8(3): 4616-4619.
- [31] Lal, G., Singh, B., Mehta, R. S., Singh, R., Maheria, S. P. (2015): Performance of fenugreek (*Trigonella foenum-graecum* L.) as influenced by sulphur and zinc. – International Journal of Seed Spices 5(1): 29-33.
- [32] Manesh, A. K., Armin, M., Moeini, M. J. (2013): The effect of sulfur application on yield and yield components of corn in two different planting methods in saline conditions. – International Journal of Agronomy and Plant Production 4(7): 1474-1478.
- [33] Mathur, L., Yadav, R. K. (2011): Effect of salicylic acid on trigonelline production in *Trigonella foenum- graecum* L. cell suspension culture. – International Referred Research Journal 1(17): 137-138.
- [34] Meena, R. K., Meena, R. N., Yadav, L., Sharma, O. P. (2012): Effect of phosphorus, molybdenum, PSB levels on growth, productivity and nutrient uptake of fenugreek (*Trigonella foenum-graecum* L.). – Environment & Ecology 30(3B): 825-829.

- [35] Mehrafarin, A., Qaderi, A., Rezazadeh, Sh., Naghdi Badi, H., Noormohammadi, Gh., Zand, E. (2010): Bioengineering of important secondary metabolites and metabolic pathways in fenugreek (*Trigonella foenum-graecum* L.). – Journal of Medicinal Plants 9(35): 1-18.
- [36] Mehta, R. S., Anwer, M. M., Aishwath, O. P., Meena, R. S. (2012): Growth, yield and quality of fenugreek (*Trigonella foenum-graecum* L.) as influenced by nitrogen, phosphorus and bio-fertilizers. – Indian Journal of Horticulturae 69(1): 94-97.
- [37] Mitoo, S. M. M. R., Sujana, M. H. K. (2018): Faijul, I. Effect of phosphorous and sulphur fertilization on seed yield of fenugreek (*Trigonella foenum-graecum* L.). – IOSR Journal of Agriculture and Veterinary Science 11(1): 5-17.
- [38] Mohamed El-Sayed Ali, M. (2018): Efficiency of elemental sulfur and phosphorus fertilizer in enhancing soybean (*Glycine max* L.) growth and yield in a clayey soil. – Egyptian Journal of Soil Science 58(2): 233-244.
- [39] Mutlu, S. (2011): Determination of some phenological, morphological, yield and quality characteristics of fenugrecks (*Trigonella foenum-graceum* L.) from different origins. – Master Thesis. OMÜ, Graduate School of Natural and Applied Sciences. 103p. (In Turkish).
- [40] Nawange, D. D., Yadav, A. S., Singh, R. V. (2011): Effect of phosphorus and sulphur application on growth, yield attributes and yield of chickpea (*Cicer arietinum* L.). – Legume Research 34(1): 48-50.
- [41] Nehara, K. C., Kumawat, P. D., Singh, B. P. (2006): Response of fenugreek (*Trigonella foenum-graecum*) to phosphorus, sulphur and plant-growth regulators under semi-arid eastern plains zone of Rajasthan. – Indian Journal of Agronomy 51(1): 73-76.
- [42] Parchin, R. A., Ghomi, A. A. N., Hasanali, N. B., Eskandari, A., Navabpour, S., Mehrafarin, A. (2019): Growth characteristics and phytochemical responses of Iranian fenugreek (*Trigonella foenum-graecum* L.) exposed to gamma irradiation. – Industrial Crops & Products 139: 111593.
- [43] Pooran-Chand, P. (1999): Character association and path analysis in rajmash. – Madras Agricultural J. 85: 188-190.
- [44] Ramkishor, V. H. P., Kumawat, B. L. (2015): Effect of clay mixing, irrigation and sulphur on growth and yield of fenugreek on loamy sand soil. – Asian Journal of Soil Science 10(1): 29-33.
- [45] Sammuria, R., Yadav, R. S. (2008): Effect of phosphorus and zinc application on growth and yield of fenugreek (*Trigonella foenum-graecum*) and their residual effect on succeeding pearl millet (*Pennisetum glaucum*) under irrigated conditions of north west Rajasthan. – Indian Journal of Agricultural Sciences 78(1): 61-64.
- [46] Singh, N., Singh, G. (2016): Response of lentil (*Lens culinaris* Medikus) to phosphorus-A review. – Agricultural Reviews 37(1): 27-34.
- [47] Singh, A., Pandey, V. P., Kumar, P., Kumar, M., Kumar, S., Vaishya, D. V. (2019): To study of correlation coefficient and path coefficient analysis for different characters in germplasm of fenugreek (*Trigonella foenum graecum* L.). – Journal of Pharmacognosy and Phytochemistry 8(1): 857-861.
- [48] Singh Manohar, M., Shivran, A. C., Choudhary, A. (2017): Effect of sulphur and molybdenum on yield attributes, yield and economics of fenugreek. – Chemical Science Review and Letters 7(21): 94-97.
- [49] Soylu, S., Sade, B., Atalay, E., Pilgir, Ç., Çetinkaya, Ü. (2000): Effects of different sowing times on yield and yield components of fenugreek (*Trigonella foenum graecum* L.) genotypes. – Selcuk University Journal of Agriculture 14(22): 131-142. (In Turkish).
- [50] Srivastava, M., Niraj Kumar, J. H. A., Singh, B. H. (2015): Effect of nitrogen and phosphorus on growth, leaf and seed yield of fenugreek (*Trigonella foenum-graecum* L.) cv. 'Pusa early bunching'. – Annals of Agri-Bio-Research 20(1): 51-53.

- [51] Şehirali, S. (1980): The effect of sowing frequency on some yield related traits in dwarf bean (*Ph. vulgaris* L. var. *nanus* Dekap). – A. Ü. Agr. Fac. Pub: 738. Scientific Research and Studies: 29, Ankara, Turkey.
- [52] Tonguç, M., Şanlı, A., Karakurt, Y., Ünlü, H. (2017): Determining the effects of sulphur applications on yield and yield components of bean. – *Alatarım* 16(1): 44-51. (In Turkish).
- [53] Tunçtürk, R. (2011): The effects of varying row spacing and phosphorus doses on the yield and quality of fenugreek (*Trigonella foenum-graecum* L.). – *Turkish Journal of Field Crops* 16(2): 142-148.
- [54] Tunçtürk, R., Çelen, A. E., Tunçtürk, M. (2011): The effects of nitrogen and sulphur fertilizers on the yield and quality of fenugreek (*Trigonella foenum-graecum* L.). – *Turkish Journal of Field Crops* 16(1): 69-75.
- [55] Turan, M., Horuz, A. (2012): Plant nutrition. – In: Rüştü Karaman, M. (ed.) Basic principles of plant nutrition. pp. 123-347. (In Turkish).
- [56] Verma, S. R., Shivran, A. C., Bhanwaria, R., Singh, M. (2017): Effect of vermicompost and sulphur on growth, yield and nutrient uptake of fenugreek (*Trigonella foenum-graecum* L.). – *Trends in Biosciences* 10(20): 3978-3981.
- [57] Yadav, S. K., Patel, A. G., Yadav, B. L. (2014): Yield, quality and soil fertility of cluster bean (*Cyamopsis tetragonoloba* L.) as influenced by various row spacing and levels of phosphorus. – *Advance Research Journal Of Crop Improvement* 5(2): 101-104.
- [58] Yurtsever, N. (1984): Experimental Statistical Methods. Ministry of Agriculture, Forestry and Rural Affairs. – General Directorate of Rural Services Publication No: 623. (In Turkish).
- [59] Zhao, Y., Xiao, X., Bi, D., Hu, F. (2008): Effects of sulfur fertilization on soybean root and leaf traits and soil microbial activity. – *Journal of Plant Nutrition* 31: 473-483.