

ANALYZING THE IMPACT OF BURNING ON VEGETATION IN HIMALAYAN CHIR PINE (*PINUS ROXBURGHII* SARG.) FORESTS, INDIA

KUMAR, R.^{1*} – PANDEY, R.²

¹*Himalayan Forest Research Institute, Conifer Campus, Panthaghati, Shimla-171013, India*

²*Indian Council of Forestry Research and Education (ICFRE), Dehradun-248195, India*

**Corresponding author*

e-mail: ranjeet@icfre.org; phone: +91-898-837-8338

(Received 13th Sep 2021; accepted 15th Feb 2022)

Abstract. The study was conducted in three Forest Divisions i.e. Solan, Nahan and Rajgarh of Himachal Pradesh, India to determine the impact of controlled burning on the diversity of shrubs and herbs in chir pine forests. A total 2.00 ha was selected for study on each site. The controlled burning was conducted in 1.5 ha (B₁) during January 2017 and 1.0 ha (B₂) in January 2018. A 0.50 ha was designated for control treatments on each site. The field observations of phytosociology were recorded during November, 2018. A total number of 49 species was found at all three sites. The number of species varied from 27 at Lawasachowki (B₂) to 35 at Mangotimor (C). The diversity of shrubs ranged from 2.17 at Bagpashog (B₂) to 2.66 at Mangotimor (C), whereas for herbs it ranged from 2.82 at Mangotimor (C) and Lwasachowki (C) to 3.25 at Bagpashog (B₂). The density of shrubs and herbs of control (C) were significantly different from burning treatments (B₁ and B₂). The density and diversity of shrubs were found higher in control as compared to burning treatments on each sites and vice versa for herbs.

Keywords: *controlled burning, shrubs, herbs, density, diversity*

Introduction

Himalaya is one of the hotspots of the biodiversity (Myers et al., 2000; Singh et al., 2009) and provides ecological niches for plants and human beings (Sharma et al., 2018). Himalaya provides various ecosystem services, sustain biodiversity, regulate hydrological integrity and maintain slope stability (Chawla et al., 2008; Sharma et al., 2009). The plant diversity and communities in Himalaya changes with environmental gradient and also due to variation in geological conditions, topography, river systems, rainfall, altitude and climate (Sharma et al., 2014). The climate change, human and cattle population pressure, over grazing, habitat loss (Sharma et al., 2014) and development of infrastructures like roads, are the major threats to Himalayan ecosystems leading to increased rate of ecological degradation and disturbance to the equilibrium of Himalayan mountain ecosystem (Ahmad et al., 1990). Forest fire is one of the major challenges and requires proper planning of land use management in the region (Avesani et al., 2000).

The forest fire impacts the ecosystems through variety of ways as biogeochemical cycling, atmospheric chemistry and ecosystem structures (Joseph et al., 2009) and also modifies the atmosphere by mixing the greenhouse gases (Simmonds et al., 2005; Chiriaco et al., 2013) and contribute in acceleration of climate change (Stocks et al., 1998; Kiem et al., 2006; Gavin et al., 2007). The forest fires destroy the biomass (Chiriaco et al., 2013; Jharya et al., 2014) and impacts nutrients availability in the soil (Verma et al., 2019) resulting into the accelerating the soil erosion on the slopes in the

Himalaya. The forest fire contributes for altering the plant communities (Whelan, 1995; Arevalo, 2014) changes the dynamics and structure of vegetation of landscape (Baeza, 2007; Arevalo, 2014). The woody and non woody vegetation are prone to forest fire which lead to decrease in the moisture content on plants (West, 1965; Gandiwa, 2011) leading to changes into the functioning of the plants. The forest fire can be prevented by improving awareness among the people about damages caused by the fire and also by employing the legal enforcement and application of suitable silviculture practices such as fire lines and controlled burning (Bhardwaj and Narkhede, 1997).

Chir pine (*Pinus roxburghii* Sarg.) forests in Western Himalayas are prone to forest fire with recurrent fire incidences (Negi et al., 2016). Fire in chir pine forest influence plant community structure and also causing huge economic and environmental losses (Negi et al., 2016; Konsam et al., 2017; Sannigrahi et al., 2020). The fire affected area in the chir pine forest constitutes 2.36% of forest area of India (FSI, 2019). The incidences of the forest fire is decreased by the old age practice of controlled burning during winter when surface temperature is low (Anderson, 1983; Rego et al., 1991; Brose et al., 1998; Blake et al., 2000; Allain and Grace, 2001; Hutchinson et al., 2005; Baeza et al., 2007; Knapp et al., 2007; Gairola et al., 2008; Brose, 2010; Gandiwa, 2011; Balch et al., 2013; Arevalo et al., 2014; Edgar and Griscom, 2017; Kumar, 2019; Kumar et al., 2020; Alba et al., 2021; Jaffe et al., 2021; Jang et al., 2021; Perles et al., 2021). However, succinct analysis and authenticated database on impact of controlled burning on woody and non woody vegetation in Chir pine forests is not available inspite of the fact that the controlled burning can assist to decrease the chances of forest fire in chir pine forest (Kumar et al., 2020).

The lack of information about the impacts and control of forest fire undermine the efforts of the forest department to control the recurrent fire. Moreover, forest fire leads huge losses to exchequer leading to loss of properties and lives (Negi et al., 2016). The impact of controlled burning on understorey of forests is poorly known, therefore, controlled burning evaluations are a requirement for the effective management of vegetation in the forest ecosystems. Moreover, under the current global changes, the evaluations of impacts of forest fire would guide for fire control mechanism and also led to refrain the losses of properties and lives.

With this in view, the study hypothesizes that the controlled burning affects the density and diversity of the forest fire fuel specially the shrubs and herbs of the forest ecosystems. Therefore, objective of the present study was to analyze the impact of controlled burning conducted twice on phytosociology of understory vegetation in chir pine forests. The result of the study would supplement to the current understanding about the forest fire control mechanism along with generation of specific knowledge about forest fire management.

Material and methods

Selection of sites

The study focuses the Indian Himalayan forests of the Nahan and Solan Forest Circles of Himachal Pradesh (HP), India. The focus of the study was higher altitude zone (>1200 m above mean sea level) in Upper Himalayan Chir pine Forests (Group 9/C1b as per Champion and Seth, 1968 classification of Forests of India). Total three sites i.e., Mangotimor, Bagpashog and Lawasachowki were selected randomly for investigations in the forests (*Table 1, Figs. 1 and 2*) in Parwanoo, Sarahan and Jamta

Forest Ranges, respectively. The monthly weather details of the areas are reported in *Figs. 3 and 4*. The dominant species at study sites were *Pinus roxburghii* with associate species as *Quercus leucotrichophora*, *Myrica esculenta* and *Pyrus pashia*. The climate of the study sites was transition between sub-temperate and sub-tropical. The winter, summer and rainy seasons are found in the area and the precipitation in the region is received in the form of the rains. The calcareous shales, carbonaceous shales, dolomite limestone are found on the ground of chir pine forests (Gupta et al., 2009). The depth of the soil in the study sites varied from 40 cm to 70 cm. The slope of the sites ranged from 10° to 25°.

Table 1. Site details of the selected sites in Nahan and Solan Circles of Himachal Pradesh Forest Department

Name of the site	Name of Forest Division	Geo-coordinate		Elevation (amsl)	Aspect
		Latitude (N)	Longitude (E)		
Mangotimor	Solan	30° 53' 49.3"	077° 00' 06.5"	1550m	Sothern Eastern
Bagpashog	Rajgarh	30° 45' 41.5"	077° 09' 52.6"	1522m	North West
Lawasachowki	Nahan	N30° 40' 32.8"	E077° 11' 45.7"	1344m	Sothern Eastern




Figure 1. Map of selected study sites in Solan and Sirmour Districts in Himachal Pradesh




Figure 2. Photographs of the study sites, a. Mangotimor b. Bagpashog and c. Lwasachowki




Figure 3. Monthly average humidity (%) and total rainfall (mm) from January 2017 to December, 2018




Figure 4. Monthly max. temp (°C) and min. temp (°C) from January 2017 to December, 2018

Fire management treatment

The forest fire can be managed by using controlled burning. Controlled burning is controlled firing of forest fuel either in their natural or modified state under specific environmental condition to fulfill the objective of plan resource management (Bhardwaj and Narkhede, 1997). The three treatments for forest fire evaluations were control (C), once burnt (B_1) and twice burnt (B_2). A total area of 2.00 ha homogenous patch of forests on each site was selected for the study and out of which the controlled burning (B_1) was conducted once in 1.50 ha on each site with 0.50 ha designated as control (C)

during January, 2017. In January 2018, controlled burning was again conducted only in 1.00 ha out of the burnt area in 2017 on each site and considered as twice burnt (B_2), and remaining 0.50 ha of the burnt area in 2017 remains intact and treated as B_1 . The designated control site of 0.50 of 2017 ha was control (C) on each site i.e. controlled burning was not undertaken on each sites in the area.

Collection of data

Total 20 quadrates for shrubs and 40 quadrates for herbs on each site were laid out randomly in control (C), once burnt (B_1) and twice burnt (B_2) areas. The size of the quadrates was 5 m x 5 m for shrubs and 1mx1m for herbs (Mishra, 1968). The number of individual and diameter of shrubs, saplings of trees and herbs including natural regeneration of trees for phyto-sociological analysis were recorded in 0.50 ha, 0.50 ha and 1.00 ha for C, B_1 and B_2 , respectively during November, 2018.

Analysis of data

The density (D), frequency (F %), abundance (A) and importance value index (IVI) were determined as per Curtis and McIntosh, 1950. The sum total of relative value of density, basal area, and frequency was used for calculating IVI. Abundance to frequency ratio (A/F) was calculated and the distribution pattern (A/F) was used to categories as contiguous (>0.050), random (0.025-0.050) and regular (<0.050) (Curtis and Cottam, 1956). The Shannon-Wiener Diversity Index (H) was calculated for determining the plant diversity (Eq. 1) (Shannon-Wiener, 1963).

$$H = - \sum_{i=1}^S (ni/N) \ln (ni/N) \quad (\text{Eq.1})$$

Concentration of dominance (Cd) was measured by Simpson's Index (Eq. 2) (Simpson, 1949).

$$Cd = \sum_{i=1}^S (ni/N)^2 \quad (\text{Eq.2})$$

where ni = Importance value of species i and N = total importance value of all the species in both the indices.

Statistical analysis

The effect of controlled burning on number of individuals of shrubs and herbs were evaluated as per the analysis of variance (ANOVA) with 3 treatments i.e. control (C), once burnt (B_1) and twice burnt (B_2). The mean of the four quadrates selected randomly from each patch i.e. for the three treatments was considered as one replication for herbs, whereas mean of two quadrates was treated as replication for shrubs. Total replications were ten for each treatment i.e., control (C), once burnt (B_1) and twice burnt (B_2) for shrubs and herbs on each site. The Least Significant Difference (LSD) test was applied for evaluation of critical differences between two treatment means.

Results

The phytosociological analysis was conducted for collected field data. The differences in the impacts of the treatments were compared through ANOVA. The details of the analysis have been described in subsequent paragraphs for each site for shrubs and herbs.

A total of 49 species was found across the three sites which belong to 49 genera and 28 families. The number of species was 45 at C, 49 at B₁ and 45 at B₂. A total number of species in control was 28 at Lawasachowki and 32 at Mangotimor. The total number of species in once burnt (B₁) varied from 28 (Lawasachowki) to 35 (Mangotimor), whereas in twice burnt (B₂), the number of species ranged from 27 (Lawasachowki) to 33 (Mangotimor).

Carissa carandas, *Myrsine africana*, *Rubus ellipticus* and saplings of *Pinus roxburghii* were present across three sites at all three treatments. *Buddleja asiatica* was present only in B₂ at Mangotimor (Table 2). The saplings of *Myrica esculenta* was present only in C at Bagpashog. The saplings of *Pistacia integerimma* and *Rosa moschata* were present in C and B₁ at Mangotimor. The saplings of *Punica granatum* at Mangotimor were present in B₁ respectively. The saplings of *Sapium insigne* were present only in C and B₂ at Lawasachowki (Table 2).

Table 2. Density/25 m² of shrubs under control (C) and burnt (B₁ and B₂) sites in Himalayan Chir Pine Forests

S.No.	Name of species	Mangotimor			Bagpashog			Lawasachowki		
		C	B ₁	B ₂	C	B ₁	B ₂	C	B ₁	B ₂
1	<i>Berberis lycium</i>	1.05	0.95	0.5	1.5	1	0.75			0.75
2	<i>Buddleja asiatica</i>			0.4						
3	<i>Carissa carandas</i>	3.7	1.3	0.75	0.95	0.75	0.4	1.75	1.25	0.65
4	<i>Colebrookea oppositifolia</i>	0.9		0.4	1	0.8	0.65	0.9	1.2	0.5
5	<i>Ficus palmata</i> *	0.5	0.55	0.5						
6	<i>Flacourtia indica</i> *	0.6						0.95	0.95	0.25
7	<i>Lantana camara</i>	1.25	0.6	0.85						
8	<i>Maesa indica</i>							1.45	1.45	0.95
9	<i>Mallotus philippensis</i> *					0.35	0.2	0.7	0.6	0.3
10	<i>Murraya koenigii</i>				0.7	1.45	0.45	1.5	1.25	0.95
11	<i>Myrica esculenta</i> *				0.55					
12	<i>Myrsine africana</i>	2.4	2.85	0.65	1.5	1	0.95	1.25	1.1	0.6
13	<i>Pinus roxburghii</i> *	0.45	0.3	0.5	0.4	0.3	0.2	0.6	0.4	0.25
14	<i>Pistacia integerimma</i> *	0.25	0.5							
15	<i>Prinsepia utilis</i>	1.15	1.45	0.4						
16	<i>Punica granatum</i> *		0.3							
17	<i>Pyrus pashia</i> *	0.5	0.65	0.4	0.55	0.4	0.25	1	0.6	
18	<i>Rosa moschata</i>	0.35	0.65							
19	<i>Rubus ellipticus</i>	1.4	1.05	0.5	1	0.6	0.4	1.95	1.1	0.4
20	<i>Sapium insigne</i> *							0.9		0.2
21	<i>Toona ciliata</i> *	0.35		0.25	0.35			0.4	0.5	
22	<i>Woodfordia fruticosa</i>	0.45	0.5	0.6				0.6	0.45	
23	<i>Zanthoxylum armatum</i>	0.75	0.95	0.35	0.9					
TOTAL		16.05	12.6	7.05	9.4	6.65	4.25	13.95	10.85	5.8

* Saplings of trees

The data on groundflora (herbs, seedlings/natural regeneration of trees and shrubs) of burnt and unburnt treatments revealed that *Adiantum lunulatum*, *Anaphalis triplinervis*, *Cheilanthes farinosa*, *Chrysopogon montanus*, *Cirsium wallichii*, *Cissampelos pareira*, *Dicliptera bupleuroides*, *Fragaria vesca*, *Heteropogon contortus*, *Oxalis corniculata*, *Viola serpens* along with seedlings of *Pinus roxburghii* were present in all the treatments and sites (Table 3). The seedlings of *Ficus palmata* was present in B₁ and B₂ only at Bagpashog and Lwasachowki. The seedlings of *Flacourtia indica* were present in B₁ and B₂ at Lwasachowki. The seedlings of *Lantana camara* and *Sapium insigne* were present in B₁ and B₂ at Mangotimor. *Thymus linearis*, *Pteris cretica* and seedlings of *Myrica esculenta* and *Prinsepia utilis* were present in B₁ and B₂ at Bagpashog (Table 3).

Density of shrubs and herbs

A. Study site: Mangotimor

Density (Ind./25 m²) of shrubs

The highest density (3.70) at C was recorded for *Carissia carandas* followed by *Myrsine africana* (2.40) and lowest (0.25) for *Pistacia integerrima* (Table 2). The maximum density (2.85) at once burnt (B₁) was found for *Myrsine africana* followed by *Prinsepia utilis* (1.45) and minimum (0.30) was recorded for *Pinus roxburghii* and *Punica granatum*. On twice burnt (B₂) site, maximum density (0.85) was recorded for *Lantana camara* followed by *Carissia carandas* (0.75) and minimum (0.25) for *Toona ciliata* (Table 2). Significance difference in total number of individuals/25 m² of all shrubs were found between C (16.05), B₁ (12.60) and B₂ (7.05) (Table 4).

Density (Ind./m²) of herbs

The highest density (3.03) at C was recorded for *Carex meiogyna*, followed by *Chrysopogon montanus* (2.55) and lowest (0.08) was recorded for *Solanum xanthocarpum* and *Parthenium hysterophorus* (Table 3). At once burnt (B₁), highest density (5.88) was recorded for *Heteropogon contortus*, followed by *Chrysopogon montanus* (5.85) and lowest (0.18) was recorded for *Parthenium hysterophorus* at Mangotimor (B₁). The highest density (3.90) at twice burnt (B₂) was recorded for *Heteropogon contortus* followed by *Chrysopogon montanus* (3.83) and lowest (0.15) for *Sapium insigne* and *Anaphalis triplinervis* (Table 3). Significance difference in total number of individuals of all herbs/m² was observed between C (12.98), B₁ (36.23) and B₂ (22.93) (Table 4).

B. Study site: Bagpashog

Density (Ind./25 m²) of shrubs

The highest density (1.50) at C was recorded for *Myrsine africana* and *Berberis lycium* followed by *Colebrookea oppositifolia* (1.00), *Rubus ellipticus* (1.00) and lowest (0.35) for *Toona ciliata* (Table 2). The highest density (1.45) at once burnt (B₁) was recorded for *Murraya koenigii*, followed by *Berberis lycium* (1.00), *Myrsine africana* (1.00) and least dominant (0.30) was *Pinus roxburghii*. The maximum density (0.95) at twice burnt (B₂) was recorded for *Myrsine africana* followed by *Berberis lycium* (0.75) and least dominant (0.20) was *Mallotus philippensis* and *Pinus roxburghii* (Table 2). Significance difference in number of individual/25 m² of all shrubs was found between C (9.40), B₁ (6.65) and B₂ (4.25) (Table 4).

Table 3. Density /m² of herbs under control (C) and burnt (B₁ and B₂) sites in Himalayan Chir Pine Forests

S.No.	Name of species	Mangotimor			Bagpashog			Lawasachowki		
		C	B ₁	B ₂	C	B ₁	B ₂	C	B ₁	B ₂
1	<i>Adiantum lunulatum</i>	1.03	1.65	2.03	0.45	0.53	0.73	0.45	0.53	0.73
2	<i>Ageratum conyzoides</i>		3.28	1.13				0.3	0.53	0.65
3	<i>Ajuga bracteosa</i>				0.3	0.53	0.65			
4	<i>Anaphalis triplinervis</i>	0.33	0.43	0.15	0.38	0.65	0.75	0.38	0.65	0.75
5	<i>Artemisia vulgaris</i>				0.23	0.5	0.63			
6	<i>Berberis lycium**</i>		0.7	0.45	0.23	0.73	0.83			
7	<i>Carex meigyna</i>	3.03	3.98	2.25						
8	<i>Carissa carandas</i>	0.2	0.43	0.3				0.23	0.5	0.63
9	<i>Cheilanthes farinosa</i>	0.2	1.73	0.85	0.28	0.55	0.3	0.25	0.73	0.83
10	<i>Chrysopogon montanus</i>	2.55	5.85	3.83	2.3	5.38	4.73	1.78	5.88	4.73
11	<i>Cirsium wallichii</i>	0.2	0.35	0.2	0.23	0.63	0.85	0.23	0.63	0.85
12	<i>Cissampelos pareira</i>	0.15	0.35	0.23	0.13	0.4	0.4	0.23	0.55	0.3
13	<i>Dicliptera bupleuroides</i>	0.23	1.48	0.7	0.4	0.35	0.5	0.38	0.35	0.5
14	<i>Ficus palmata*</i>					0.53	0.55		0.43	0.4
15	<i>Flacourtia indica**</i>								0.53	0.55
16	<i>Fragaria vesca</i>	0.38	0.83	0.63	0.65	0.48	0.58	0.4	0.45	0.58
17	<i>Galium aparine</i>		0.58	0.43	0.38	0.58	0.43			
18	<i>Geranium wallichianum</i>		0.3	0.2	0.18	0.45	0.58	0.08	0.58	0.43
19	<i>Heteropogon contortus</i>	2.28	5.88	3.9	2.3	5.88	5.33	2.3	5.38	5.33
20	<i>Hydrocotyle asiatica</i>	0.4	0.75	0.35						
21	<i>Lantana camara**</i>		4.98	0.65						
22	<i>Leucas lanata</i>		0.45	0.6	0.25	0.73	0.65	0.38	0.73	0.65
23	<i>Mallotus philippensis*</i>							0.18	0.35	0.5
24	<i>Murraya koenigii**</i>							0.38	0.78	0.65
25	<i>Myrica esculenta**</i>					0.33	0.5			
26	<i>Myrsine africana**</i>				0.3	0.58	0.5	0.3	0.45	0.5
27	<i>Oxalis corniculata</i>	1.13	0.28	2.23	0.28	0.35	0.5	0.3	0.7	0.63
28	<i>Parthenium hysterophorus</i>	0.08	0.18	0.45						
29	<i>Pinus roxburghii*</i>	0.13	0.23	0.18	0.2	0.43	0.45	0.23	0.3	0.45
30	<i>Prinsepia utilis**</i>					0.63	0.48			
31	<i>Pteris cretica</i>					0.78	0.63			
32	<i>Pyrus pashia*</i>	0.1	0.53	0.23	0.18	0.45	0.55			
33	<i>Rubia cordifolia</i>				0.18	0.63	0.5			
34	<i>Rubus ellipticus**</i>	0.15			0.25	0.35	0.65	0.2	0.38	0.48
35	<i>Sapium insigne</i>		0.23	0.15						
36	<i>Solanum xanthocarpum</i>	0.08			0.3	0.78	0.5			
37	<i>Sonchus asper</i>				0.28	0.45	0.63	0.23	0.53	0.55
38	<i>Thalictrum foliolosum</i>	0.15	0.43	0.25						
39	<i>Thymus linearis</i>					0.7	0.53			
40	<i>Viola serpens</i>	0.13	0.43	0.6	0.28	0.38	0.63	0.2	0.83	0.8
41	<i>Zanthoxylum armatum**</i>	0.1			0.23	0.53	0.45			
TOTAL		13.03	36.31	22.97	11.17	26.27	25.99	9.41	22.77	22.47

*Natural regeneration of trees and ** Natural regeneration of the shrubs

Table 4. Mean and result of ANOVA for density of shrubs and herbs under control (C) and burnt (B₁ and B₂) treatment in the selected sites in Himalayan Chir Pine Forests

Name of site	Treatment (Mean)				F value		LSD _(0.05)
	C	B ₁	B ₂	SE	Calculated	Critical value	
Shrubs							
Mangotimor	16.05	12.6	7.05	1.222	13.151	3.554	2.568
Bag Pashog	9.40	6.65	4.25	0.73	24.894	3.554	1.53
LawasaChowki	13.95	10.85	5.80	0.75	60.860	3.554	1.57
Herbs							
MangotiMor	12.98	36.23	22.93	2.791	34.919	3.554	5.865
Bag Pashog	11.10	26.175	25.92	2.861	18.201	3.554	6.011
LawasaChowki	9.35	22.70	22.43	1.899	32.269	3.554	3.990

Density (Ind./m²) of herbs

The highest density (2.30) at C was recorded for *Heteropogon contortus* and *Chrysopogon montanus* followed by *Fragaria vesca* (0.65) and least dominant (0.13) was *Cissampelos pareira* (Table 3). The maximum density (5.88) at once burnt (B₁) was recorded for *Heteropogon contortus*, followed by *Chrysopogon montanus* (5.38) and minimum (0.33) for *Myrica esculenta*. The maximum density (5.33) at twice burnt (B₂) was recorded for *Heteropogon contortus* followed by *Chrysopogon montanus* (4.73) and minimum (0.30) for *Cheilanthes farinosa* (Table 3). The number of individuals of all herbs/m² was significantly different between C (11.10), B₁ (26.17) and B₂ (25.92). No significant difference was found between B₁ (26.17) and B₂ (25.92) (Table 4).

C. Study site: Lawasachowki

Density (Ind./25 m²) of shrubs

The highest density (1.95) at C was shown by *Rubus ellipticus* followed by *Carissa carandas* (1.75) and lowest (0.40) for *Toona ciliata* (Table 2). The maximum density (1.45) at once burnt (B₁) was recorded for *Maesa indica* followed by *Carissa carandas* (1.25) and minimum (0.40) for *Pinus roxburghii*. *Maesa indica* (0.95) and *Murraya koenigii* (0.95) at twice burnt (B₂) showed highest value for density (Ind/25 m²) followed by *Berberis lycium* (0.75) and least dominant (0.20) was *Sapium insigne* (Table 2). Significant difference in number of individuals/25 m² of all shrubs was found between C (13.95), B₁ (10.85) and B₂ (5.80) (Table 4).

Density (Ind./m²) of herbs

The maximum density (2.30) at C was recorded for *Heteropogon contortus* followed by *Chrysopogon montanus* (1.78) and minimum (0.08) for *Geranium wallichianum* (Table 3). The highest density (5.88) at once burnt (B₁) was recorded for *Chrysopogon montanus* followed by *Heteropogon contortus* (5.38) and lowest (0.30) for *Pinus roxburghii*. Maximum value of density at twice burnt (B₂) was observed for *Heteropogon contortus* (5.33), followed *Chrysopogon montanus* (4.73) and lowest (0.30) for *Cissampelos pareira* (Table 3). The number of individuals of all herbs/m² between C (9.35) was significantly different from B₁ (22.70) and B₂ (22.43), whereas no significant difference was found between B₁ (22.70) and B₂ (22.43) (Table 4).

Dominance of shrubs and herbs the basis of importance value index (IVI)

A. Dominance of shrubs

Carissa carandas (44.04), *Myrsine africana* (38.56) and *Carissa carandas* (30.84) were dominant species at C, B₁ and B₂, respectively at Mangotimor. *Pinus roxburghii* had the least IVI (10.70) at C and 10.69 at B₁ at Mangotimor. The least dominant was *Toona ciliata* (11.11) at B₂ at Mangotimor (Table 5).

Table 5. Importance value index (IVI) of shrubs under control (C) and burnt (B₁ and B₂) sites in Himalayan Chir Pine Forests

S.No.	Name of species	Mangotimor			Bagpashog			Lawasachowki		
		C	B ₁	B ₂	C	B ₁	B ₂	C	B ₁	B ₂
1	<i>Berberis lycium</i>	18.26	19.04	17.75	41.64	32.88	43.07			32.72
2	<i>Buddleja asiatica</i>			16.28						
3	<i>Carissa carandas</i>	44.04	23.81	30.84	28.91	32.43	32.23	31.06	25.81	29.6
4	<i>Colebrookea oppositifolia</i>	13.06		20.15	30.4	31.2	38.99	15.82	27.56	26.49
5	<i>Ficus palmata</i> *	12.09	14.78	20.4						
6	<i>Flacourtia indica</i> *	13.96						22.82	25.5	21.4
7	<i>Lantana camara</i>	16.24	13.46	25.28						
8	<i>Maesa indica</i>							31.23	34.89	34.53
9	<i>Mallotus philippensis</i> *					23.54	22.98	16.84	18.94	25.74
10	<i>Murraya koenigii</i>				23.37	44.5	36.98	27.98	32.76	33.16
11	<i>Myrica esculenta</i> *				20.87					
12	<i>Myrsine africana</i>	34.89	38.56	27.3	38.12	35.68	45.74	25.6	30.52	31.81
13	<i>Pinus roxburghii</i> *	10.7	10.69	23.67	14.4	32.26	22.61	21.39	18.85	21.32
14	<i>Pistacia integerimma</i> *	15.55	18.02							
15	<i>Prinsepia utilis</i>	21.13	32.64	18.69						
16	<i>Punica granatum</i> *		30.23							
17	<i>Pyrus pashia</i> *	17.79	21.92	21.12	22.86	31.65	26.24	24.5	22.81	
18	<i>Rosa moschata</i>	11.16	17.57							
19	<i>Rubus ellipticus</i>	22.54	25.19	21.57	30.26	35.87	31.17	26.92	24.2	24.84
20	<i>Sapium insigne</i> *							26.23		18.38
21	<i>Toona ciliata</i> *	12.41		11.11	20.15			15.34	19.02	
22	<i>Woodfordia fruticosa</i>	10.76	15.47	21.52				14.28	19.13	
23	<i>Zanthoxylum armatum</i>	25.43	18.63	24.33	29.03					
TOTAL		300	300	300	300	300	300	300	300	300

*Saplings of trees

Berberis lycium (41.64), *Murraya koenigii* (44.5) and *Myrsine africana* (45.74) was dominant species at C, B₁ and B₂, respectively at Bagpashog. The least dominant at Bagpashog was *Pinus roxburghii* (14.40), *Mallotus philippensis* (23.54) at B₁ and *Pinus roxburghii* (22.61) at B₂ (Table 5).

Maesa indica was dominant species in treatments i.e., C (31.23) and B₁ (34.89) at Lawasachowki, whereas *Maesa indica* (34.53) showed highest value of importance value index (IVI) for dominant species at B₂. The least dominant was *Woodfordia fruticosa* (14.28) at C, *Pinus roxburghii* (18.85) at B₁ and *Sapium insigne* (18.38) at B₂ at Lawasachowki (Table 5).

B. Dominance of herbs

Heteropogon contortus showed the highest IVI value (42.87) at Mangotimor (C) and 44.82 at Lawasachowki (B₁) (Table 6). The least dominant at Mangotimor was *Cissampelos pareira* (4.04) at C, *Leucas lanata* (4.53) at B₁, *Cirsium wallichii* (3.48) at B₂. At Lawasachowki, the least dominant was *Geranium wallichianum* (3.46) at C, *Myrsine africana* (6.75) at B₁ and *Geranium wallichianum* (5.32) at B₂ (Table 6).

Table 6. Importance value index (IVI) of herbs under control (C) and burnt (B₁ and B₂) sites in Himalayan Chir Pine forests

S.No.	Name of species	Mangotimor			Bagpashog			Lawasachowki		
		C	B ₁	B ₂	C	B ₁	B ₂	C	B ₁	B ₂
1	<i>Adiantum lunulatum</i>	17.58	16.44	20.08	9.57	12.35	9.87	9.88	14.19	12.51
2	<i>Ageratum conyzoides</i>		14.97	13.24				7.67	7.12	9.67
3	<i>Ajuga bracteosa</i>				7.52	5.98	7.74			
4	<i>Anaphalis triplinervis</i>	6.03	10.23	6.58	17.03	10.01	10.5	36.71	11.61	13.27
5	<i>Artemisia vulgaris</i>				13.99	8.59	11.73			
6	<i>Berberis lycium**</i>		9.27	10.58	5.5	6.68	6.27			
7	<i>Carex meiogyna</i>	39.61	22.12	21.34						
8	<i>Carissa carandas</i>	15.53	12.96	8.45				13.05	10.01	15.1
9	<i>Cheilanthes farinosa</i>	10.24	11.85	9.25	8.03	14.07	7.21	5.96	7.94	7.67
10	<i>Chrysopogon montanus</i>	35.91	26.88	27.59	34.65	37.99	30.9	36.95	42.58	36.86
11	<i>Cirsium wallichii</i>	10.22	7.97	3.48	7.08	7.95	7.96	5.95	9.33	9.76
12	<i>Cissampelos pareira</i>	4.04	9.11	7.49	9.04	5.77	8.62	11.58	16.1	9.35
13	<i>Dicliptera bupleuroides</i>	9.11	12.26	12.49	14.69	7.85	6.11	8.38	9.01	7.57
14	<i>Ficus palmata*</i>					8.22	9.69		9.81	11.06
15	<i>Flacourtia indica**</i>								9.62	12.33
16	<i>Fragaria vesca</i>	11.43	9.99	7.19	12.09	7.18	7.26	14.23	8.55	9.08
17	<i>Galium aparine</i>		12.19	5.82	20.05	8.97	4.31			
18	<i>Geranium wallichianum</i>		6.82	4.3	11.57	7.34	7.26	3.46	10.45	5.32
19	<i>Heteropogon contortus</i>	42.87	29.35	29.35	33.55	35.89	36.03	41.9	44.82	43.11
20	<i>Hydrocotyle asiatica</i>	11.16	8.39	4.7						
21	<i>Lantana camara**</i>		20.53	9.35						
22	<i>Leucas lanata</i>		4.53	8.58	11.11	6.87	7.22	18.77	8.15	8.9
23	<i>Mallotus philippensis*</i>							10.78	9.42	7.54
24	<i>Murraya koenigii**</i>							12.01	8.93	12.17
25	<i>Myrica esculenta**</i>					6.87	6.08			
26	<i>Myrsine africana**</i>				7.96	8.97	6.08	8.05	6.75	11.56
27	<i>Oxalis corniculata</i>	15.17	4.9	21.03	7.71	8.15	9.05	8.03	9.49	12.17
28	<i>Parthenium hysterophorus</i>	7.09	5.8	11.17						
29	<i>Pinus roxburghii*</i>	13.08	6.54	8.19	9.77	7.25	11.06	15.87	7.87	14.32
30	<i>Prinsepia utilis**</i>					7.95	10.51			
31	<i>Pteris cretica</i>					7.49	9.57			
32	<i>Pyrus pashia*</i>	8.04	9	14.9	5.62	7.75	6.2			
33	<i>Rubia cordifolia</i>				11.57	7.95	9.01			
34	<i>Rubus ellipticus**</i>	6.8			11.11	8.15	9.58	9.28	8.87	13.56
35	<i>Sapium insigne</i>		13.74	18.71						
36	<i>Solanum xanthocarpum</i>	10.81			7.96	7.49	9.05			
37	<i>Sonchus asper</i>				7.71	5.65	9.57	6.1	10.35	7.67
38	<i>Thalictrum foliolosum</i>	6.15	6.76	7.6						
39	<i>Thymus linearis</i>					8.01	9.63			
40	<i>Viola serpens</i>	7.49	7.42	8.54	8.03	7.72	9.57	15.39	19.03	9.44
41	<i>Zanthoxylum armatum**</i>	11.64			7.08	8.89	6.35			
TOTAL		300	300	300	300	300	300	300	300	300

*Natural regeneration of trees and ** Natural regeneration of the shrubs

At Bagpashog, *Chrysopogon montanus* showed highest value of IVI (34.65) at C and 37.99 at B₁, whereas *Heteropogon contortus* (36.03) was dominant species at B₂. The least dominant was *Berberis lycium* (5.5) at C, *Sonchus asper* (5.65) at B₁, *Galium aparine* (4.31) at B₂ at Bagpashog (Table 6).

Distribution pattern

The distribution pattern was contiguous ($A/F > 0.05$) for all the shrubs, saplings and herbs at all three sites.

Concentration of dominance (Cd)

The highest value of Cd for shrubs (Table 7) was 0.117 at Bagpashog (B₂) followed by 0.114 at Bagpashog (B₁) site and lowest (0.075) at Mangotimor (B₂). The highest value of Cd for herbs (Table 8) was 0.075 at Mangotimor (C) preceded by 0.074 at Lwasachowki (C) site and lowest (0.048) at Bagpashog (B₂).

Table 7. Concentration of dominance (C) and diversity index (H) for shrubs under control (C) and burnt (B₁ and B₂) sites in Himalayan Chir Pine Forests

Name of site	Control (C)		Burnt (B ₁)		Burnt (B ₂)	
	Cd	H	Cd	H	Cd	H
Mangotimor	0.079	2.66	0.080	2.58	0.075	2.61
Bagpashog	0.098	2.36	0.114	2.19	0.117	2.17
Lwasachowki	0.082	2.53	0.087	2.46	0.094	2.38

Table 8. Concentration of dominance (C) and diversity index (H) for herbs under control (C) and burnt (B₁ and B₂) sites in Himalayan Chir Pine Forests

Name of site	Control (C)		Burnt (B ₁)		Burnt (B ₂)	
	Cd	H	Cd	H	Cd	H
Mangotimor	0.075	2.82	0.052	3.09	0.054	3.06
Bagpashog	0.055	3.07	0.052	3.22	0.048	3.25
Lwasachowki	0.074	2.82	0.068	2.94	0.063	2.97

Shannon-Wiener index (H)

The highest value of Shannon-Wiener index (H) for shrubs (Table 7) was 2.66 at Mangotimor (C) followed by 2.61 at Mangotimor (B₂) and lowest was 2.17 at Bagpashog (B₂). The highest value of Shannon-Wiener Index (H) for herbs was 3.25 at Bagpashog (B₂) followed by 3.22 at Bagpashog (B₁) and lowest was 2.82 at Mangotimor (C) and Lwasachowki (C) (Table 8).

Discussion

The floristic composition and structure of communities are the main attributes of the forest ecosystems and depend upon environment and anthropogenic activities (Gairola et al., 2008; Shaheen et al., 2012; Bisht and Bhatt, 2013; Dar and Sundarapandian, 2016) like burning. The control burning in forests is a mechanism applied for

management of forest fire (Whelan, 1995; Arevalo et al., 2014) and has been significantly affected by the number of individuals of shrubs and herbs of chir pine forests and depend on time, intensity and way of controlled burning undertaken. The intensity of controlled burning also decides the floristic composition as the low intensity fire substantially changes the understory composition and density (Brown, 1960; Swan, 1970; Nyland et al., 1982; Reich et al., 1990; Blake and Schuetz, 2000). The change in composition of woody plants after burning is observed by Blake and Schuetz (2000). In the study, controlled burning did not completely burn all available fuel loads on the ground rather total number of shrubs varied from 9 at Bagpashog (B₁ and B₂) to 16 at Mangotimor (C). The probable cause may be the associated factors which restrict the complete loss of fuel loads on the site. Moreover, the intensity of burning depends on temperature, wind speed, humidity, aspect, slope, plant type and fuel load (Maclean, 1992; Pyne and Bryant, 1994; Pyne, 1997; Blake and Schuetz, 2000).

The result indicates that the controlled burning has varied impacts on woody vegetation of understory and herbs on the ground floor in chir pine forests. However, literature on the aspects for the chir pine forests is lacking. The consistent pattern for IVI was not observed for shrubs and herbs in all the three treatments and similar as Kumar et al. (2020). Density and diversity of shrubs were higher in control as compared to fire occurrence for both the burnt treatments and similar to Allain and Grace (2001). The new shoots of shrubs and natural regeneration of trees which emerge after controlled burning are easily palatable and grazed by cattle, however, these new shoots attract the herbivores in the burnt sites. The burning with intensity of the grazing affects the natural regeneration of trees and shrubs and ultimately affecting the diversity of woody vegetation of understory. The controlled burning also destroys the woody plants and has also been reported by various workers (Enslin et al., 2000; Gandiwa and Kativu, 2009; Gandiwa, 2011). The re-sprouting of woody vegetation has also been reported for woody plants after burning (Marrinan et al., 2005; Gandiwa, 2011). The higher temperature during burning decreases physiological functions, growth and increases the mortality of seedlings (Smith et al., 2017; Sharma et al., 2020). These are probable reasons of low density and diversity of shrubs and saplings of trees. The establishment of natural regeneration after burning was poor due to grazing pressure on the sites. The density and diversity after burning can be increased by following rotational grazing.

Total number of herbs varied from 21 at Lawasachowki (C) and Mangotimor (C) to 30 at Bagpashog (B₁ and B₂). Total density and diversity of herbs were higher in burnt sites (B₁ and B₂) as compared to C and is supported by various researchers (Elliot et al., 1999; Royo et al., 2010; Kumar et al., 2020). Density of the herbs was increased after controlled burning and may be due to decrease in competition among different species for soil nutrient and increase in solar insolation. The frequent occurrence of forest fire in chir pine forests (Kumar and Thakur, 2008; Singh et al., 2016; Attri et al., 2020) led to adaptation to forest fire. The various researchers have also reported the higher stems per plant in burnt sites as compared to unburnt sites (San Jose and Farinas, 1983; Scholes and Walker, 1993; Enslin et al., 2000). The higher density of herbs also changes or improves the microclimate of burnt sites by reducing soil erosion. The present result reveals that controlled burning significantly affect the number of the individuals of shrubs and herbs. The contiguous distribution pattern was observed for shrubs and herbs and has also been reported by as a common phenomenon in forest (Kershaw, 1973; Singh and Yadava, 1974; Kunhikannan et al., 1998; Gairola et al., 2011; Dar and

Sundarpandian, 2016; Kumar et al., 2020). The variation in distribution pattern depends upon microenvironment and biotic factors (Joshi and Tiwari, 1990; Dar and Sundarpandian, 2016). The various factors like slope, soil type and soil moisture affect abundance and distribution pattern of the species (Nigh et al., 1985; Pallardy et al., 1988; Taft et al., 1995; Blake and Schuette, 2000). The higher value of concentration of dominance for herbs in control sites and shrubs in burnt sites revealed the homogeneous nature of community (Kohli et al., 2004; Kumar et al., 2020). The effect of the burning is mainly determined by intrinsic nature of understory vegetation which tends to regain its original structure (Fernandes and Botelho, 2003; Marino et al., 2010).

Conclusions

The diversity and density of shrubs were higher in control as compared to burnt sites. The diversity and density of herbs were higher in burnt sites as compared to control. The study concludes that controlled burning may be beneficial for forest management as burning may modify the diversity of the species while maintaining the density for the chir pine forests of the Indian Himalayan region as frequent forest fire has been noted in the recent years in the region. The significance of the study is many folds as the comparative analysis reveals the importance of burning for the forest management alongwith generating the information about the controlled burning impact analysis. The study may be applicable at local level however for larger spatial domain needs further evaluation on actual ground level basis. Interestingly, the forest fire occurs in small patch of land but in absence of proper management practices it converts into a large fire incidences leading to huge loss, however checking the fire at local level by controlled burning may reduce the chances of big fire in forests.

The study concludes that the burning has immediate impacts on the diversity and density of the herbs and shrubs of the forest. The recording of the observations on same phytosociological parameter after long duration on these sites may be made to observe the responses and for specific recommendation. Therefore, forest management should be reorient for immediate silvicultural treatment for strengthening and managing the understory.

Acknowledgements. The authors are grateful to Director, HFRI for supporting to conduct the study and also obliged to Indian Council of Forestry Research and Education (ICFRE), Dehradun for funding to execute the study. The authors are also thankful to Himachal Pradesh Forest Department (HPFD) for assisting to conduct the study. The authors are also thankful to staff of Forest Ecology and Climate Change Division of HFRI for assisting in the study.

REFERENCES

- [1] Ahmad, A., Rawat, J. S., Rai, S. C. (1990): An analysis of the Himalayan environment and guidelines for its management and ecologically sustainable development. – Environmentalist 10: 281-298.
- [2] Alba, C. E. D., Hjalten, J., Sjogren, J. (2021): Restoration strategies in boreal forests: Differing field and ground layer response to ecological restoration by burning and gap cutting. – Forest Ecology and Management 494: 119357.
<https://doi.org/10.1016/j.foreco.2021.119357>.

- [3] Allain, L., Grace, J. B. (2001): Changes in density and height of the shrub *Baccharis balimifolia* following burning in coastal tallgrass prairie. – In: Proceeding 17th N.A. Prairie conference, pp. 66-72.
- [4] Anderson, H. W. (1983): Regenerating yellow birch with prescribed fire. – In: Proceedings of the 1982 Society of American Foresters Conference, 19-22 September 1982, Cincinnati, Ohio, Society of American Foresters, Washington, DC, pp. 168-172.
- [5] Arevalo, J. R., Fernandez-Lugo, S., Garcia-Dominguez, C., Naranjo-Cigala, A., Grillo, F., Calvo, L. (2014): Prescribed burning and clear-cutting effects on understory vegetation in a *Pinus canariensis* stand (Gran Canaria). – The Scientific World Journal 2014(3). <https://doi.org/10.1155/2014/215418>.
- [6] Attri, V., Dhiman, R., Sarvade, S. (2020): A review on status, implications and recent trends of forest fire management. – Archives of Agriculture and Environment Science 5(4): 592-602
- [7] Avesani, P., Perini, A., Ricci, F. (2000): Interactive case based planning for forest fire management. – Applied Intelligence 13(1): 41-57.
- [8] Baeza, M. J., Valdecantos, A., Alloza, J. A., Vallejo, V. R. (2007): Human disturbance and environmental factors as drivers of long-term post-fire regeneration patterns in Mediterranean forests. – Journal of Vegetation Science 18(2): 243-252.
- [9] Balch, J. K., Massad, T. J., Brando, P. M., Nepstad, D. C., Curran, L. M. (2013): Effects of high-frequency understorey fires on woody plant regeneration in Southeastern Amazonian forests. – Royal Society Publishing 368(1619): 20120157. <https://doi.org/10.1098/rstb.2012.0157>.
- [10] Bhardwaj, S. D., Narkhede, S. (1997): Guide for Management of Forest Fires. – Dr. Y. S. Parmar University of Horticulture and Forestry. Nauni, Solan (H.P.) 173230. 1-47.
- [11] Bisht, A. S., Bhat, A. B. (2013): Vegetation structure and plant diversity relation in a sub-alpine region of Garhwal Himalaya, Uttarakhand, India. – African Journal of Plant Science 7(8): 401-406.
- [12] Blake, J. G., Schuette, B. (2000): Restoration of an oak forest in east-central Missouri early effects of prescribed burning on woody vegetation. – Forest Ecology and Management 139: 109-126.
- [13] Brose, P. H., Van Lear, D. H. (1998): Responses of hard wood advance regeneration to seasonal prescribed fires in oak-dominated shelterwood stands. – Canadian Journal of Forest Research 28(3): 331-339.
- [14] Brose, P. H. (2010): Long-term effects of single prescribed fires on hardwood regeneration in oak shelterwood stands. – Forest Ecology and Management 260: 1516-1524.
- [15] Brown, J. H. (1960): The role of fire in altering the species composition of forest in Rhode Island. – Ecology 41: 310-316.
- [16] Champion, H. G., Seth, S. K. (1968): A Revised Survey of Forest Types of India. – Managers of Publications, Delhi, 404p.
- [17] Chawla, A., Rajkumar, S., Singh, K. N., Lal, B., Thukral, A. K., Singh, R. D. (2008): Plant species diversity along an altitudinal gradient of Bhabha Valley in western Himalaya. – Journal of Mountain Science 5(2): 157-177.
- [18] Chiriaco, M. V., Perugini, L., Cimini, D., D' Amato, E., Velentini, R., Bovio, G., Corona, P., Barbati, A. (2013): Comparison of approaches for reporting forest fire related biomass loss and green house gas emissions in southern Europe. – International Journal of Wildland Fire 22(6): 730-738.
- [19] Curtis, J. T., McIntosh, R. P. (1950): The interrelations of certain analytic and synthetic phytosociological characters. – Ecology 31(3): 434-455.
- [20] Curtis, J. T., Cottam, G. (1956): Plant ecology work book-laboratory field reference manual. – Burgess Publishing Co., Minnesota, 193p.

- [21] Dar, J. A., Sundarapandian, S. (2016): Patterns of plant diversity in seven temperate forest types of Western Himalaya, India. – Journal of Asia-Pacific Biodiversity 9(3): 280-292.
- [22] Edgar, B. E., Griscom, H. P. (2017): The effect of controlled burns on abundance of woody species in Appalachian pine- Oak forest at buck mountain, West Virginia. – Natural Areas Journal 37(1): 30-38.
- [23] Elliot, K. J., Hendrick, R. L., Major, A. E., Vose, J. M., Swank, W. T. (1999): Vegetation dynamics after a prescribed fire in the southern Appalachians. – Forest Ecology and Management 114: 199-213.
- [24] Enslin, B. W., Potgieter, A. L. F., Biggs, H. C., Biggs, R. (2000): Long-term effects of fire frequency and season on the woody vegetation dynamics of the *Sclerocarya birrea*/*Acacia nigrescens* savanna of the Kruger National Park. – Koedoe 43(1): 27-37.
- [25] Fernandes, P. M., Botelho, H. S. (2003): A review of prescribed burning effectiveness in fire hazard reduction. – International Journal of Wildland Fire 12: 117-128.
- [26] FSI (2019): India State of Forest Report. – Forest Survey of India. Ministry of Environment Forest and Climate Change, Kaulagarh Road, Dehradun.
- [27] Gairola, S., Rawal, R. S., Todaria, N. P. (2008): Forest vegetation patterns along an altitudinal gradient in sub-alpine zone of west Himalaya, India. – African Journal Plant Science 2(6): 42-48.
- [28] Gairola, S., Sharma, C. M., Suyal, S., Ghildiyal, S. K. (2011): Composition and diversity of five major forest types in moist temperate climate of the Western Himalaya, Uttarakhand, India. – Forestry studies in China 13: 139-153.
- [29] Gandiwa, E., Kativu, S. (2009): Influence of fire frequency on *Colophospermum mopane* and *Combretum apiculatum* woodland structure and composition in northern Gonarezhou National Park, Southeast Zimbabwe. – Koedoe 51(1): 685. DOI: 10.4102/koedoe, v51 i1.685.
- [30] Gandiwa, E. (2011): Effects of repeated burning on woody vegetation structure and composition in a semi-arid southern African savanna. – International Journal of Environmental Sciences 2(2): 458-471.
- [31] Gavin, D. G., Hallett, D. J., Hu, F. S., Lertzman, K. P., Prichard, S. J., Brown, K. J., Lynch, J. A., Bartlein, P., Peterson, D. L. (2007): Forest fire and climate change in Western North America: Insights from sediment charcoal records. – Frontiers in Ecology and the Environment 5(9): 499-506.
- [32] Gupta, B., Mehta, R., Mishra, V. K. (2009): Fire ecology of ground vegetation in *Pinus roxburghii* Sargent plantations in North-West Himalayas: Floristic composition and species diversity. – Caspian Journal of Environmental Science 7(2): 71-78.
- [33] Hutchinson, T. F., Sutherland, E. K., Yaussy, D. A. (2005): Effect of repeated prescribed fires on the structure, composition, and regeneration of mixed-oak forest in Ohio. – Forest Ecology and Management 218: 210-228.
- [34] Jaffe, M. R., Collins, B. M., Levine, J., Northrop, H., Malandra, F., Krofcheck, D., Hurteau, M. D., Stephens, S. L., North, M. (2021): Prescribed fire shrub consumption in Sierra Nevada mixed-conifer forest. – Canadian Journal of Forest Research 13 April 2021. <https://doi.org/10.1139/cjfr-2020-0454>.
- [35] Jang, W., Croteau, J. S., Ortega, Y. K., Hood, S. M., Keyes, C. R., Pearson, D. E., Lutes, D. C., Sala, A. (2021): Native and non-native understory vegetation responses to restoration treatments in a dry conifer forest over 23 years. – Forest Ecology and Management 481: 118684.
- [36] Jharya, M. K., Bargali, S. S., Swamy, S. L., Kittur, B., Bargali, K., Pawar, G. V. (2014): Impact of forest fire on biomass and carbon storage pattern of tropical deciduous forest in Bhoramdeo Wildlife Sanctuary, Chattisgarh. – International Journal of Ecology and Environmental Science 40(1): 57-74.
- [37] Joseph, S., Anitha, K., Murthy, M. S. R. (2009): Forest fire in India: A review of the knowledge base. – The Japanese Forest Society and Springer 14: 127-134.

- [38] Joshi, N. K., Tiwari, S. C. (1990): Phytosociological analysis of Woody vegetation along an altitudinal gradient in Gharhwal Himalaya. – Indian Journal of Forestry 13(4): 322-328.
- [39] Kershaw, K. A. (1973): Quantitative and Dynamic Plant Ecology. – Edward Arnold Ltd., London, 308.
- [40] Kiem, A. S., Franks, S., Verdon, D. C. (2006): Climate variability in the land of fire and flooding rain. – The Australian Journal of Emergency Management 21(2): 52-56.
- [41] Knapp, E. E., Schwilk, D. W., Kane, J. M., Keeley, J. E. (2007): Role of burning season on initial understory vegetation response to prescribed fire in a mixed conifer forest. – Canadian Journal of Forest Research 37(1): 11-22.
- [42] Kohli, R. K., Dogra, K. S., Batish, D. R., Singh, H. P. (2004): Impact of Invasive plant on the structure and composition of natural vegetation of Northwestern Indian Himalayas. – Weed Technology 18(sp1): 1296-1300.
- [43] Konsam, B., Phartyal, S. S., Kumar, M., Todaria, N. P. (2017): Life after fire for understory plant community in subtropical Chir Pine Forest of Garhwal Himalaya. – The Indian forester 143(8): 759-766.
- [44] Kumar, R., Thakur, V. (2008): Effect of forest fire on trees, shrubs and regeneration behavior in chirpine forest in northern aspect under Solan forest division, Himachal Pradesh. – Indian Journal of Forestry 31(1): 19-27.
- [45] Kumar, R. (2019): Prescribed burning: An important tool in management of forest resources. – Indian Journal of Forestry 42(2): 123-131.
- [46] Kumar, R., Verma, R. K., Kumari, K., Kumar, V., Gulshan, K. R. (2020): Impact of controlled burning on herbs in high altitude zone of chirpine forests in Solan and Nahan circles, Himachal Pradesh. – International Journal of Bio-resource and Stress Management 11(4): 414-424.
- [47] Kunhikannan, C., Verma, R. K., Verma, R. K., Khatri, P. K., Totey, N. G. (1998): Ground flora, soil microflora and fauna diversity under plantation ecosystem in Bhata Land of Bilaspur, Madhya Pradesh. – Environment and Ecology 16(3): 539-548.
- [48] Maclean, N. (1992): Young Men and Fire. – University of Chicago Press, Chicago, IL, 352p.
- [49] Marino, E., Madrigal, M., Hernando, C., Diez, C., Fernandes, C. (2010): Flammability descriptors of fine dead fuels resulting from two mechanical treatments in shrubland: A comparative laboratory study. – International Journal of Wildland Fire 19: 314-324.
- [50] Marrinan, M. J., Edwards, W., Landsberg, J. (2005): Resprouting of saplings following a tropical rainforest fire in north-east Wueensland, Australia. – Austral Ecology 30(8): 817-826.
- [51] Mishra, R. (1968): Ecology Word Book. – Oxford and IBH, Publication. Calcutta, 244p.
- [52] Myers, N., Mittermeier, R. A., Mittermeier, C. A., Da Fonseca, A. B. G., Kent, J. (2000): Biodiversity hotspots for conservation priorities. – Nature 403: 853-858.
- [53] Negi, M. S., Kumar, A. (2016): Assessment of increasing threat of forest fires in Uttarakhand, using remote sensing and GIS techniques. – Global Journal of Advanced Research 3(6): 457-468.
- [54] Nigh, T. A., Pallardy, S. G., Garrett, H. E. (1985): Sugar maple environment relationships in the River Hills and central Ozark Mountains of Missouri, USA. – The American Midland Naturalist Journal 114: 235-251.
- [55] Nyland, R. D., Abrahamson, L. P., Adams, K. B. (1982): Proceedings of the 1982 society of American Foresters Conference. – 19-22 September 1982, Cincinnati, Ohio, Society of American Foresters, Washington DC, pp. 163-167.
- [56] Pallardy, S. G., Nigh, T. A., Garrett, H. E. (1988): Changes in forest composition in central Missouri, USA, 1968-1982. – The American Midland Naturalist Journal 120: 21-24.
- [57] Perles, S. J., Niu, X. M., Ruth, A. D., Gibbons, L. D. (2021): Initial conditions influence effects of prescribed burns and deer exclosure fences on tree regeneration and understory

- diversity in Appalachian oak-dominated forests. – *Forest Ecology and Management* 495(2): 119353. <https://doi.org/10.1016/j.foreco.2021.119353>.
- [58] Pyne, N. F., Bryant, F. C. (1994): *Techniques for wildlife habitat management of uplands*. – McGraw Hill, New York, NY, 840p.
- [59] Pyne, S. J. (1997): *Fire in America: A cultural history of wildland and rural fire*. – University of Washington Press, Seattle, WA 7(3): 297-299.
- [60] Rego, F. C., Bunting, S. C., da Silva, J. M. (1991): Changes in understory vegetation following prescribed fire in maritime pine forests. – *Forest Ecology and Management* 41(1-2): 21-31.
- [61] Reich, P. B., Abrams, M. D., Ellsworth, D. S., Kruger, E. L., Tabone, T. J. (1990): Fire affects ecophysiology and community dynamics of central Wisconsin oak forest regeneration. – *Ecology* 71: 2179-2190.
- [62] Royo, A. A., Collins, R., Adams, M. B., Kirschbaum, C., Carason, W. P. (2010): Pervasive interactions between ungulate forest herbaceous diversity. – *Ecology* 91: 93-105.
- [63] San Jose, J. J., Farinas, M. R. (1983): Changes in tree density and species composition in a protected *Trachypogon* savanna, Venezuela. – *Ecology* 64: 447-453.
- [64] Sannigrahi, S., Pilla, F., Basu, B., Basu, A. S., Sarkar, K., Chakraborti, S., Joshi, P. K., Zhang, Q., Wang, Y., Bhatt, S., Bhatt, A., Jha, S., Keesstra, S., Roy, P. S. (2020): Examining the effects of forest fire on terrestrial carbon emission and ecosystem production in India using remote sensing approaches. – *Science of The Total Environment* 725. <https://doi.org/10.1016/j.scitotenv.2020.138331>.
- [65] Scholes, R. J., Walker, B. H. (1993): *An African savanna: synthesis of the Nylsvley study*. – Cambridge University Press, Cambridge, 306p.
- [66] Shaheen, H., Ullah, Z., Khan, S. M., Harper, D. M. (2012): Species composition and community structure of western Himalayan moist temperate forests in Kashmir. – *Forest Ecology and Management* 278: 138-145.
- [67] Shannon, C. E., Wiener, W. (1963): *The Mathematical theory of Communication*. – University of Illinois Press, Urbana, U.S.A.
- [68] Sharma, E., Tse-ring, K., Chettri, N., Shrestha, A. (2009): Biodiversity in the Himalayas-trends, perception and impacts of climate change. – IMBC-Plenary Session 1: Climate Change and its Implications for Mountain, International Centre for Integrated Mountain Development, Kathmandu, Nepal.
- [69] Sharma, P., Rana, J. C., Devi, U., Randhawa, S. S., Kumar, R. (2014): Floristic diversity and distribution pattern of plant communities along altitudinal gradient in Sangla valley, Northwest Himalaya. – *The Scientific World Journal* 2014: 264878. <http://dx.doi.org/10.1155/2014/264878>.
- [70] Sharma, L., Samant, S. S., Kumar, A., Lal, M., Devi, K., Tewari, L. M. (2018): Diversity, distribution pattern, endemism and indigenous uses of wild edible plants in Cold Desert Biosphere Reserve of Indian Trans Himalaya. – *Indian Journal of Traditional Knowledge* 17(1): 122-131.
- [71] Sharma, A., Brethaeuer, D. K., McKeithen, J., Bohn, K. K., Vogel, J. G. (2020): Prescribed burning effects of natural regeneration in Pine flatwoods: Implications for uneven-aged stand conversion from a Florida study. – *Forests* 11: 328. doi: 10.3390/f11030328.
- [72] Simmonds, P. G., Mannings, A. J., Derwent, R. G., Ciais, P., Ramonet, M., Kazan, V., Ryall, D. (2005): A burning question. Can recent growth rate anomalies in the greenhouse gases can be attributed to large-scale biomass burning events. – *Atmospheric Environment* 39(14): 2513-2517.
- [73] Simpson, E. H. (1949): Measurement of diversity. – *Nature* 163: 688.
- [74] Singh, J. S., Yadava, P. S. (1974): Seasonal variation in composition, plant biomass and net primary productivity of a tropical grassland at Kurukshetra, India. – *Ecology Monograph* 44(3): 351-376.

- [75] Singh, A., Lal, M., Samant, S. S. (2009): Diversity, indigenous uses and conservation prioritization of medicinal plants in Lahaul valley, proposed Cold Desert Biosphere Reserve, India. – International Journal of Biodiversity Science and Management 5(3): 132-154.
- [76] Singh, R. D., Gumber, S., Tewari, P., Singh, S. P. (2016): Nature of forest fires in Uttarakhand: Frequency, size and seasonal patterns in relation to pre-monsoonal environment. – Current Science 111(2): 398-403.
- [77] Smith, A. M., Talhelm, A. F., Johnson, D. M., Sparks, A. M., Kolden, C. A., Yedinak, K. M., Apostol, K. G., Tinkham, W. T., Abatzoglou, J. T., Lutz, J. A. (2017): Effects of fire radiative energy density dose on *Pinus contorta* and *Larix occidentalis* seedlings physiology and mortality. – International Journal of Wildland Fire 26: 82-94.
- [78] Stocks, B. J., Fosberg, M. A., Lynham, T. J., Mearns, L., Wotton, B. M., Yang, Q., Jin, J-Z., Lawrence, K., Hartley, G. R., Mason, J. A., McKenney, D. W. (1998): Climate change and Forest fire potential in Russian and Canadian Boreal Forests. – Climatic Change 38: 1-13.
- [79] Swan Jr., F. R. (1970): Post-fire response of four plant communities in south central New York State. – Ecology 51: 1074-1082.
- [80] Taft, J. B., Shwartz, M. W., Philippe, L. R. (1995): Vegetation ecology of flatwoods on the Illinoian till plain. – Journal of Vegetation Science 6: 647-666.
- [81] Verma, S., Singh, D., Singh, A. K., Jayakumar, S. (2019): Post-fire soil nutrient dynamics in a tropical dry deciduous forest of Western Ghats, India. – Forest Ecosystems 6(6): 1-9. <https://doi.org/10.1186/s40663-019-0168-0>.
- [82] West, O. (1965): Fire in vegetation and its use in pasture management with special reference to tropical and sub-tropical Africa. – Commonwealth Agricultural Bureau, Farnham Royal, England, 53p.
- [83] Whelan, R. J. (1995): The Ecology of Fire. – Cambridge University Press, Cambridge, UK, 360p.