

SPATIAL AND TEMPORAL DISTRIBUTION CHARACTERISTICS OF PM_{2.5} AND VARIATION FACTORS OF THE AQI IN THE BEIJING-TIANJIN-HEBEI REGION FROM 2015 TO 2018

CHENG, M. Y.^{1,2} – ZHANG, H.¹ – WAN, L. H.^{1,2*}

¹College of Geographical Sciences, Harbin Normal University, Harbin 150025, Heilongjiang, China

²Heilongjiang Provincial Key Laboratory of Geographic Environment Monitoring and Spatial Information Service in Cold Regions, Harbin Normal University, Harbin 150025, Heilongjiang, China

*Corresponding author

e-mail: wanluhe@hrbnu.edu.cn; phone: +86-1321-4618-158

(Received 6th May 2022; accepted 23rd Aug 2022)

Abstract. The current air pollution situation in northern China, especially in the Beijing-Tianjin-Hebei region, is particularly critical. How to prevent and control air quality in the Beijing-Tianjin-Hebei region has become the focus of China's environmental protection departments. Based on the hourly monitoring data from available monitoring stations in the Beijing-Tianjin-Hebei region from 2015 to 2018, The annual pollution frequency of the city was calculated, and the temporal and spatial distribution characteristics of the PM_{2.5} concentration are analyzed in the study area from 2015 to 2018 by using the spatial interpolation method. The impact of various air pollutants and ground meteorological factors on air quality are studied using the Pearson correlation analysis method. The results are as follows: (1) With the increase of time, the pollution situation of PM_{2.5} in the Beijing-Tianjin-Hebei region has steadily improved. (2) PM_{2.5} and PM₁₀ were the decisive pollutants that had the greatest impact on the AQI. (3) The influence of ground pressure and temperature on the PM_{2.5} concentration and the AQI were more closely related than other meteorological factors. These research results help to deepen the understanding and prediction of air quality changes in the Beijing-Tianjin-Hebei region and provide theoretical support for policy-makers to improve air quality in the region.

Keywords: variation trend, multiple factors, spatial interpolation, correlation analysis

Introduction

With increasing awareness of environmental protection, people are paying more attention to air pollution because these problems will not only have an adverse impact on the environment but also damage human health (Kuo et al., 2021). The concentration of fine particles has always been a key indicator of China's environmental quality and a core factor that restricts sustainable development (Qian et al., 2021). Therefore, particulate matter (PM_{2.5}) pollution has become a major and urgent challenge facing China and the focus of the central government (Liu et al., 2021). PM_{2.5} pollution in the environment has been associated with a variety of adverse health effects (Ru et al., 2021). Many studies have reported that long term exposure to fine particulate matter (PM_{2.5}) will increase the risk of chronic obstructive pulmonary disease (COPD) (Bo et al., 2021). Globally, the number of COPD deaths and the daily deaths caused by environmental PM_{2.5} increased by more than 90% from 1990 to 2019 (Xiaorong et al., 2021). Exposure to long term air pollution, especially fine particulate matter, is a contributor to incidence rate and mortality worldwide and is also known as risk factor for coronary heart disease (CAD) and myocardial infarction (MI) (Slawsky et al., 2021).

Moreover, long term PM_{2.5} exposure is closely related to incidence rate and mortality rate of cancer (Pei et al., 2021). Therefore, understanding and analyzing the temporal and spatial distribution characteristics of PM_{2.5} and the relevant influencing factors of the PM_{2.5} concentration will help to deepen people's understanding of PM_{2.5} and provide a reference for formulating air pollution control policies based on health effects. This is of great significance for controlling air pollution and protecting people's health (Yu et al., 2019).

The air quality index (AQI) can undoubtedly reflect the air quality of a region more directly, and its level directly affects the survival of human beings in a region, although the definition and scope of the AQI vary from country to country (Meng et al., 2021). Scholars all over the world have never stopped studying the regional trend changes and influencing factors of the AQI and its pollutants. Dutta et al. (2021) analyzed the air pollution trends and patterns of the three major cities in India under a temporal and spatial framework and pointed out that the seasonal distribution of the AQI showed that the pollution concentration was high in winter. Bhutiani et al. (2021), based on the temporal and spatial changes of air quality in integrated industrial estate, Haridwar, and its surrounding areas, demonstrated the important role of highway transportation in environmental air quality. Farhadi et al. (2018) analyzed the sensitivity of meteorological parameters and the instability index to the carbon monoxide concentration, particulate matter, and the AQI and discovered that particulate matter was the most important influence on the AQI. Mangayarkarasi et al. (2021) utilized PM_{2.5} as the core factor, then, using the seasonal autoregressive comprehensive moving average and Facebook's Prophet database, proposed an AQI prediction model that provides strong support for the AQI prediction. In particular, during the worldwide novel coronavirus pandemic, regulators need to consider PM_{2.5} and PM₁₀ in monitoring ambient air quality so as to prevent potential hazards associated with human exposure (Richard et al., 2021). As the economic and political center of China, the Beijing-Tianjin-Hebei region has a large population density. An analysis of the change trend factors of the AQI in the Beijing-Tianjin-Hebei region not only is conducive for the formulation of environmental protection policies but also plays a reference role in the prevention and control of COVID-19 in the region.

This study set out to: (1) determine the decisive pollutants that affect the AQI and analyze the internal correlations of air pollutants; (2) analyze the temporal and spatial variation law of PM_{2.5} in the Beijing-Tianjin-Hebei region from 2015 to 2018; and (3) explore the important influencing factors of meteorological factors on the AQI and PM_{2.5} concentration changes.

Data and Methods

Study area

The Beijing-Tianjin-Hebei region is located in the north of the North China Plain at a west longitude of 113°7' 20"–119°50'47" and a north latitude of 36°3'–42°36'58" (Fig. 1). It is adjacent to the Yanshan Mountains in the north, the North China Plain in the south, the Taihang Mountains in the west, and Bohai Bay in the east. The terrain in the northwest and north is high, and the terrain in the south and east is relatively flat. The area includes a variety of geomorphic features, but it is still dominated by a plain landform with an altitude of -52 to 2836 m (Nuan et al., 2021). The Beijing-Tianjin-Hebei region is one of the four major industrial zones in China that includes two

municipalities directly under the central government (Beijing and Tianjin) and 13 cities in Hebei Province such as Zhangjiakou, Chengde, Qinhuangdao, Tangshan, Langfang, Baoding, Cangzhou, Shijiazhuang, Hengshui, Xingtai, and Handan. It has an extremely important economic and political status (Meng et al., 2021).




Figure 1. Study area and monitoring station locations

Data sources

The hourly air quality data of the Beijing-Tianjin-Hebei region from 2015 to 2018 included the AQI, PM_{2.5}, O₃, PM₁₀, NO₂, SO₂, and CO. There exist 72 available air quality monitoring points in the study area (*Fig. 1*), and they publish their data on the national urban air quality real-time release platform of the China Environmental Monitoring Station (<http://www.cnemc.cn>). The pollutant concentration data used in the calculation in this paper is the hourly data. The IAQI of single category pollutants is calculated by using the concentration limit of single category pollutants and the concentration of pollutants respectively, and the maximum value in the calculation result is the air quality index of this hour or that day.

The hourly meteorological data observed in the Beijing-Tianjin-Hebei region from 2015 to 2018 included the temperature, ground pressure, relative humidity, two-minute average wind speed, two-minute average wind direction, precipitation, and other parameters. There are 171 meteorological monitoring stations in the study area (*Fig. 1*). The China Meteorological Data Network (<http://data.cma.cn>) Release contains data that are quality controlled. Therefore, actual rate of each data parameter exceeds 99.9%, and the accuracy of the data is near 100%.

Study methods

Correlation analysis

A Pearson correlation analysis is suitable for measuring the degree of correlation between two variables, and this is defined as the quotient of the covariance and standard deviation between two variables. The correlation coefficient is between -1 and 1. When the absolute value is closer to 0, the correlation is weaker. When the absolute value is closer to 1, the correlation is stronger (Yao, 2021). The correlation grades are shown in Table 1.

Table 1. Correlation strength comparison

Correlation coefficient (r)	Degree of correlation between variables
$1.0 \geq r \geq 0.8$	Extremely strong correlation
$0.8 > r \geq 0.6$	Strong correlation
$0.6 > r \geq 0.4$	Moderate correlation
$0.4 > r \geq 0.2$	Weak correlation
$0.2 > r $	Weak correlation, Basically irrelevant

Spatial interpolation method

The spatial interpolation algorithm is based on the data of known sample points and is used to deduce unknown data in the same area. An inverse distance weighted interpolation (IDW) combines the advantages of the multiple regression gradient method and the natural proximity method. It is a global interpolation method. Experiments have demonstrated that the predicted sample values on a continuous surface generated by the inverse distance weighted interpolation method are completely equal to the actual measured sample values. Hence, the IDW is an accurate interpolation method (Rahman and Lateh, 2016).

Results

Subsection

The correlations between the concentrations of various kinds of atmospheric pollutants and air quality are shown in Figure 2.

The AQI had a very significant and very strong positive correlation with PM_{2.5} and PM₁₀, with correlation coefficients of 0.877 and 0.947, respectively. Among the six types of pollutants, PM_{2.5} and PM₁₀ had the greatest impact on the AQI and were determined to be decisive pollutants. With increases in the PM_{2.5} and PM₁₀ concentrations, the air quality became significantly worse.

The correlation coefficient of PM_{2.5} with PM₁₀ with CO passed the significance level test of $P < 0.05$. PM_{2.5} showed a strong positive correlation with PM₁₀ and the CO and NO₂ concentrations, and the correlation coefficients were 0.775, 0.814, and 0.642, respectively.

There was a significant and strong positive correlation between the SO₂ concentration and the NO₂ concentration, and the correlation coefficient was 0.743. There was a significant and strong negative correlation with the O₃ concentration, and the correlation coefficient was -0.777.




Figure 2. Correlations between the atmospheric pollutant concentration and the AQI

There was a significant and strong positive correlation between the NO₂ concentration and the CO concentration, with a correlation coefficient of 0.783, and a very significant and moderate negative correlation with the O₃ concentration, with a correlation coefficient of -0.585. The concentrations of CO and O₃ showed insignificant and moderate negative correlations, respectively. The concentration of O₃ was negatively correlated with the concentration of the other five pollutants, among which the correlation coefficient was the strongest with the concentration of SO₂ and the smallest with the concentration of PM₁₀. The correlation degree between SO₂ and PM₁₀ was the weakest.

The annual temporal and spatial distribution characteristics of PM_{2.5}

The inverse distance weighted spatial interpolation method was used to interpolate the annual average concentration of PM_{2.5} of 72 air quality monitoring stations in the Beijing-Tianjin-Hebei region from 2015 to 2018, and the continuous surface generated by interpolation is shown in Figure 3. From 2015 to 2018, the annual average concentration range of PM_{2.5} in Beijing Tianjin Hebei region was 28–114 ug/m³ (2015, Fig. 3a), 26–101 ug/m³ (2016, Fig. 3b), 25–90 ug/m³ (2017, Fig. 3c), 7–75 ug/m³ (2018, Fig. 3d).

From 2015 to 2018, the southern regions of the Beijing-Tianjin-Hebei region, primarily Baoding, Xingtai, Shijiazhuang, Hengshui, and Handan, had the highest annual average concentration of PM_{2.5} compared with other regions that were seriously affected by fine particles. The central region primarily includes Cangzhou, Tianjin, Langfang, Tangshan, and Beijing. The annual average concentration of PM_{2.5} was low and was less affected by fine particles. Zhangjiakou, Chengde, and Qinhuangdao in the northern region had the lowest annual average concentrations of PM_{2.5} and were less affected by fine particles (Fig. 3).

Generally speaking, the distribution of PM_{2.5} high concentration areas in the Beijing-Tianjin-Hebei region from 2015 to 2018 was basically unchanged, but the highest value of PM_{2.5} concentration showed a decreasing trend with time. In terms of the spatial

distribution, the annual average concentration of PM_{2.5} from high to low showed distribution characteristics of the highest in the south, the second in the middle, and the lowest in the north.




Figure 3. The annual average concentration distribution of PM_{2.5} from 2015 to 2018

Seasonal temporal and spatial distribution characteristics of PM_{2.5}

According to the climate status of the Beijing-Tianjin-Hebei region in China, the PM_{2.5} concentration data of all the air quality monitoring stations in the Beijing-Tianjin-Hebei region from 2015 to 2018 were divided into four seasons according to the time attribute as shown in Table 2.

Table 2. Season determination

Seasons	Spring	Summer	Autumn	Winter
Month	3, 4, 5	6, 7, 8	9, 10, 11	12, 1, 2

The seasonal spatial and temporal distribution of the PM_{2.5} concentration in the Beijing-Tianjin-Hebei region during 2015–2018 is shown in *Figure 4*.




Figure 4. Seasonal spatial distribution of the PM_{2.5} average concentration from 2015 to 2018

In the spring of 2015, the minimum concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 28 µg/m³, and the maximum value was 97 µg/m³. The average concentration of PM_{2.5} was the highest in the south, followed by the middle, and the lowest in the north (*Fig. 4a*). The minimum average concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region in summer was 24 µg/m³, and the maximum value was 91 µg/m³. The atmospheric environment in the southern part of the Beijing-Tianjin-

Hebei region was most seriously affected by the concentration of PM_{2.5}, and the atmospheric environment in the northern portion was least affected by the concentration of PM_{2.5}. The concentration of PM_{2.5} was low in most areas, and the overall air quality of the region was the best (*Fig. 4b*). The minimum average concentration of PM_{2.5} in autumn was 23 ug/m³, and the maximum value was 102 ug/m³. The maximum average concentration of PM_{2.5} had an obvious upward trend compared with summer. The high value area of the PM_{2.5} concentration was primarily distributed in the south, and the PM_{2.5} concentration in the north was the lowest (*Fig. 4c*). In winter, the minimum average concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 27 ug/m³, and the maximum value was 211 ug/m³. The highest PM_{2.5} concentration was primarily distributed in the south, the lowest PM_{2.5} concentration in the north, and the Beijing-Tianjin-Hebei region has the worst air quality in winter (*Fig. 4d*).

In the spring of 2016, the minimum average concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 25 ug/m³, and the maximum value was 80 ug/m³. The atmospheric environment in the south was most affected by the concentration of PM_{2.5}, and the concentration of PM_{2.5} in the north was the lowest (*Fig. 4e*). In summer, the minimum average concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 22 ug/m³, and the maximum value was 70 ug/m³. The high value areas were primarily distributed in Hengshui and Beijing, and the concentrations of PM_{2.5} in the other areas were low (*Fig. 4f*). The minimum average concentration of PM_{2.5} in autumn was 30 ug/m³, and the maximum value was 126 ug/m³. Compared with summer, the maximum value increased by 56 ug/m³, and the minimum value increased by 8 ug/m³. The high P value area was primarily distributed in the south, and the PM_{2.5} concentrations in other areas were low (*Fig. 4g*). The average concentration in winter was 27 ug/m³, and the maximum value was 170 ug/m³. In addition, the highest concentration displayed a rapid upward trend. The high value area was primarily distributed in the south of the Beijing-Tianjin-Hebei region, followed by the central region (*Fig. 4h*).

In the spring of 2017, the minimum concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 27 ug/m³, and the maximum value was 77 ug/m³. The high value area was primarily distributed in Baoding, and the concentrations of PM_{2.5} in Zhangjiakou and Chengde in the north were the lowest (*Fig. 4i*). In summer, the minimum average concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 19 ug/m³, and the maximum value was 73 ug/m³. The high value areas were primarily concentrated in the south, of which the atmospheric environmental pollution in Handan was most seriously affected by the concentration of PM_{2.5}, and the concentration of PM_{2.5} in the north was low (*Fig. 4j*). The minimum average concentration of PM_{2.5} in autumn was 12 ug/m³, and the maximum value was 85 ug/m³. The high value areas were primarily distributed in Handan and Xingtai in the south. The impact of the PM_{2.5} concentration on the atmospheric environment during autumn was significantly more serious than during spring and summer (*Fig. 4k*). In winter, the minimum average concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 25 ug/m³, and the maximum value was 157 ug/m³. Compared with other seasons, the average concentration of PM_{2.5} was the largest in winter, and the pollution degree of the atmospheric environment was also the most seriously affected by the PM_{2.5} concentration. The high value area was primarily distributed in the south, and the PM_{2.5} concentrations in Zhangjiakou and Chengde in the north were the lowest (*Fig. 4l*).

In the spring of 2018, the minimum average concentration range of PM_{2.5} in the Beijing-Tianjin-Hebei region was 9 ug/m³, and the maximum value was 79 ug/m³. The high value areas were primarily distributed in Baoding. The atmospheric environment of Zhangjiakou and Chengde in the north was the least affected by the PM_{2.5} concentration, and the PM_{2.5} concentration was the lowest (*Fig. 4m*). In summer, the minimum average concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 17 ug/m³, and the maximum value was 50 ug/m³. In summer, the concentration of PM_{2.5} in the study area was low, and the maximum value pm significantly (*Fig. 4n*). The minimum average concentration of PM_{2.5} in autumn was 19 ug/m³, and the maximum value was 73 ug/m³. The high value areas were primarily distributed in Handan, Baoding, and Shijiazhuang in the south, and the atmospheric environment in the north was least affected by the PM_{2.5} concentration (*Fig. 4o*). In winter, the minimum average concentration of PM_{2.5} in the Beijing-Tianjin-Hebei region was 26 ug/m³, and the maximum value was 113 ug/m³. In winter, the concentration of PM_{2.5} reached the maximum value in a year, and the high value areas were primarily distributed in the south. Zhangjiakou and Chengde in the north were the least affected by the concentration of PM_{2.5}, and the concentration of PM_{2.5} was the lowest (*Fig. 4p*).

Overall, from 2015 to 2018, the regional distribution of high PM_{2.5} concentrations in each corresponding season in the Beijing-Tianjin-Hebei region was roughly the same, with high concentrations in the south, followed by the middle and low concentrations in the north. With increased time, the pollution during the four seasons eased from 2015 to 2018, and the overall PM_{2.5} concentration decreased. According to the chronological order of different years, the PM_{2.5} concentration was the highest in winter, with the widest impact on the distribution range, and the pollution was the most serious. The concentration of PM_{2.5} was the lowest in summer. The ranking of the PM_{2.5} concentration from low to high was the following: summer < spring < autumn < winter.

The frequency of exceedance of PM_{2.5} concentration in cities

The 24-hour mean value data of PM_{2.5} in each city in the study area from 2015 to 2018 are statistically analyzed and compared with the air quality standard of PM_{2.5} (0-35 µg/m³ is good, 35-75 µg/m³ is moderate, and more than 75 µg/m³ is polluted) specified in 《Ambient Air Quality Standard》 (GB 3095—1996). *Figure 5a,b* is obtained.

Taking Beijing as an example, from 2015 to 2018, the annual frequency of air quality (PM_{2.5}) above moderate level has increased year by year. The annual frequency of good level increases year by year, from 29.32% in 2015 to 34.43% in 2018. The frequency of pollution level days decreases year by year, from 39.18% in 2015 to 20.60% in 2018, with a decrease of 18.58%. The air quality is significantly improved (*Fig. 5*).

Among the cities in Beijing Tianjin Hebei region, the air quality of PM_{2.5} in Hengshui City improved most significantly. The frequency of polluted level days in the whole year decreased by 35.24% from 55.77% in 2015 to 19.76% in 2018, and the good level frequency also increased from 36.26% to 55.35%, an increase of 19.09%. Zhangjiakou is relatively unique, from 2015 to 2018, although the air quality has improved slightly, the change range is the smallest. In 2016, the polluted level frequency even increased in Zhangjiakou, but in 2015, most of the pollution frequencies of other cities in Beijing Tianjin Hebei region were more than 30%. The polluted frequency of Zhangjiakou was 4.93%, and the frequency of good air quality was as high

as 63.84% (Fig. 5). The high quality of air quality and the stability of change, it may benefit from the excellent geographical location of Zhangjiakou in Shankou.




Figure 5(a). Annual frequency of air quality(PM_{2.5}) from 2015 to 2016




Figure 5(b). Annual frequency of air quality(PM_{2.5}) from 2017 to 2018

Influence of meteorological factors on air quality

A correlation analysis between the meteorological factors, air quality, and air pollutants was conducted, and the results are shown in *Figure 6*.




Figure 6. Correlation of the meteorological factors with the AQI and air pollutants

Wind speed had a very significant and very strong negative correlation with SO₂ and NO₂, with correlation coefficients of -0.815 and -0.859, respectively. Wind speed had a significant and strong negative correlation with O₃, with correlation coefficient of -0.918, and had no significant negative correlation with the AQI and other pollutants (Fig. 6). On the whole, the average wind speed in the study area had a negative correlation with the AQI and the concentrations of six pollutants. This means that when the average wind speed was greater, the smaller the concentration of various pollutants, and the better the air quality.

Wind direction had no significant correlation with the AQI and the six types of pollutants. Wind direction had a medium positive correlation with CO, a medium negative correlation with SO₂, and a weak positive correlation with the AQI and other pollutants (Fig. 6). On the whole, the wind direction value is negatively correlated with SO₂ and positively correlated with other air pollutants. The smaller the wind direction (the larger the value means that the wind direction was more east and south) value, the greater the SO₂ content, the smaller the content of other air pollutants, and the smaller the AQI, and the better the air quality. Compared with other pollutant concentrations, the influence of wind direction on SO₂ and CO was more significant.

Temperature had a very significant and very strong negative correlation with the AQI and PM₁₀, and the correlation coefficients were -0.929 and -0.924, respectively. Temperature had a significant and very strong negative correlation with PM_{2.5}, and the correlation coefficient was -0.806, but it had no significant negative correlation with other pollutants (Fig. 6). The research demonstrated that temperature had a negative correlation with the AQI and the concentration of six pollutants. When the temperature increased, this was conducive to the diffusion of pollutants. In addition, the concentrations of various pollutants were reduced, improving air quality.

Humidity had a significant and very strong positive correlation with PM_{2.5}, NO₂, and CO, and the correlation coefficients were 0.812, 0.823, and 0.833, respectively. Humidity had a significant and strong negative correlation with O₃, and the correlation coefficient was -0.790. Humidity had insignificant and medium positive correlations with the AQI, which was related to PM₁₀ (Fig. 6). The results showed that the relative

humidity was positively correlated with the AQI and the concentrations of five pollutants except O₃. This means that the greater the relative humidity, the smaller the O₃ concentration, but the concentration of other pollutants increased, especially the concentrations of PM_{2.5}, NO₂, and CO, which accelerate the accumulation of pollutants and easily cause heavy pollution.

There was a very significant and very strong negative correlation between pressure and the AQI and PM_{2.5}, and the correlation coefficients were -0.894 and -0.868, respectively. There was a significant and strong negative correlation between pressure and PM₁₀, and the correlation coefficient was -0.790. The pressure had a moderate positive correlation with NO₂ and CO, a weak positive correlation with O₃, and a very weak positive correlation with SO₂ (Fig. 6). The results showed that the greater the near surface pressure, the more favorable the diffusion of the pollutant concentration, the lower the concentrations of various pollutants, and the better the air quality.

Precipitation had a very weak positive correlation with the AQI, a weak positive correlation with NO₂ and CO, a moderate negative correlation with O₃, a weak negative correlation with PM_{2.5}, and a very weak negative correlation with PM₁₀ and SO₂. The correlations between precipitation and the AQI and the six types of pollutants were not significant (Fig. 6). The research showed that when the precipitation increased, the increasing and decreasing trends of various pollutants were different. However, generally speaking, the air quality was developing a favorable trend. In general, changes in temperature, humidity, and pressure had the greatest impact on the changes in the PM_{2.5} concentration in the study area, and temperature and air pressure had the greatest contributions to changes in air quality in the study area.

Discussion

Currently, with the rapid industrial development in the world, the concentration of particulate matter and various air pollutants in the air continues to increase. The impact of various pollutants on air quality has always been a focus of world atmospheric environmental researchers. Mavrakakis et al. (2021) studied the effects of early heat wave events on human thermal discomfort and urban air quality in an industrialized plain in the Mediterranean region, and they determined that particulate matter (diameter < 10 µm) had a significant impact on poor air quality. Vehicle emissions, dust events, and the combustion of fossil fuels and other organic compounds can increase the urban PM concentration (He et al., 2021; Grmasha et al., 2021). However, there have been different conclusions regarding the types of primary pollutants in the ambient air due to different regions and seasons. In the autumn of 2014–2018, the ozone concentration in the Pearl River Delta increased sharply and became the primary pollutant in the ambient air (Huang et al., 2021). Liu et al. (2020) determined that in Nanchong City, the PM_{2.5} value showed a downward trend with each year, and PM_{2.5} was the primary atmospheric pollutant. Bogomolova et al. (2021) concluded that PM_{2.5} has the potential to predict novel corona pneumonia. From 2015 to 2018, the impact of PM₁₀ and PM_{2.5} on the AQI in the Beijing-Tianjin-Hebei region was far greater than that of other air pollutants (Fig. 2). Because PM_{2.5} causes more direct harm to the human body, the primary pollutant affecting air quality in the Beijing-Tianjin-Hebei region was identified as PM_{2.5}, which is understandable because the Beijing-Tianjin-Hebei region is one of China's four major industrial zones.

The large-scale temporal and spatial distribution characteristics of PM_{2.5} not only represent the temporal and spatial distribution characteristics of the AQI to a certain extent but also more intuitively reflect the harm caused by regional atmospheric environmental changes to the human body. Wang et al. (2019) and others used the PM_{2.5} data of 338 cities in China from 2014 to 2017 for real-time monitoring, and they found that the annual average value of the PM_{2.5} concentration decreased annually. However, more than two thirds of cities still exceeded the standard value specified by the Chinese government (35 µg/m³). The high concentration of PM_{2.5} was primarily distributed in the Henan Province in the middle of east China (including Shandong, Jiangsu, Anhui, and other provinces) and the Beijing-Tianjin-Hebei region (Li et al., 2019; Zhao et al., 2021). The concentrations of PM_{2.5} in northern and eastern China were higher than that in southern and western China (Zhou et al., 2019), and the primary air pollutant in northern China was also PM_{2.5} (Huimin et al., 2021). There were obvious regional differences in the urban air quality. The average annual concentration of the AQI and PM_{2.5} showed a ranking of the following: northern cities > southern cities, inland cities > coastal cities (Jia and Ye, 2019). The primary reason for the formation of haze is that the concentration of PM_{2.5} is too high. Determining the spatial heterogeneity of the PM_{2.5} concentration and its influencing factors is of great significance for regional air quality control and management. Zhou et al. (2019) showed that the decreasing trend of the PM_{2.5} concentration in the Beijing-Tianjin-Hebei region from 2015 to 2018 agreed with the change trend of China's air quality this year that showed the distribution characteristics of the highest in the south, the second in the middle, and the lowest in the north (*Fig. 3*). This was also caused by the industrial distribution in the Beijing-Tianjin-Hebei region and the decrease in altitude when the terrain in the region decreased from the north to the south (*Fig. 1*). From 2015 to 2018, most cities in Beijing Tianjin Hebei region showed that the frequency of annual pollution days decreased year by year, and the frequency of annual excellent air quality days increased year by year. The frequency of PM_{2.5} in the whole region was roughly the same. The emission of PM_{2.5} was significantly controlled and improved year by year, and the air quality was getting better and better (*Fig. 5*). Different from other cities, Zhangjiakou has a unique and stable change law, which also confirms that the change trend of PM_{2.5} concentration has spatial differences and regional characteristics. Seasonal variation was very evident in the temporal and spatial variation of the PM_{2.5} concentration. Zhang et al. (2019) showed that the air quality in summer and autumn was better than that in spring and winter by using data from national monitoring stations in 2015. In Tibet, the pollutant concentration in winter was even 38% higher than that in summer (Deqing et al., 2021). There were obvious seasonal differences in the air quality and pollutant concentrations in the Beijing-Tianjin-Hebei region (Cheng et al., 2019). The variation in the PM_{2.5} concentration in Shijiazhuang City in the Beijing Tianjin Hebei region was ranked as the following: winter > autumn > spring > summer (Yue et al., 2021). From 2015 to 2018, the spatial distribution of high PM_{2.5} concentration areas in each season in the Beijing-Tianjin-Hebei region was basically the same. The seasonal change was basically consistent with the national PM_{2.5} change trend and was ranked as the following: winter > autumn > spring > summer (*Fig. 4*).

A study that examines the driving factors of air quality change and PM_{2.5} concentration change is conducive for formulating targeted environmental improvement policies. Secondary aerosols contribute greatly to the PM_{2.5} concentration. Low temperature, low wind speed, and high relative humidity will aggravate the

accumulation of regional pollutants in winter (He et al., 2021). In Chengdu, the air quality had a significant positive correlation with the air temperature, air pressure, visibility, and sunshine hours (Cheng et al., 2018). Sindosi et al. (2021) used the city of Ivanina in northern Greece as an example, and they showed that, in addition to meteorological factors, socioeconomic factors also determined the level of atmospheric particulate matter. Bowen et al. (2021) confirmed that changes in the PM_{2.5} concentration were affected by many factors by using the daily air pollutant concentration and meteorological element data of Lanzhou from 2013 to 2020. By utilizing the air pollution index and surface meteorological elements of the Beijing-Tianjin-Hebei region from 2001 to 2010, Shi et al. (2018) confirmed that meteorological factors had an important impact on air pollution. The novel coronavirus pneumonia outbreak occurred at the end of 2019, and the air quality and PM_{2.5} concentration have been shown to be significantly associated with the spread of novel coronavirus pneumonia. Changes in human activity have had a significant impact on air quality during the novel coronavirus pneumonia period (Bogomolova et al., 2021; Gao et al., 2021). The AQI of most cities decreased significantly in NO₂, SO₂, CO, PM_{2.5}, and PM₁₀, but the change in O₃ was not significant (Fu et al., 2020). The mortality of the novel coronavirus pneumonia was found to be positively correlated with the mean temperature and the AQI. However, mortality was found to be independent of wind speed, relative humidity, and precipitation (Huang et al., 2020). This suggests that the novel coronavirus pneumonia case fatality rate (CFR) may be predicted by PM_{2.5} and other air pollutants (Hou et al., 2020). The increases in temperature and air pressure in the Beijing-Tianjin-Hebei region from 2015 to 2018 led to a significant decrease in the air quality index in the study area. Wind speed, wind direction, humidity, and precipitation had no significant impact on the air quality index in the study area. The change in the PM_{2.5} concentration in the study area had strong correlations with the temperature, humidity, and air pressure (*Fig. 6*), and the variation in the regional temperature and atmospheric pressure played an important role in the regional air quality trend.

Conclusions

(1) PM_{2.5} and PM₁₀ had the greatest impact on the air quality index and were decisive pollutants. PM₁₀ and CO had the greatest impact on PM_{2.5}, and CO had the greatest impact on PM₁₀.

(2) From 2015 to 2018, the overall air quality of the Beijing-Tianjin-Hebei region obviously recovered and developed in a good direction. From the perspective of the spatial distribution, the PM_{2.5} high value area was primarily concentrated in the south, and the areas least affected were primarily concentrated in Chengde and Zhangjiakou. Overall, the annual average concentrations of PM_{2.5} areas from low to high in the study area were ranked in the following order: north < central < south.

(3) From 2015 to 2018, the concentration of PM_{2.5} decreased during each season, but the spatial distribution of the high value area of the PM_{2.5} concentration during each season remained basically the same. The pollution degree and pollution range were similar in different seasons, ranking as the following: winter > autumn > spring > summer. There was no significant difference in the regional distribution characteristics of the high PM_{2.5} value during the same season.

(4) From 2015 to 2018, the PM_{2.5} air quality of cities in Beijing Tianjin Hebei region improved year by year, the frequency of annual pollution days decreased year by year, and the frequency of annual high-quality air quality days increased year by year.

(5) An increase in temperature and air pressure significantly reduced the air quality index in the study area, and the air quality improved. The influences of wind speed, wind direction, humidity, and precipitation on the air quality index in the study area were not significant. However, a change in the PM_{2.5} concentration in the study area had a strong correlation with temperature, humidity, and air pressure. An increase in temperature and air pressure and a decrease in humidity led to a significant decrease in the air quality index in the study area, and the air quality developed a good trend. Changes in wind speed, wind direction, and precipitation had no significant correlations with changes in the PM_{2.5} concentration in the study area.

REFERENCES

- [1] Bhutiani, R., Kulkarni, D. B., Khanna, D. R., Tyagi, V., Ahamad, F. (2021): Spatial and seasonal variations in particulate matter and gaseous pollutants around integrated industrial estate (IIE), SIDCUL, Haridwar: a case study. – *Environment, Development and Sustainability* 23: 15619-15638.
- [2] Bo, Y., Chang, L-Y., Guo, C., Lin, C., Lau, A. K. H., Tam, T., Lao, X. Q. (2021): Reduced ambient PM_{2.5}, better lung function, and decreased risk of chronic obstructive pulmonary disease. – *Environment International* 156: 10676.
- [3] Bogomolova, A., Balk, I., Polkovnikova, N., Ivaschenko, N. (2020): The impact of the COVID-19 related lockdowns on air quality. – *International Conference on Sustainability and Climate Change* 534: 012010.
- [4] Bowen, C., Yuxia, M., Fengliu, F., Yifan, Z., Jiahui, S., Hang, W., Yongtao, G., Yifan, C. (2021): Influence of weather and air pollution on concentration change of PM_{2.5} using a generalized additive model and gradient boosting machine. – *Atmospheric Environment* 255: 118437.
- [5] Cheng, Y., Zhang, D., Yang, R., Ni, C., Sun, L. (2018): A Study on the Relationship between Meteorological Condition and Air Quality of Chengdu. – *Open Journal of Nature Science* 6(01): 53-62.
- [6] Cheng, Y., Liu, T., Zhao, Y., Wang, Y. (2019): Spatiotemporal Evolution and Socioeconomic Driving Mechanism of Air Quality in Beijing-Tianjin-Hebei and Surrounding Areas ("2+26" Cities). – *Economic Geography* 39(10): 183-192.
- [7] Deqing, Z., Tang, S., Ren, C., Da, Q. (2021): Analysis of the Air Pollution Index and Meteorological Factors and Risk Assessment for Tibet. – *Journal of Physics: Conference Series* 1838(1): 012047.
- [8] Dutta, S., Ghosh, S., Dinda, S. (2021): Urban Air-Quality Assessment and Inferring the Association Between Different Factors: A Comparative Study Among Delhi, Kolkata and Chennai Megacity of India. – *Aerosol Science and Engineering* 5(1): 93-111.
- [9] Farhadi, R., Hadavifar, M., Moeinaddini, M., Amintoosi, M. (2018): Sensitivity Analysis of Meteorological Parameters and Instability Indices on Concentration of Carbon Monoxide, Particulate Matter, and Air Quality Index in Tehran. – *ECOPERSIA* 6(2): 91-100.
- [10] Fu, F., Purvisroberts, K. L., Williams, B. (2020): Impact of the COVID-19 Pandemic Lockdown on Air Pollution in 20 Major Cities around the World. – *Atmosphere* 11(11): 1189.
- [11] Gao, C., Li, S., Liu, M., Zhang, F., Cai, C. (2021): Impact of the COVID-19 Pandemic on Air Pollution in Chinese Megacities from the Perspective of Traffic Volume and Meteorological Factors. – *Science of The Total Environment* 773(1-4): 145545.

- [12] Grmasha, R. A., Al-Azzawi, S., Al-Sareji, O. J., Alardhi, T., Alkhayyat, A. (2021): Engineering, Analysis of Air Quality Index Distribution of PM 10 and PM 2.5 Concentrations in Ambient Air of Al-Hillah City, Iraq. – IOP Conference Series Materials Science and Engineering 1058(1): 012014.
- [13] He, C., Hong, S., Mu, H., Tu, P., Huang, J. (2021): Characteristics and Meteorological Factors of Severe Haze Pollution in China. – Advances in Meteorology 2021: 1-15.
- [14] Hou, C. K., Qin, Y. F., Liu, Q. L., Yang, X. Y., Wang, H. (2020): Impact of Long-Term Air Pollution on the Case Fatality Rate of COVID-19. – Research Square.
- [15] Huang, H., Liang, X., Huang, J., Yuan, Z., Ouyang, H., Wei, Y., Bai, X. (2020): Correlations between Meteorological Indicators, Air Quality and the COVID-19 Pandemic in 12 Cities across China. – Journal of Environmental Health Science and Engineering 18(2): 1491-98.
- [16] Huang, C., Liao, Q., Lu, J. (2021): Analysis of Temporal and Spatial Variation Characteristics of Air Quality in the Pearl River Delta during 2015-2019. – IOP Conference Series Earth and Environmental Science 687(1): 012026.
- [17] Huimin, Z., Rende, W., Xiaomiao, F., Cheney, A., Jinping, Q. (2021): Spatial characteristics of the PM_{2.5}/PM₁₀ ratio and its indicative significance regarding air pollution in Hebei Province, China. – Environmental Monitoring and Assessment 193: 486.
- [18] Jia, Q., Ye, C. (2019): Canonical Correlation Analysis of Composition of Urban Development Land and Air Quality in 35 Large and Medium-sized Cities in China. – Chinese Agricultural Science Bulletin 35(25): 84-93.
- [19] Kuo, C. H., Hsiu, L. Y., Yen, L. C. (2021): Convex nonparametric least squares and stochastic semi-nonparametric frontier to estimate the shadow prices of PM_{2.5} and NO_x for Taiwan's transportation modes. – Ecological Indicators 15(9): 659-677.
- [20] Li, D., Zhao, Y., Wu, R., Dong, J. (2019): Spatiotemporal Features and Socioeconomic Drivers of PM_{2.5} Concentrations in China. – Sustainability 11(4): 1201.
- [21] Liu, C. L., Guo, C. H., Jiang, Z. B., Chen, L. Y. (2020): Characteristics and Time Series Analysis of Primary Air Pollutants in Nanchong City from 2014 to 2018. – Sichuan Environment 39(02): 42-48.
- [22] Liu, G., Dong, X., Kong, Z., Dong, K. (2021): Does national air quality monitoring reduce local air pollution? The case of PM_{2.5} for China. – Journal of Environmental Management 296(4): 113232.
- [23] Mangayarkarasi, R., Vanmathi, C., Khan, M. Z., Noorwali, A., Agarwal, P. (2021): COVID19: Forecasting Air Quality Index and Particulate Matter (PM_{2.5}). – Computers, Materials and Continua 67(3): 3363-3380.
- [24] Mavrikis, A., Kapsali, A., Tsiros, I. X., Pantavou, K. (2021): Air quality and meteorological patterns of an early spring heatwave event in an industrialized area of Attica, Greece. – Euro-Mediterranean Journal for Environmental Integration 6: 25.
- [25] Meng, J., Wang, M., Xuekelaiti, X. (2021): Characteristics of air pollution and environmental economic efficiency in Beijing-Tianjin-Hebei and surrounding areas. – Arabian Journal of Geosciences 14(12)
- [26] Nuan, W., Jie, Y., Lin, Z., Yanbing, W., Zhengyang, H. (2021): Spatial Downscaling of Remote Sensing Precipitation Data in the Beijing-Tianjin-Hebei Region. – Journal of Computer and Communications 09(06): 191-202.
- [27] Pei, Y., Xu, R., Coelho, M. S. Z. S., Saldiva, P. H. N., Li, S., Zhao, Q., Mahal, A., Sim, M., Abramson, M. J., Guo, Y. (2021): The impacts of long-term exposure to PM_{2.5} on cancer hospitalizations in Brazil. – Environment International 154: 106671.
- [28] Qian, L., Zheyu, Z., Chaofeng, S., Run, Z., Yang, G., Chen, C. (2021): Spatio-temporal variation and driving factors analysis of PM_{2.5} health risks in Chinese cities. – Ecological Indicators 129: 107937.

- [29] Rahman, M. R., Lateh, H. (2016): Meteorological drought in Bangladesh: assessing, analysing and hazard mapping using SPI, GIS and monthly rainfall data. – *Environmental Earth Sciences* 75: 1026.
- [30] Richard, G., Sawyer, W. E., Izah, S. (2021): Health Implications of Air Quality Index of Fine and Coarse Particulates During Outdoor Combustion of Biomass in the Niger Delta, Nigeria. – *International Journal of World Policy and Development Studies* 72: 21-26.
- [31] Ru, M., Brauer, M., Lamarque, J.-F., Shindell, D. (2021): Exploration of the Global Burden of Dementia Attributable to PM_{2.5}: What Do We Know Based on Current Evidence? – *GeoHealth* 5(5): e220GH000356.
- [32] Shi, H., Critto, A., Torresan, S., Gao, Q. (2018): The Temporal and Spatial Distribution Characteristics of Air Pollution Index and Meteorological Elements in Beijing, Tianjin, and Shijiazhuang, China. – *Integrated Environmental Assessment and Management* 14(5): 710-721.
- [33] Sindosi, O. A., Hatzianastassiou, N., Markozannes, G., Rizos, E. C., Ntzani, E., Bartzokas, A. (2021): PM₁₀ Concentrations in a Provincial City of Inland Greece in the Times of Austerity and Their Relationship with Meteorological and Socioeconomic Conditions. – *Water Air and Soil Pollution* 232: 77.
- [34] Slawsky, E., Ward-Caviness, C. K., Neas, L., Devlin, R. B., Cascio, W. E., Russell, A. G., Huang, R., Kraus, W. E., Hauser, E., Diaz-Sanchez, D., Weaver, A. M. (2021): Evaluation of PM_{2.5} air pollution sources and cardiovascular health. – *Environmental Epidemiology* 5(3): e157.
- [35] Wang, Y., Duan, X., Wang, L. (2019): Spatial-Temporal Evolution of PM_{2.5} Concentration and its Socioeconomic Influence Factors in Chinese Cities in 2014–2017. – *International Journal of Environmental Research and Public Health* 16(6): 985.
- [36] Xiaorong, Y., Tongchao, Z., Yuan, Z., Hao, C., Shaowei, S. (2021): Global burden of COPD attributable to ambient PM_{2.5} in 204 countries and territories, 1990 to 2019: A systematic analysis for the Global Burden of Disease Study 2019. – *Science of the Total Environment* 796: 148819.
- [37] Yao, W., Gui, K., Wang, Y., Che, H., Zhang, X. (2021): Identifying the dominant local factors of 2000-2019 changes in dust loading over East Asia. – *Science of The Total Environment* 777(D23): 146064.
- [38] Yu, G., Wang, F., Hu, J., Liao, Y., Liu, X. (2019): Value Assessment of Health Losses Caused by PM_{2.5} in Changsha City, China. – *International Journal of Environmental Research and Public Health* 16(11): 2063.
- [39] Yue, T., Qiu, J., Wang, J., Fang, C. (2021): Analysis of Spatio-Temporal Variation Characteristics of Main Air Pollutants in Shijiazhuang City. – *Sustainability* 13(2): 941.
- [40] Zhang, Z., Chen, L., Zhang, H., Liu, P. (2019): The Analysis of Spatial-Temporal Distribution Characteristics and Correlation on Air Quality in Beijing-Tianjin-Hebei Region. – *Geomatics World* 26(03): 31-35.
- [41] Zhao, C., Sun, Y., Zhong, Y., Xu, S., Liang, Y., Liu, S., He, X., Zhu, J., Shibamoto, T., He, M. (2021): Health, Spatio-temporal analysis of urban air pollutants throughout China during 2014–2019. – *Air Quality Atmosphere & Health* 14(10): 1619-1632.
- [42] Zhou, L., Zhou, C., Yang, F., Che, L., Wang, B. (2019): Spatio-temporal evolution and the influencing factors of PM_{2.5} in China between 2000 and 2015. – *Journal of Geographical Sciences* 29(2): 253-270.