

ASSESSMENT OF BIOCHEMICAL MARKERS AND BEHAVIOR RESPONSE OF NON-TARGET FRESHWATER TELEOST *GAMBUSIA AFFINIS* TO CARBOFURAN TOXICITY

ROUACHDIA, R.¹ – TREA, F.² – TICHATI, L.^{2,3} – OUALI, K.^{2*}

¹Department of Natural and Life Sciences, Faculty of Exact Sciences and Natural and Life Sciences, University of Tebessa, Tebessa 12002, Algeria
(e-mail: crouachdia@yahoo.fr)

²Environmental Biomonitoring Laboratory (LBSE), Department of Biology, Faculty of Sciences, Badji-Mokhtar University, BP 12, Annaba 23000, Algeria
(e-mail: fouzia.trea@univ-annaba.dz; <https://orcid.org/0000-0002-9950-8239>)

³Environmental Research Center (ERC), Campus Sidi Amar. BP N° 2024. 23005. Annaba
(e-mail: lazherbio@hotmail.fr; <https://orcid.org/0000-0002-1503-7354>)

*Corresponding author

e-mail: kheireddine.ouali@univ-annaba.dz; <http://orcid.org/0000-0002-5842-0767>

(Received 11th Aug 2022; accepted 11th Nov 2022)

Abstract. The objective of this study is to evaluate the toxicity of a carbamate pesticide, carbofuran, on a non-target species *Gambusia affinis* females. Fish were exposed for 15 and 40 days to two concentrations 0.191 and 0.255 mg/L corresponding respectively ¼ and 1/3 of the LC₅₀ of carbofuran. After parturition, the frequency of malformations in offspring of treated females was evaluated. In addition, we studied the effect of acute toxicity of the carbofuran at 2, 4 and 6 hours on swimming activity, and on brain AChE response. A significant increase in enzymatic activity, and Malondialdehyde (MDA) concentration, along with decreased the activity of acetylcholinesterase (AChE), reduced glutathione (GSH) and total lipid contents in a time - and dose-dependent manner was observed in carbofuran group compared to controls. Acute exposure of *Gambusia* to carbofuran induced a decrease in swimming activity. This behavioral effect was significantly correlated with the decrease in the brain's AChE activity. Exposure of females to relatively low doses of carbofuran also led to a marked increase in the level of larvae skeletal alterations. Hence, the results proved that females strongly accumulate this pesticide during the reproductive cycle, and thus its effect can be transmissible to their offspring.

Keywords: fish, carbamate pesticide, stress biomarkers, brain AChE response, swimming activity, deformities

Introduction

Pesticides are wide-used chemicals in current agriculture activities and are considered one of the main factors involved in environmental pollution in the world (Hamed et al., 2021; Tissot et al., 2022). Aquatic environments are principally susceptible to pesticide pollution following the processes of leaching and runoff and treated water. Seasonal agricultural and weathering activities, including precipitation, photodegradation contribute to an uneven spatial and temporal distribution of pesticide concentrations in the natural water bodies (Amri et al., 2017; Bouzahouane et al., 2018). Despite their selectivity and specific mode of action, pesticides exert their harmfulness towards organisms involuntarily exposed, following the contamination of the environment and the food chain. They are cytotoxic, neurotoxic, embryotoxic, mutagenic, teratogenic, or carcinogenic. The genotoxic effects of these chemicals can go through activation of cell metabolism, and the formation of electrophilic intermediates able to interact with nucleic

acids, or through induction of oxidative stress, inhibition of the cell communication, formation of activated receptors, or others (Abdulelah et al., 2020; Archer et al., 2020; Kasonga et al., 2021).

Carbofuran (2,3-dihydro- 2,2-dimethyl-7-benzofuranyl-methyl carbamate) is a worldwide used chemical and a non-persistent spectrum systemic insecticide, nematicide and acaricide (Mishra et al., 2020). Water bodies can be contaminated with carbofuran due to its groundwater ubiquity index (GUS) of approximately 4.5, which increases their risk of reaching groundwater (Mishra et al., 2020). A set of recent works have been showed the impact of insecticides and in particular of carbofuran on growth (Mansano et al., 2016; Cheghib et al., 2020), on biochemical and hematological parameters of *Clarias batrachus* (Begum, 2008; Narra, 2016) and, on fish acetylcholinesterase in brain and muscle (Rendón-von Osten et al., 2005; Mahboob et al., 2014).

However, there is a lack of information on the genotoxic, teratogenic, and neurobehavioral effects of carbofuran on non-target organisms is available. In fish, nutritional imbalances involving deficiencies of tryptophan, essential fatty acids, and vitamins have been shown to result in skeletal malformations (Roy and Lall, 2007; Garcês et al., 2020; Liu et al., 2020). Environmental factors, like water temperature, heavy metal contamination, or hydrodynamic conditions have been reported to induce marked alterations during development (Sfakianakis et al., 2015; Mendes et al., 2021; Darvishi et al., 2022). Thus, the present study was aimed to assess the toxic effects of sub-chronic exposure to sub-lethal concentrations of carbofuran on the behavioral activity via measurement of swimming and acetylcholinesterase (AChE) activity, and the antioxidant activity evaluation via determination of the major antioxidant markers, including GSH, GST, CAT, and MDA.

Materials and Methods

Site description

Gambusia affinis, is a fish belonging to the family Poeciliidae, it is viviparous of small size varying from 2 to 6 cm. The species studied was fished in the wadi kherraza (36° 53' 60" N and 7° 46' 0.001" E located in Annaba (North-East, Algeria) this area is far from any industrial activity, housing and crop fields.

Biological material

The harvested fish are transported to the laboratory in a cooler. All our experiments were conducted in aquariums with a capacity of 70 liters whose water is dechlorinated and continuously filtered (500 ml/min) and under laboratory conditions, a photoperiod 12L: 12D, temperature 17.5 ± 0.73 °C, pH 7.51 ± 0.33 , dissolved oxygen 13.83 ± 4.23 % and Salinity 239.5 ± 21.12 mg/L. A commercial prepared diet called TETRAMIN (composed of shrimp and dehydrated fish) was given to the experimental fishes.

Experimental protocol

A static nonrenewable toxicity bioassay was conducted according to a standard method APHA-AWWA-WEF (1998) to determine the lethal concentration. The fish were divided into three experimental batches of 60 fish per batch; a control batch composed of untreated fish, two batches treated with carbofuran (99.9% purity, Cluzeau Info Labo, France) dissolved in acetone (20 mg/ml). The carbofuran solution is homogenized in the

aquarium 15 min before the introduction of the fish. The exposure times are 15 d and 40 d at two doses of 0.191 mg/L ($\frac{1}{4}$ LC₅₀) and 0.255 mg/L ($\frac{1}{3}$ LC₅₀) according to tests previously performed using a battery of concentrations from 10 to 100 mg/L. After each exposure period, the hepatopancreas, and brains of the fishes are removed under ice, weighed and stored at -80°.

Preparation of homogenate and measurement of enzyme activities

Hepatopancreas and brains from each batch were pooled at 100 mg and homogenized in saccharose sodium phosphate buffer at pH 6.5. The sample homogenates were centrifuged at 5000 rpm for 15 minutes; the resulting supernatant was recovered for assay of total protein, lipids, malondialdehyde (MDA), reduced glutathione (GSH), GST activity, catalase (CAT), and acetylcholinesterase (AChE).

Metabolite analyses

Protein levels were quantified by the method of Bradford (1976) method using coomassie brilliant blue (BBC). Total lipid levels have been determined as previously described (Goldsworthy et al., 1972) using sulfo-phospho-vanillin assay. The absorbance was read in the dark at 530 nm. Lipid peroxidation was estimated by the method of Draper and Hadley (1990) based on the determination of thiobarbituric acid reactive species (TBARS), using 1, 1, 3, 3 tetra ethoxypropane as standard. The content of MDA was spectrophotometrically determined and expressed as nmol /mg protein.

Biomarkers analysis

The concentration of reduced glutathione (GSH) is quantified by the colorimetric method of Weckberker and Cori (1988), based on measurement of 2-nitro-5-mercaptopuric acid derived from the reduction of 5-5'-dithio-bis-2-nitrobenzoic acid (DTNB) by the SH thiol groups of glutathione measured at a wavelength of 412 nm. Glutathione S-transferase (GST) activity was measured by as described elsewhere (Habig et al, 1974) using a mixture solution of 0.1 M potassium phosphate buffer (pH 6.5) and 1 mM GSH, 1 mM CDNB. The enzymatic activity is displayed in nmol min⁻¹ mg⁻¹ protein. The catalase activity was determined as described by Aebi (1984). The absorbances were measured at 240 nm by the change in optical density following the hydrolysis of hydrogen peroxide (H₂O₂) by reacting 200 µL of H₂O₂ and 20 µL of homogenate in phosphate buffer (100 mM, pH 7.4) for 1 min at an incubation temperature of 25°C. The enzymatic activity was given in nmoles of H₂O₂/ minute/ milligram of protein. Acetylcholinesterase (AChE) activity was evaluated as previously described (Ellman et al., 1961). through recording the sample absorbance after 5 min at 412 nm in the presence of 1mM acetylthiocholine, which would be hydrolyzed to acetic acid and thiocholine providing yellow product named TNB (5-thio-2 nitrobenzoic acid) in the presence of DTNB (5'-dithio-bis-2-nitrobenzoic acid). AChE activity was expressed in nmol min⁻¹ mg⁻¹ protein.

Measurement of swimming activity

The swimming activity of control and treated fish was determined by the method of Bretau et al. (2001), Sousa and Nunes (2020). Visual cues were placed on the aquarium windows to determine 4 observation areas of 40x40x31cm to allow quantification of movements. Each observation involved groups of 5 naïve individuals (tested once). During 6 hours of exposure, fish movements were monitored every 2 hours for a period

of 15 min, with a positioning record every 30 seconds. During the exposure, fish movements were tracked and recorded using high-speed recording system with a camera (Uxsiya, Ref xetwbk62fg-02) and LED lights, which allowed us to accurately calculate the movements of the specimens. The movement activity is given by the number of individuals that moved from one area to another during the 15mn of observation. The concentrations tested and the controls were the same as for the neurochemical activity measurements. Each treatment was carried out five times.

Skeletal deformity

The estimation of skeletal malformations was carried out on aged fry. For the identification of skeletal malformations, the method improved by Staples (1964) based on the use of alizarin red was used. Fry from control and treated females were placed in absolute ethanol (99%) for 48h and then placed in a solution composed of ethanol (85%), KOH (1.5%) and alizarin red (0.0015%) for 5 days. The fry was macerated in an aqueous solution of KOH (1%) until total degradation of the tissues and their skeletons will be clearly visible. The specimens will be preserved in glycerol (50%) for microscopic observation.

Statistical analysis

The results were displayed as means and standard error. We used the two-criteria analysis of variance test "ANOVA" followed by the "Tukey" test to compare the means between them on the one hand and between those of the controls on the other hand. The variations are considered significant when, after this test, the significance has a probability greater than 95% ($p < 0.05$). All tests were performed using Statistica version 10.1 software.

Results

Effect of carbofuran on metabolic parameters (total lipids and MDA)

The results shown in *Figure 1* reveal that treatment of *Gambusia affinis* females with carbofuran induces a significant decrease in hepatopancreatic lipids. This variation was found to be dose and time-dependent ($p < 0.001$). This decrease in lipid levels is associated with lipids peroxidation revealed by an increase in MDA levels after 15 days at $LC_{50} \frac{1}{4}$ and $LC_{50} \frac{1}{3}$ respectively. The ANOVA test shows a highly significant dose and time effect ($p < 0.001$) and a dose-time interaction ($p < 0.001$).

Effect of carbofuran on biomarkers

Reduced glutathione level

Our results show a reduction in hepatopancreatic GSH content in females exposed for 15 and 40 days to both doses 0.191 mg/L and 0.255 mg/L, with reductions ranging from 27% to 33% after 15 days and 37% and 48% after 40 days of treatment in comparison with the control ($p < 0.05$, $p < 0.001$, *Figure 2A*).




Figure 1. Lipids (A) and malondialdehyde (B) content in *Gambusia affinis* hepatopancreas as response to carbofuran exposure. Values are expressed as means $\pm$ SEM ($n = 5$). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs control and # $p < 0.05$ vs days 15




Figure 2. GSH content (nmol/mg protein) and enzymatic activity of GST (nmoles C-DNB conjugate formed/min/mg protein) in *Gambusia affinis*. Values are given as the mean $\pm$ SEM of 5 specimens. Significant difference compared to the control group (** $p < 0.01$, * $p < 0.05$); Significant difference compared to the carbofuran LC50/3 treated group (# $p < 0.05$) and † $p < 0.05$ vs 15 days treated group

Glutathione S-transferase activity

According to Figure 2, the results reveal an increase in hepatopancreatic GST activity in treated females compared with a control. The GST activity increased significantly from day 15 for both two concentrations tested ($p < 0.0001$; Figure 2B). The highest activity was noticed at day 40 for 0.255 mg/L concentration. The ANOVA test reveals a highly significant time effect ($F_{4,39} = 189.7$ $p < 0.0001$), treatment effect ($F_{2,32} = 113.8$; $p < 0.0001$) and time/treatment interaction ($F_{9,48} = 21.31$, $p < 0.0001$).

Catalase activity

We found an increase in catalase activity in females treated with 0.191 mg/L and 0.255 mg/L of carbofuran compared to control females. Tukey's test showed a significant difference ($p < 0.05$) for 0.191 mg/L after 40 days of treatment, while highest activity was recording at 0.255 mg/L after 15 and 40 days ($p < 0.01$, Figure. 3A).




Figure 3. Catalase (CAT) activity ($\mu\text{mol H}_2\text{O}_2/\text{min}/\text{mg}$ protein) and acetylcholinesterase (AChE) ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in *Gambusia affinis*. Values are given as the mean $\pm$ SEM of 5 specimens. Significant difference compared to the control group (** $p < 0.01$, * $p < 0.05$); Significant difference compared to the carbofuran treated group # $p < 0.05$ and † $p < 0.05$ vs 15 days treated groups

Acetylcholinesterase activity

Brain acetylcholinesterase activity was strongly inhibited after exposure of females to 0.191 and 0.255 mg/L carbofuran compared to the control. This inhibition corresponded to a reduction of about 28.84% and 52.18%, respectively after 15 days and about 40.1% and 62.2% after 40 days of carbofuran exposure (Figure 3B).

Neurochemical and behavioral response to carbofuran

AChE activity and swimming activity were determined in the fish brain at different time (2, 4, and 6 hours) during the exposure period as shown in Figure 4. The activity of AChE was reduced as the concentration and duration of carbofuran exposure increased. The inhibition of AChE activity was increased from 26% to 49.1% with the dose of 0.191 mg/L after 6 hours of exposure. At the same periods, inhibition was more significantly increased from 42% to 60.3% with the highest concentration (0.255 mg/L). This variation in AChE was affected the swimming activity of treated females, where a reduction in swimming activity was recorded after 2 hours of exposure in females exposed to 0.191 mg/L and 0.255 mg/L. A decreased in swimming activity was a time and concentration-dependent (Figure 4). The ANOVA two-way test show a significant decrease after 4 h of exposure to 0.191 mg/L, 0.255 mg/L ($p < 0.05$) and very significant after 6 h ($p < 0.001$).

Effect of carbofuran on offspring deformities

The spinal deformities observed in the larvae are shown in Figure 5. Two types of spinal deformations were found, a kyphosis (dorsal curvature) and lordosis (Lordosis ventral curvature). The percentage of deformities represented in Table 1, show that the kyphosis is highest in the fry of females treated with both two concentrations of carbofuran (20% at 0.255 mg/L and 12.5% at 0.191 mg/L). Furthermore, comparison of the frequency of malformed fry shows that the highest concentration (0.255 mg/L) causes more deformities.




Figure 4. Swimming and AchE activity of adult female *Gambusia affinis* after 2, 4 and 6 hours of exposure to carbofuran. Values are given as the mean $\pm$ S.E.M of 5 observations. Significant difference compared to the control group (* $p < 0.05$, ** $p < 0.01$); Significant difference compared to the treated group w (# $p < 0.05$, ## $p < 0.01$)




Figure 5. Red alizarin stained slides of spine representation, shown deformities in juvenile *Gambusia affinis* exposed for to carbofuran. A and B Normal spines; C, D and E bifurcated neural and hemal spines (arrow). C and F broken hemal spine (small arrow). Manification 50X

Table 1. Spinal deformity incidences of juveniles from control and exposed females to 0.191, and 0.255.66 mg/L carbofuran

	Control		$\frac{1}{4}$ LC ₅₀ 0.191 mg/l		$\frac{1}{3}$ LC ₅₀ 0.255 mg/l	
Class of deformities	Kyphosis	Lordosis	Kyphosis	Lordosis	Kyphosis	Lordosis
% of deformities	0%	2.5%	12.5% *	7.50% *	21%*#	10%*#
Total deformities	2.5%		20%*		31%*	

Number of specimens examined = 30 in each group; * p< 005: significance from Control group; #P<0.05: significance from $\frac{1}{4}$ LC₅₀ group

Discussion

In this work we tested two doses of carbofuran $\frac{1}{3}$ and $\frac{1}{4}$ LC₅₀, a carbamate pesticide that can be extremely toxic to fish (Barbieri et al., 2016; Mendes et al., 2021). Our results reveal that exposure of *G. affinis* to carbofuran-induced concentrations and time-dependent variations in GSH levels and GST activity; we found a decrease in GSH levels in the hepatopancreas by both two concentration tested ($\frac{1}{3}$ and $\frac{1}{4}$ CL₅₀), starting at 15 days of exposure, this level continues to decrease until the end of the treatment. This result is similar with that observed by Akram et al. (2021) in fresh water bighead carp. A decrease of hepatic and renal glutathione (GSH) was also detected in the African catfish *Clarias gariepinus* after its exposure to carbofuran (Hamed et al., 2017), and in liver, kidney, gills, gonads and muscles after 35 days of exposure (Ibrahim and Harabawy, 2014).

Glutathione S-transferases (GST) are belonged to the key antioxidant enzymes of the antioxidant defense system (Yang and Lee, 2015). Indeed, in phase II of biotransformation, GST enters mainly in detoxification process of xenobiotics, including herbicides and pesticides (Van der Oost et al., 2003; Higgins and Hayes, 2011) and endogenous chemicals able to conjugate with glutathione (Domingues et al., 2010). In this study, we noted an elevation in GST activity accompanied with a reduction in glutathione content. This adaptive response could be due to the solicitation of the phase II process of pesticide biotransformation in response to a substantial increase in ROS, as recently reported by Demirci and Güngördü (2020). Similar results were found in *Gambusia holbrooki* after 24 h and 96 h of exposure to acetamiprid and in *Oreochromis niloticus* exposed to 2-4- D (Oruc and Uner, 2004; Demirci, and Güngördü, 2020). Furthermore, Jiang et al. (2016) found that mRNA level of gstp1 correlates better with the activity of the GST enzyme, which hence can be controlled by the interaction of phase II products activating the regulation of antioxidant enzyme transcription and mainly that of GST.

The mechanisms of free radical detoxification are associated with a cascade of biochemical events involving the mixture of several enzymes such as CAT. CAT with superoxide dismutase SOD can be defined as the main defense line against the damaging effect caused by ROS overproduction and in particular the superoxide anion in response to chemical aggression (Hermes-Lima, 2004; Laouati et al., 2021). In addition, results showed a significant induction of CAT activity in *G. affinis* exposed to carbofuran, this increase is concentration-dependent. This response is likely explained by the catalytic activity of this enzyme to convert hydrogen peroxide H₂O₂ to water and oxygen (Amri et al., 2017). These results are in agreement with those of Nunes et al. (2004), reporting that

exposure to clofibrate and led to significant CAT activity in *Gambusia* and *Daphnia magna*. Other previous studies conducted on adult fish showed that exposure to 2,4-D and zinthosmethyl lead to increase CAT activity (Oruç and Üner, 2000; Oruc et al., 2004; Atamaniuk et al., 2013).

The variations that were recorded during this study regarding biomarker responses clearly indicate that carbofuran induce an imbalance between antioxidants and pro oxidants, this is supported in our results by a marked decrease in total lipids and an increased level of MDA in fish exposed to different concentrations of carbofuran. Lipid peroxidation (LPO) is a potential parameter of oxidative stress damage (Ibrahim and Harabawy, 2014). MDA is the end product of lipids peroxidation, and a marker of oxidative stress and cell membrane components impairment (Kuder and Gundala, 2017). In this study, the MDA concentrations was increased in time-and concentration dependent manner. Consequently, the lipid tissue of *G. affinis* can highly accumulate the pesticide. Previous studies have also reported that *Clarias batrachus* and zebrafish *Danio rerio* exposed respectively to LC₅₀ (7.66 mg/L) of carbofuran and 25 mg/L of 2,4-D induce malondialdehyde (MDA) accumulation (Begum, 2008; Li et al., 2017).

Acetylcholinesterase is a good indicator of neurotoxicity, at the neuromuscular and interneuronal junctions; the nerve impulse is transmitted through the release of acetylcholine. The inhibition of AChE by toxicants leads to a storage of chemical mediated in the synaptic spaces, maintaining a permanent transmission of the nerve impulse, which usually induces muscle tetany and death (Hernández-Moreno et al., 2011; Harabawy and Ibrahim, 2014). AChE activity is used in toxicology as organophosphates and carbamates toxicity marker (Fossi et al., 2001; Narra, 2016). In this study, we observed an inhibition of AChE activity of about 30% started from the 15th day of treatment in *G. affinis* exposed to carbofuran, this inhibition is progressive until the end of treatment. Carbamate insecticides are specific inhibitors of AChE in several freshwater fish (Pessoa et al., 2011; Nunes et al., 2014; Ensibi et al., 2014). Similar results to ours have shown that treatment with organophosphate insecticides cause a decrease of AChE activity in the hemolymph after 24 h of exposure in the blue mussel *Mytilus edulis* (Rickwood and Galloway, 2004). In the same context Narra (2016), reported a decrease in AChE activity in the brain of *Clarias batrachus* after Sub-acute exposure 1/10th of LC₅₀ for 60 days to carbofuran.

Our findings also revealed that carbofuran appears to work in a time-and concentration dependent manner, as the inhibition of AChE rises with increasing carbofuran concentration and exposure time (30% at 0.191 mg/L on day 15 and 50% at 0.255 mg/L on day 40). Bretau et al. (2001), revealed that the inhibition of AChE by carbofuran is concentration dependent, but several studies have reported that the inhibition of AChE activity is time dependent (Amblard et al., 1998; Legierse et al., 1999; Liu et al., 2013).

In this study, we tried to investigate the correlation between carbofuran AChE inhibition and locomotors behavior. Acute exposure to carbofuran causes a significant decrease in swimming activity after 4 h for both doses in *G. affinis*, this decrease is very significant after 6 h of exposure to the highest concentration, this disorder of swimming activity is accompanied by an inhibition of AChE activity, the activity of AChE was increased from 26% to 49.1% at 0.191 mg/L respectively after 2 h, 4 h and 6 h of exposure. At the same periods the inhibition rates are more important in the poisons treated with the highest dose 0.255 mg/L. These observations suggest the existence of a correlation between the concentration of carbofuran, AChE activity and swimming behavior. Similar results have been reported in *Carassius auratus* treated with various

concentrations of carbofuran (Bretaud et al., 2001) and in *Gambusia yucatana* exposed to chlorpyrifos, carbofuran and glyphosate (Rendón-von Osten et al., 2005). We found through the results obtained that the activity of the brain AChE in *G. affinis* is more sensitive to carbofuran compared to swimming activity. Indeed, the impact of carbofuran on swimming activity is only observed when AChE activity is inhibited at 44.2%. In this sense, a few works have reported that at concentrations near to lethal toxicity values, disturbances of locomotor behavior and swimming speed were observed in fish in response to exposure to carbofuran (Hernandez-Moreno et al., 2011; Pessoa et al., 2011; Campos-Garcia et al., 2015; Mendes et al., 2021).

Can the injurious effects observed after exposure *G. affinis* to carbofuran affect their offspring? Undeniably, the results obtained on the morphology of the juveniles of fish exposed to different concentrations of carbofuran clearly confirm this hypothesis. The effects of carbofuran can be transmitted to the offspring. Our results show an induction of spinal skeletal deformations (Kyphosis and Lordosis) in the larvae of *G. affinis* treated females compared to control and that the rate of these skeletal deformations is concentration-dependent.

In fish, the young stages are particularly sensitive to environmental contamination and can lead to the appearance of malformations, in particular deformations of the vertebral column (Messaoudi et al., 2009). Several studies have shown that skeletal deformation is strongly correlated with external stresses (Messaoudi et al., 2008). Thus, exposure to certain toxic, potentially teratogenic substances such as organophosphate herbicides and pesticides, carbamates, dioxin, polyaromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs), and heavy metals such as zinc and cadmium can alter the processes of bone formation and development and consequently lead to skeletal deformities (Koyama, 1996; Olsson et al., 1999; Cheng et al., 2000; Teraoka et al., 2002; Teh et al., 2002). These skeletal deformations are relatively well described in several fish species; however, the mechanisms of their induction are poorly studied. Some Studies have shown that these deformations are caused by an imbalance in phosphocalcic metabolism leading to decalcification of bones and resulting in their fragility (Muramoto, 1981; Sato et al., 1983; Akiyama et al., 1986; Dabrowski et al., 1990; Kessabi et al., 2009).

Our results indicated an increase in the rate of skeletal deformations in larvae of the females treated with different concentrations of carbofuran. This suggests that carbofuran may have a potential negative effect on the molecular processes and mechanisms regulating the development and growth of bone tissue in fish. As a result, carbofuran could have teratogenic effects on *Gambusia affinis* larvae. Our results are confirmed by the work of Pawar and Katdre (1984) that showed a toxic and a teratogenic effect in *Microhyla ornat tadpoles* exposed to carbofuran, namely, body axis body curvatures, growth retardation and behavioral disturbances. In the same way Sassi et al. (2010) showed that cadmium treatment at high temperature (32°C) affects the embryos of *G. affinis* and causes an increase in frequency of Kyphosis and Lordosis.

In this study, the multi-marker approach at various levels of biological organization enabling us to understand the potential of *G. affinis* as a bioindicator species of pollution and gives a reliable prediction of the adverse effects of pesticides and emerging contaminants in aquatic organisms. In summary, it can be concluded that low dose exposure to carbofuran can induce oxidative stress in *G. affinis*. We observe increased activity of enzymatic antioxidants in response to carbofuran, justifying the adaptive capacities of this non-target species to environmental stresses. In this study, fish swimming activity and AChE activity are correlated and support a neurotoxic effect of

carbofuran however, AChE is more sensitive to the pesticide even at a low concentration. These effects may alter the health of aquatic organisms and suggest that carbofuran may be a potentially dangerous aggressor on non-target organisms.

Conclusions

The exposure of females of *Gambusia affinis* to carbofuran can affect their offspring, resulting in an increased incidence of spinal deformities. Therefore, the effects of carbamate become important in assessing the teratogenic effects on non-target organisms. Additionally, these results provide sufficient knowledge for further research studies on the impact of environmental factors responsible for bone deformations and in particular parameters related to phosphocalcic metabolism in fish.

REFERENCES

- [1] Abdulelah, S. A., Crile, K. G., Almouseli, A., Awali, S., Tutwiler, A. Y., Tien, E. A., Manzo, V. J., Hadeed, M. N., Belanger, R. M. (2020): Environmentally relevant atrazine exposures cause DNA damage in cells of the lateral antennules of crayfish (*Faxonius virilis*). – *Chemosphere* 239: 124786.
- [2] Aebi, H. (1984): Catalase in vitro. – *Methods in Enzymology* 105: 121-126.
- [3] Akiyama, T., Murai, T., Mori, K. (1986): Role of tryptophan metabolites in inhibition of spinal deformity of chum salmon fry caused by tryptophan deficiency. – *Nippon Suisan Gakkaishi* 52(7): 1255-1259.
- [4] Akram, R., Iqbal, R., Hussain, R., Jabeen, F., Ali, M. (2021): Evaluation of oxidative stress, antioxidant enzymes and genotoxic potential of bisphenol A in fresh water bighead carp (*Aristichthys nobilis*) fish at low concentrations. – *Environmental pollution* 268: 115896.
- [5] Amblard, G., Bry, C., Toutant, J. P., Arpagaus, M. (1998): Effets de l'exposition à un insecticide carbamate, le carbofuran, sur les performances de jeunes stades de brochet (*Esox lucius* L., 1758): résultats préliminaires. – *Bulletin Français de la Pêche et de la Pisciculture* 350-351: 529-534.
- [6] Amri, S., Samar, M. F., Sellem, F., Ouali, K. (2017): Seasonal antioxidant responses in the sea urchin *Paracentrotus lividus* (Lamarck 1816) used as a bioindicator of the environmental contamination in the South-East Mediterranean. – *Marine pollution bulletin* 122(1-2): 392-402.
- [7] APHA-AWWA-WEF (1998): Standard methods for the examination of water and waste water. – Clesceri, L. S., Greenberg, A. E., Eaton, A. D. (eds.) 20th ed. American Public Health Association, American Water Works Association, Water Environment Federation, Washington, DC.
- [8] Archer, E., Wolfaardt, G. M., van Wyk, J. H., van Blerk, N. (2020): Investigating (anti) estrogenic activities within South African wastewater and receiving surface waters: Implication for reliable monitoring. – *Environmental Pollution* 263: 114424.
- [9] Atamaniuk, T. M., Kubrak, O. I., Storey, K. B., Lushchak, V. I. (2013): Oxidative stress as a mechanism for toxicity of 2, 4-dichlorophenoxyacetic acid (2, 4-D): studies with goldfish gills. – *Ecotoxicology* 22(10): 1498-1508.
- [10] Barbieri, E., Moreira, P., Luchini, L. A., Ruiz Hidalgo, K., Munoz, A. (2016): Assessment of acute toxicity of carbofuran in *Macrobrachium olfersii* (Wiegmann, 1836) at different temperature levels. – *Toxicology and Industrial Health* 32(1): 7-14.
- [11] Begum, G. (2008): Assessment of biochemical markers of carbofuran toxicity and recovery response in tissues of the freshwater teleost, *Clarias batrachus* (Linn). – *Bulletin of environmental contamination and toxicology* 81(5): 480-484.

- [12] Bouzahouane, H., Barour, C., Sleimi, N., Ouali, K. (2018): Multi-biomarkers approach to the assessment of the southeastern Mediterranean Sea health status: Preliminary study on *Stramonita haemastoma* used as a bioindicator for metal contamination. – *Chemosphere* 207: 725-741.
- [13] Bradford, M. M. (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizes the principle of protein-dye binding. – *Anal. Biochem* 72(1-2): 248-254.
- [14] Bretaud, S., Saglio, P., Toutant, J. P. (2001): Effets du carbofuran sur l'activité de l'acétylcholinestérase cérébrale et sur l'activité de nage chez *Carassius auratus* (Cyprinidae). – *Cybium* 25(1): 33-40.
- [15] Campos-Garcia, J., Martinez, D. S. T., Alves, O. L., Leonardo, A. F. G., Barbieri, E. (2015): Ecotoxicological effects of carbofuran and oxidised multiwalled carbon nanotubes on the freshwater fish *Nile tilapia*: Nanotubes enhance pesticide ecotoxicity. – *Ecotoxicology and Environmental Safety* 111: 131-137.
- [16] Cheghib, Y., Chouahda, S., Soltani, N. (2020): Side-effects of a neonicotinoid insecticide (actara®) on a non-target larvivorous fish *Gambusia affinis*: growth and biomarker responses. – *The Egyptian Journal of Aquatic Research* 46(2): 167-172.
- [17] Cheng, S. H., Wai, A. W. K., So, C. H., Wu, R. S. S. (2000): Cellular and molecular basis of cadmium-induced deformities in zebrafish embryos. – *Environmental Toxicology and Chemistry* 19(12): 3024-3031.
- [18] Dabrowski, K., El-Fiky, N., Köck, G., Frigg, M., Wieser, W. (1990): Requirement and utilization of ascorbic acid and ascorbic sulfate in juvenile rainbow trout. – *Aquaculture* 91(3-4): 317-337.
- [19] Darvishi, M., Safari, R., Hoseinifar, S. H., Shabani, A., Dadar, M., Jarayedi, Z., Paolucci, M. (2022): Sublethal doses of diazinon affected reproductive, immune, and oxidative status in female zebrafish (*Danio rerio*). – *Aquaculture Reports* 22: 100944.
- [20] Demirci, Ö., Güngördü, A. (2020): Evaluation of the biochemical effects of an acetamiprid-based insecticide on a non-target species, *Gambusia holbrooki*. – *Water and Environment Journal* 34: 481-489.
- [21] Domingues, I., Oliveira, R., Lourenço, J., Grisolia, C. K., Mendo, S., Soares, A. M. V. M. (2010): Biomarkers as a tool to assess effects of chromium (VI): comparison of responses in zebrafish early life stages and adults. – *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 152(3): 338-345.
- [22] Draper, H. H., Hadley, M. (1990): Malondialdehyde determination as index of lipid Peroxidation. – *Methods in Enzymology* 186: 421-431.
- [23] Ellman, G. L., Courtney, K. D., Andres Jr, V., Featherstone, R. M. (1961): A new and rapid colorimetric determination of acetylcholinesterase activity. – *Biochemical pharmacology* 7(2): 88-95.
- [24] Ensibi, C., Hernández-Moreno, D., Míguez Santiyán, M. P., Daly Yahya, M. N., Rodríguez, F. S., Pérez-López, M. (2014): Effects of carbofuran and deltamethrin on acetylcholinesterase activity in brain and muscle of the common carp. – *Environmental Toxicology* 29(4): 386-393.
- [25] Fossi, M. C., Minutoli, R., Guglielmo, L. (2001): Preliminary results of biomarker responses in zooplankton of brackish environments. – *Marine pollution bulletin* 42(9): 745-748.
- [26] Garcês, A., Pires, I., Rodrigues, P. (2020): Teratological effects of pesticides in vertebrates: A review. – *Journal of Environmental Science and Health, Part B* 55(1): 75-89.
- [27] Goldsworthy, G. J., Mordue, W., Guthkelch, J. (1972): Studies on insect adipokinetic hormones. – *General and Comparative Endocrinology* 18(3): 545-551.
- [28] Habig, W. H., Pabst, M. J., Jakoby, W. B. (1974): Glutathione S-transferases: the first enzymatic step in mercapturic acid formation. – *Journal of biological Chemistry* 249(22): 7130-7139.

- [29] Hamed, H. S., Osman, A. G. (2017): Modulatory effect of lycopene against carbofuran toxicity in African catfish, *Clarias gariepinus*. – Fish physiology and biochemistry 43(6): 1721-1731.
- [30] Hamed, H. S., Ismal, S. M., Faggio, C. (2021): Effect of allicin on antioxidant defense system, and immune response after carbofuran exposure in *Nile tilapia*, *Oreochromis niloticus*. – Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology 240: 108919.
- [31] Harabawy, A. S., Ibrahim, A. T. A. (2014): Sublethal toxicity of carbofuran on the African catfish *Clarias gariepinus*: Hormonal, enzymatic and antioxidant responses. – Ecotoxicology and Environmental Safety 106: 33-39.
- [32] Hermes-Lima, M. (2004): Oxygen in biology and biochemistry: role of free radicals. Functional metabolism: regulation and adaptation. – John Wiley & Sons, Inc., Hoboken: 319-368.
- [33] Hernandez-Moreno, D., Perez-Lopez, M., Soler, F., Gravato, C., Guilhermino, L. (2011): Effects of carbofuran on the sea bass (*Dicentrarchus labrax* L.): Study of biomarkers and behaviour alterations. – Ecotoxicology and Environmental Safety 74(7): 1905-1912.
- [34] Higgins, L. G., Hayes, J. D. (2011): Mechanisms of induction of cytosolic and microsomal glutathione transferase (GST) genes by xenobiotics and pro-inflammatory agents. – Drug Metabolism Reviews 43(2): 92-137.
- [35] Ibrahim, A. T. A., Harabawy, A. S. (2014): Sublethal toxicity of carbofuran on the African catfish *Clarias gariepinus*: Hormonal, enzymatic and antioxidant responses. – Ecotoxicology and Environmental Safety 106: 33-39.
- [36] Jiang, J., Chen, Y., Yu, R., Zhao, X., Wang, Q., Cai, L. (2016): Pretilachlor has the potential to induce endocrine disruption, oxidative stress, apoptosis and immunotoxicity during zebrafish embryo development. – Environmental Toxicology and Pharmacology 42: 125-134.
- [37] Kasonga, T. K., Coetzee, M. A., Kamika, I., Ngole-Jeme, V. M., Momba, M. N. B. (2021): Endocrine-disruptive chemicals as contaminants of emerging concern in wastewater and surface water: A review. – Journal of Environmental Management 277: 111485.
- [38] Kessabi, K., Kerkeni, A., Saïd, K., Messaoudi, I. (2009): Involvement of Cd bioaccumulation in spinal deformities occurrence in natural populations of Mediterranean killifish. – Biological trace element research 128(1): 72-81.
- [39] Koyama, J. (1996): Vertebral deformity susceptibilities of marine fishes exposed to herbicide. – Bulletin of Environmental Contamination and Toxicology 56(4): 655-662.
- [40] Kuder, R. S., Philip, G. H. (2017): Antioxidant enzymatic activities and lipid peroxidation in liver and ovary of zebrafish (*Danio rerio*) exposed to deltamethrin. – Chemistry and Ecology 33(8): 739-749.
- [41] Laouati, I., Rouane-Hacene, O., Derbal, F., Ouali, K. (2021): The mussel caging approach in the assessment of trace metal contamination in southern Mediterranean coastal waters: a multi-biomarker study. – Environmental Science and Pollution Research 28(44): 63032-63044.
- [42] Legierse, K. C., Verhaar, H. J., Vaes, W. H., De Bruijn, J. H., Hermens, J. L. (1999): Analysis of the time-dependent acute aquatic toxicity of organophosphorus pesticides: The critical target occupation model. – Environmental Science & Technology 33(6): 917-925.
- [43] Li, K., Wu, J. Q., Jiang, L. L., Shen, L. Z., Li, J. Y., He, Z. H., Wei, P., Lv, Z., He, M. F. (2017): Developmental toxicity of 2, 4-dichlorophenoxyacetic acid in zebrafish embryos. – Chemosphere 171: 40-48.
- [44] Liu, X. J., Luo, Z., Zheng, J. L., Xiong, B. X. (2013): Effects of waterborne acephate exposure on antioxidant responses and acetylcholinesterase activities in *Synechogobius hasta*. – Environmental Toxicology 28(1): 42-50.
- [45] Liu, S., Yu, M., Xie, X., Ru, Y., Ru, S. (2020): Carbofuran induces increased anxiety-like behaviors in female zebrafish (*Danio rerio*) through disturbing dopaminergic/norepinephrinergic system. – Chemosphere 253: 126635.

- [46] Mahboob, S., Al-Ghanim, K. A., Fares Al-Balwi, H. A., Al-Misned, F., Ahmad, Z. (2014): Carbofuran-induced effects on acetylcholinesterase (AChE) in erythrocytes and liver of *Cyprinus carpio*. – *Toxicological & Environmental Chemistry* 96(4): 614-623.
- [47] Mansano, A. S., Moreira, R. A., Pierozzi, M., Oliveira, T. M., Vieira, E. M., Rocha, O., Regali-Seleglim, M. H. (2016): Effects of diuron and carbofuran pesticides in their pure and commercial forms on *Paramecium caudatum*: The use of protozoan in ecotoxicology. – *Environmental Pollution* 213: 160-172.
- [48] Mendes, R. C., Henriques, M. B., Barbieri, E. (2021): Carbofuran affects behavior and metabolism of the Atlantic Forest *lambari Deuterodon iguape*, a native species from Brazil. – *Environmental Science and Pollution Research* 28(43): 61128-61136.
- [49] Messaoudi, I., Deli, T., Kessabi, K., Barhoumi, S., Kerkeni, A., Saïd, K. (2008): Association of spinal deformities with heavy metal bioaccumulation in natural populations of grass goby, *Zosterisessor ophiocephalus* Pallas, 1811 from the Gulf of Gabès (Tunisia). – *Environmental Monitoring and Assessment* 156(1): 551-560.
- [50] Messaoudi, I., Kessabi, K., Kacem, A., Saïd, K. (2009): Incidence of spinal deformities in natural populations of *Aphanius fasciatus* Nardo, 1827 from the Gulf of Gabes, Tunisia. – *Afr J Ecol* 47(3): 360-366.
- [51] Mishra, S., Zhang, W., Lin, Z., Pang, S., Huang, Y., Bhatt, P., Chen, S. (2020): Carbofuran toxicity and its microbial degradation in contaminated environments. – *Chemosphere* 259: 127419.
- [52] Muramoto, S. (1981): Vertebral column damage and decrease of calcium concentration in fish exposed experimentally to cadmium. – *Environmental Pollution Series A, Ecological and Biological* 24(2): 125-133.
- [53] Narra, M. R. (2016): Single and cartel effect of pesticides on biochemical and haematological status of *Clarias batrachus*: A long-term monitoring. – *Chemosphere* 144: 966-974.
- [54] Nunes, B., Carvalho, F., Guilhermino, L. (2004): Acute and chronic effects of clofibrate and clofibric acid on the enzyme's acetylcholinesterase, lactate dehydrogenase and catalase of the mosquitofish, *Gambusia holbrooki*. – *Chemosphere* 57(11): 1581-1589.
- [55] Olsson, P. E., Westerlund, L., Teh, S. J., Billsson, K., Berg, A. H., Tysklind, M., Hinton, D. E. (1999): Effects of maternal exposure to estrogen and PCB on different life stages of zebrafish (*Danio rerio*). – *Ambio* 28(1): 100-106.
- [56] Oruç, E. Ö., Üner, N. (2000): Combined effects of 2, 4-D and azinphosmethyl on antioxidant enzymes and lipid peroxidation in liver of *Oreochromis niloticus*. – *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology* 127(3): 291-296.
- [57] Oruc, E. O., Sevgiler, Y., Uner, N. (2004): Tissue-specific oxidative stress responses in fish exposed to 2, 4-D and azinphosmethyl. – *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 137(1): 43-51.
- [58] Pawar, K. R., Katdare, M. (1984): Toxic and teratogenic effects of fenitrothion, BHC and carbofuran on embryonic development of the frog *Microhyla ornata*. – *Toxicology letters* 22(1): 7-13.
- [59] Pessoa, P. C., Luchmann, K. H., Ribeiro, A. B., Veras, M. M., Correa, J. R. M. B., Nogueira, A. J., Bainy, A. C. D., Carvalho, P. S. M. (2011): Cholinesterase inhibition and behavioral toxicity of carbofuran on *Oreochromis niloticus* early life stages. – *Aquatic Toxicology* 105(3-4): 312-320.
- [60] Rendón-von Osten, J., Ortiz-Arana, A., Guilhermino, L., Soares, A. M. V. M. (2005): In vivo evaluation of three biomarkers in the mosquitofish (*Gambusia yucatana*) exposed to pesticides. – *Chemosphere* 58(5): 627-636.
- [61] Rickwood, C. J., Galloway, T. S. (2004): Can acetylcholinesterase be considered a biomarker of effect? A study of the mussel *Mytilus edulis* exposed to the priority pollutant Chlorfenvinphos. – *Aquatic Toxicology* 67: 45-56.

- [62] Roy, P. K., Lall, S. P. (2007): Vitamin K deficiency inhibits mineralization and enhances deformity in vertebrae of haddock (*Melanogrammus aeglefinus* L.). – Comparative Biochem Physiol. Part B 148: 174-183.
- [63] Sassi, A., Annabi, A., Kessabi, K., Kerkeni, A., Saïd, K., Messaoudi, I. (2010): Influence of high temperature on cadmium-induced skeletal deformities in juvenile mosquitofish (*Gambusia affinis*). – Fish physiology and biochemistry 36(3): 403-409.
- [64] Sato, M., Kondo, T., Yoshinaka, R., Ikeda, S. (1983): Effect of water temperature on the skeletal deformity in ascorbic acid-deficient rainbow trout [*Salmo gairdnerii*]. – Bulletin of the Japanese Society of Scientific Fisheries 49(3): 443-446.
- [65] Sfakianakis, D. G., Renieri, E., Kentouri, M., Tsatsakis, A. M. (2015): Effect of heavy metals on fish larvae deformities: a review. – Environmental Research 137: 246-255.
- [66] Sousa, B., Nunes, B. (2020): Reliability of behavioral test with fish: How neurotransmitters may exert neuromodulatory effects and alter the biological responses to neuroactive agents. – Science of The Total Environment 734: 139372.
- [67] Staples, R. (1964): Refinements in rapid clearing technic in the KOH-alizarin red S method for fetal bone. – Stain Technology 39: 61-63.
- [68] Teh, S. J., Deng, X., Teh, F. C., Hung, S. O. (2002): Selenium-induced teratogenicity in Sacramento splittail (*Pogonichthys macrolepidotus*). – Marine Environmental Research 54(3-5): 605-608.
- [69] Teraoka, H., Dong, W., Ogawa, S., Tsukiyama, S., Okuhara, Y., Niiyama, M., Ueno, N., Peterson, R., Hiraga, T. (2002): 2, 3, 7, 8-Tetrachlorodibenzo-p-dioxin toxicity in the zebrafish embryo: altered regional blood flow and impaired lower jaw development. – Toxicological Sciences 65(2): 192-199.
- [70] Tissot, A. G., Granek, E. F., Thompson, A. W., Hladik, M. L., Moran, P. W., Scully-Engelmeyer, K. (2022): The silence of the clams: Forestry registered pesticides as multiple stressors on soft-shell clams. – Science of the Total Environment 819: 152053.
- [71] Van der Oost, R., Beyer, J., Vermeulen, N. P. (2003): Fish bioaccumulation and biomarkers in environmental risk assessment: a review. – Environmental Toxicology and Pharmacology 13(2): 57-149.
- [72] Weckbecker, G., Cory, J. G. (1988): Ribonucleotide reductase activity and growth of glutathione-depleted mouse leukemia L1210 cells in vitro. – Cancer letters 40(3): 257-264.
- [73] Yang, H. Y., Lee, T. H. (2015): Antioxidant enzymes as redox-based biomarkers: a brief review. – BMB Reports 48(4): 200.