

First check list of Serbian Trichoptera

IVANA ŽIVIĆ, ZORAN MARKOVIĆ & MILOJE BRAJKOVIĆ

ABSTRACT: One hundred and seventy-three species of Trichoptera have been reported to date in the fauna of Serbia. Of the recorded species, *Limnephilus petri* Marinkovic is endemic for the fauna of Serbia; *Chaetopteryx maxima* Kumanski and *Drusus botosaneanui* Kumanski are endemics of the Eastern Balkans zoogeographic region; and *Rhyacophila bosniaca* Schmid and *Rhyacophila vranitzensis* Marinkovic et Botosaneanu are endemics of the Dinaric Western Balkans.

Of the 173 species, 65.3% (113 species) also inhabit at least one other republic of the former Yugoslavia in addition to Serbia. These are in fact species with a limited range of distribution (in the central and western parts of the Balkan Peninsula).

Introduction

The Trichoptera represent a homogeneous group of insects with two suborders: Annulipalpia and Integipalpia. This division was supplemented by Ross (1967), who relegated all recent Trichoptera to three superfamilies: Rhyacophiloidea (with the families Rhyacophilidae, Glossosomatidae, and Hydroptilidae); Hydropsychoidea (includes all Annulipalpia except the indicated families of Rhyacophiloidea); and Limnephiloidea (all Integipalpia). About 10,000 species have been described to date, although it is estimated that around 50,000 exist HOLZENTHAL & BLAHNIK (2000). Eight hundred and ninety-five species are known in the fauna of Europe Illies (1978).

The first faunistic data on Trichoptera of the former Yugoslavia were given by RADOVANOVIC (1931), who registered 59 species. Four years later, RADOVANOVIC (1935) published a list of 170 species of Trichoptera. Nine species new for the fauna of Yugoslavia were reported and two species new to science (*Drusus discophorus* Radovanovic and *Drusus plicatus* Radovanovic) were described in 1943, RADOVANOVIC (1943). RADOVANOVIC (1953) subsequently found 45 additional species of Trichoptera, of which six (*Cyrnus cernatiocornis* Kolenati, *Leptocerus annulicornis* Stephens, *Colpotaulius incisus* Curtis, *Grammotaulius nitidus* Müller, *Limnephilus stigma* Curtis and *Oecismus monedula* Hagen) were new for the fauna of Yugoslavia.

A great contribution to knowledge about Trichoptera of the former Yugoslavia was made by MARINKOVIC-GOSPODNETIC. The indicated author in 1966 described five species new to science and for the fauna of Yugoslavia: *Chaetopteryx gonospina* Mar., *Chaetopteryx stankovici* Mar., *Hydropsyche botosaneanui* Mar., *Plectrocnemia smiljajae* Mar., *Limnophilus petri* Mar. She subsequently described six species new to science and for the fauna of Yugoslavia in 1970 (*Drusus klapaleki* Mar., *Drusus radovanovici* Mar., *Drusus ramae* Mar., *Potamophylax schmidi* Mar., *Psylopteryx bosniaca* Mar. and *Crunoecia bosniaca* Mar.) and two more in 1971 (*Drusus serbicus* Mar. and *Drusus croaticus* Mar.). Investigating the

Trichoptera fauna of the Morača River and its tributaries, MARINKOVIC-GOSPODNETIC (1981) found seven species new for the territory of Yugoslavia: *Glossosoma bifidum* Hagen, *Hydroptila vectis* Curtis, *Limnephilus graecus* Schmid, *Athripsodes dissimilis* Linne, *Triaenodes ochrellus* Ramb., *Oecetis ochracea* Curtis and *Leptocerus interruptus* Fabricus.

The number of authors who have conducted research on the Trichoptera faunas of republics of the former Yugoslavia is small.

The Trichoptera fauna of Slovenia has been little investigated: only 54 species have been recorded RADOVANOVIC (1933, 1953).

About 120 species have been recorded for the fauna of Croatia MATONICKIN (1959, 1987); PAVLETIC and MATONICKIN (1972); HABDIJA (1979); MATONICKIN et al. (1975).

Of republics of the former Yugoslavia, Bosnia-Herzegovina is the one with the most thoroughly studied fauna of Trichoptera. Thanks to the investigations of RADOVANOVIC (1953) and MARINKOVIC-GOSPODNETIC (1970a, 1978, 1979), about 300 species of Trichoptera have been recorded there.

Around 140 species of Trichoptera have been recorded in the fauna of Macedonia. The first comprehensive investigations of the Trichoptera fauna of Macedonia were carried out by RADOVANOVIC (1943, 1953), who found 46 species. The Trichoptera fauna of Macedonia was also investigated by Botosaneanu (1960), who recorded 41 species for the territory of Western Macedonia, 26 of them being new for the fauna of Macedonia. PAVLOVSKI (1991) registered 15 species of Trichoptera in the Babuna River, of which *Rhyacophila loxias* Schmid and *Rhyacophila armeniaca* Guer were new for the fauna of Macedonia. ARSOV (1991) found 41 species of Trichoptera in the Zrnovska River, eight of them new for the fauna of Macedonia.

The Trichoptera fauna of Montenegro has been only partially studied, but can be said to be rich. The greatest number of described species come from the region of Mt. Durmitor, on which about 150 species have been confirmed. The first faunistic data on the Trichoptera fauna of Montenegro (the region of Mt. Durmitor) were given by RADOVANOVIC (1935, 1953), who recorded 19 species. MALICKY (1982) described a species new to science, *Plectrocnemia mojkovacensis*, from the Bistrica River and the species *Rhyacophila diakoftensis* from the Tara River MALICKY (1983). KRUSNIK (1987) found 95 species of Trichoptera on Mt. Durmitor, 67 of them new for the fauna of Montenegro. Studying the Trichoptera of the Moraca River and its tributaries (the Zeta, Slatina, Manastir, Maljestak, and Plavnica Rivers), MARINKOVIC-GOSPODNETIC (1981) recorded 48 species of Trichoptera.

In the Trichoptera fauna of Serbia, 151 species were registered up to 1980. Data on the fauna were published in papers of RADOVANOVIC (1931, 1935, 1953) and MARINKOVIC-GOSPODNETIC (1975, 1980). In the past 20 years, information has been gathered on the larvae of this very interesting group of insects, which have been studied within the framework of research on the macrozoobenthos of aquatic ecosystems SIMIC (1993); SIMIC & OSTOJIC (1994); MARKOVIC (1995); MARKOVIC & MILJANOVIC (1995); KONTA (1997); MARKOVIC et al. (1997, 1997a, 1997b); MARKOVIC (1998); MARKOVIC et al. (1998); MARTINOVIC-VITANOVIC et al. (1998); MARKOVIC et al. (1999); MILJANOVIC (2000); STRAHINIC (2000); PAUNOVIC (2001); ŽIVIĆ et al. (2001, 2001a).

The inadequate investigation of Trichoptera in Serbia presents a challenge to researchers, and it is to be hoped that considerably more attention will be devoted to this very interesting order of insects in the period to come.

Results

One hundred and seventy-three species of Trichoptera have been reported to date in the fauna of Serbia. Subsequent to the investigations of RADOVANOVIC (1931, 1935, 1953) and MARINKOVIC-GOSPODNETIC (1975, 1980), 22 more species of Trichoptera new for the fauna of Serbia were recorded (at the larval stage) BARACKOV (1973); KONTA (1997); SIMIC (1993); MARKOVIC et al. (1997a); MARKOVIC (1998); MARKOVIC et al. (1998, 1999); STRAHINIC (2000); ŽIVIĆ et al. (2001); ŽIVIĆ et al. (2001a). Among the registered species, *Limnephilus petri* Marinkovic is endemic for the fauna of Serbia; the species *Chaetopteryx maxima* Kumanski and *Drusus botosaneanui* Kumanski are endemics of the Eastern Balkans zoogeographic region; and *Rhyacophila bosniaca* Schmid and *Rhyacophila vranitzensis* Marinkovic et Botosaneanu are endemics of the Dinaric Western Balkans ILLIES (1978).

The greatest number of recorded species of Trichoptera belong to the families Limnephilidae (32.94%) and Rhyacophilidae (13.29%). The family Hydroptilidae is represented by three species; the families Helicopsychidae and Odontoceridae are each represented by two species; and the families Ecnomidae, Phryganeidae, and Uenoidae are each represented by a single species. It has been established that 113 species also inhabit at least one other republic of the former Yugoslavia in addition to Serbia. These are in fact species with a limited range of distribution (in the western and central parts of the Balkan Peninsula).

Investigations of Trichoptera have to date been carried out on a small area of Serbia. Moreover, greater diversity of this order of insects has been recorded in neighboring countries (where more attention has been paid to the Trichoptera): about 230 species have been registered in Bulgaria KUMANSKI (1985, 1988); 240 species in Romania ARSOV (1991); 211 species in Hungary SCHMERA (2001); and 300 species in Bosnia-Herzegovina RADOVANOVIC (1953); MARINKOVIC-GOSPODNETIC (1970a, 1978, 1979). In view of these facts, it can be assumed that the diversity of Trichoptera in Serbia is significantly greater than the reported 173 species.

FAUNISTIC DATA ON TRICHOPTERA SPECIES REPORTED IN SERBIA

The list of recorded species of Trichoptera is given in accordance with the systematics of ILLIES (1978) from „Limnofauna Europea“. Given in parentheses after each species name are numbers indicating who reported it (the numbers are ordinal numbers of authors in the list of references).

RHYACOPHILIDAE

- Rhyacophila aquitanica* Mac Lachlan, 1879 (17)
- Rhyacophila balcanica* Radovanovic, 1953 (17, 40)
- Rhyacophila bosniaca* Schmid, 1970 (20)
- Rhyacophila furcifera* Klapalek, 1904 (17)
- Rhyacophila fischeri* Botosaneanu, 1957 (20)
- Rhyacophila hirticornis* Mac Lachlan (20, 40)
- Rhyacophila laevis* Pictet, 1834 (17, 40)
- Rhyacophila loxias* Schmid, 1970 (17)
- Rhyacophila nubila* Zetterstet, 1840 (2, 7, 17, 27, 35, 38, 40, 46, 48, 50)
- Rhyacophila obliterata* Mac Lachlan, 1863 (17)
- Rhyacophila obtusa* Klapalek, 1894 (**R. obtusidens**) (17, 40)
- Rhyacophila polonica* Mac Lachlan, 1879 (7, 17)
- Rhyacophila trescavicensis* Botosaneanu, 1960 (20)
- Rhyacophila tristis* Pictet, 1834 (2, 17, 42)
- Rhyacophila vranitzensis* Marinkovic et Botosaneanu, 1967 (17)

Rhyacophila fasciata Hagen, 1859 (*Rh. septentrionis*) (24, 26, 28, 29, 38, 40)
Rhyacophila evoluta Mac Lachlan, 1879 (38, 40)
Rhyacophila vulgaris Pictet (38, 40)
Rhyacophila dorsalis Curtis, 1834 (25, 35)
Rhyacophila philopotamoides Mac Lachlan, 1879 (2)
Rhyacophila pascoei Mac Lachlan, 1879 (48, 49, 50)
Rhyacophila praemorsa Mac Lachlan, 1879 (7)

GLOSSOSOMATIDAE

Glossosoma bifidum Hagen (17)
Glossosoma boltoni Curtis, 1834 (7, 17, 35, 38, 46)
Glossosoma discophorum Klapalek, 1902 (17)
Syngapetus iridipennis Mac Lachlan, 1902 (17)
Syngapetus krawanyi Ulmer, 1938 (20)
Agapetus laniger Pictet, 1834 (17, 40)
Agapetus ochripes Curtis, 1834 (**A. comatus**) (17, 40)
Agapetus slavorum Botosaneanu, 1960 (17)
Agapetus fuscipes Curtis, 1834 (27, 46)

HYDROPTILIDAE

Hydroptila vectis Curtis, 1834 (2)
Hydroptila sparsa Curtis, 1834 (46)
Agraylea multipunctata Curtis, 1834 (28)

PHILOPOTAMIDAE

Philopotamus montanus Donovan, 1813 (7, 17, 35, 38, 40, 42, 46)
Philopotamus variegatus Scopoli, 1763 (17, 42)
Wormaldia occipitalis Pictet, 1834 (17)
Wormaldia pulla Mac Lachlan, 1878 (17)
Wormaldia subnigra Mac Lachlan, 1865 (2, 17, 46)
Chimarra marginata Linne, 1767 (unpublished data)

HYDROPSYCHIDAE

Dipletrona atra Mac Lachlan (*Hydropsyche helenica*) (17)
Dipletrona felix Mac Lachlan (35, 37)
Hydropsyche angustipennis Curtis, 1834 (7, 17, 22, 25, 27, 29, 30, 35, 38, 40, 48, 49, 50, 51, 52)
Hydropsyche botosaneanui Marinkovic-Gospodnetic, 1966 (20)
Hydropsyche bulbifera Mac Lachlan, 1878 (7)
Hydropsyche contubernalis Mac Lachlan, 1865 (29, 48, 49, 50)
Hydropsyche modesta Novás, 1925 (*Hydropsyche dissimulata* Kumanski et Botosaneanu) (20, 50, 51)
Hydropsyche fulvipes Curtis, 1834 (17, 42, 51)
Hydropsyche instabilis Curtis, 1834 (7, 17, 35, 40, 50)
Hydropsyche ornata Mac Lachlan, 1878 (17, 42)
Hydropsyche pellucidula Curtis, 1834 (7, 17, 22, 24, 26, 27, 28, 29, 35, 45, 48, 49, 51, 52)
Hydropsyche peristeriaca Botosaneanu et Marinkovic, 1966 (17)
Hydropsyche saxonica Mac Lachlan, 1884 (2, 17, 27, 48, 49, 50)
Hydropsyche tabacarui Botosaneanu, 1960 (17)
Hydropsyche tjederi Botosaneanu et Marinkovic, 1966 (17)
Cheumatopsyche lepida Pictet, 1834 (2, 17, 45, 46, 48)

POLYCENTROPODIDAE

Neureclipsis bimaculata Linne, 1758 (17, 51)
Plectrocnemia minima Klapalek, 1899 (17)
Plectrocnemia conspersa Curtis, 1834 (35, 38, 40)

Plectrocnemia geniculata Mac Lachlan, 1871 (40, 46)
Polycentropus irroratus Curtis, 1835 (25, 27)
Polycentropus flavomaculatus Pictet, 1834 (17, 22, 30, 35, 38, 40, 42, 48)
Polycentropus excisus Klapalek, 1894 (20)
Holocentropus stagnalis Albarda, 1874 (17)
Cyrnus flavidus Mac Lachlan, 1864 (49)
Cyrnus crenaticornis Kolenati, 1859 (17, 42)
Cyrnus trimaculatus Curtis, 1834 (17)

PSYCHOMYIDAE

Psychomyia pusilla Fabricius, 1781 (17, 35, 40, 42)
Lype reducta Hagen, 1860 (17)
Metalype fragilis Pictet, 1834 (20)
Tinodes pallidulus Mac Lachlan, 1878 (17)
Tinodes rostocki Mac Lachlan, 1878 (17)
Tinodes unicolor Pictet, 1834 (17)

ECNOMIDAE

Ecnomus tenellus Rambur, 1842 (17, 42)

PHRYGANEIDAE

Phryganea grandis Linne, 1758 (7, 17, 42)

BRACHYCENTRIDAE

Brachycentrus montanus Klapalek, 1891 (17)
Micrasema minimum Mac Lachlan, 1876 (17, 46)
Micrasema sericeum Klapalek, 1902 (17)
Oligoplectrum maculatum Fourcroy, 1785 (20, 35, 46)

LIMNEPHILIDAE

Drusus annulatus Stephens (42)
Drusus biguttatus Pictet, 1834 (17, 38)
Drusus botosaneanui Kumanski, 1968 (20)
Drusus discolor Rambur, 1834 (17)
Drusus discophorus Radovanovic, 1942 (20)
Drusus serbicus Marinkovic, 1971 (17)
Drusus trifidus Mac Lachlan, 1868 (42)
Ecclisopteryx guttulata Pictet, 1834 (17)
Limnephilus affinis Curtis, 1834 (17)
Limnephilus auricula Curtis, 1834 (17)
Limnephilus bipunctatus Curtis, 1834 (7, 17, 35, 42)
Limnephilus extricatus Mac Lachlan, 1865 (2, 17)
Limnephilus flavicornis Fabricius, 1787 (17, 27, 35, 42)
Limnephilus fuscicornis Rambur, 1842 (20)
Limnephilus lunatus Curtis, 1834 (7, 17, 27)
Limnephilus petri Marinkovic, 1966 (17)
Limnephilus rhombicus Linne, 1758 (17, 38)
Limnephilus sparsus Curtis, 1834 (17)
Limnephilus vittatus Fabricius, 1798 (17)
Limnephilus centralis Curtis, 1834 (38, 40)

Limnephilus marmoratus Curtis (38, 40)
Grammotaulius nitidus Müller, 1764 (17, 42)
Glyptotaelius pellucidus Retzius, 1783 (17, 42, 46)
Anabolia furcata Brauer, 1857 (17)
Anabolia nervosa Curtis, 1834 (7, 17, 24, 26, 27, 28, 29, 38, 40, 46, 48, 49, 50)
Rhadicoleptus alpestris Kolenati, 1848 (17)
Parachiona picicornis Pictet, 1834 (20)
Potamophylax latipennis Curtis, 1834 (*P. cingulatus*, *P. stellatus*, *Stenophylax latipennis*) (7, 17, 26, 27, 35, 38, 40, 46, 50)
Potamophylax luctuosus Piller et Mitterpacher, 1783 (17, 40)
Potamophylax nigricornis Pictet, 1834 (17, 27, 46)
Potamophylax pallidus Klapalek, 1900 (17)
Psilopteryx montana Kumanski, 1968 (17)
Halesus digitatus Schrank, 1781 (17, 38, 40, 46)
Halesus tessellatus Rambur, 1842 (20, 38, 40)
Halesus radiatus Curtis, 1834 (*H. interpunctatus*) (26, 38, 40, 46)
Hydatophylax infumatus Mac Lachlan, 1865 (2, 20)
Stenophylax mucronatus Mac Lachlan, 1880 (7)
Stenophylax permistus Mac Lachlan, 1895 (40)
Stenophylax mitis Mac Lachlan, 1875 (17)
Stenophylax vibex speluncarum Mac Lachlan, 1875 (17, 40, 42)
Metanoea flavipennis Pictet, 1834 (*M. chapmani*) (40)
Mesophylax impunctatus Mac Lachlan, 1884 (40)
Micropterna nycterobia Mac Lachlan, 1875 (17, 27, 40, 42)
Micropterna testacea Gmelin, 1798 (27)
Micropterna sequax Mac Lachlan, 1875 (40)
Annitella triloba Marinkovic, 1955 (20)
Annitella obscurata Mac Lachlan, 1876 (27)
Allogamus auricollis Pictet, 1834 (7, 17)
Allogamus uncatius Brauer, 1857 (17)
Chaetopteryx cissylvanica Botosaneanu, 1960 (17)
Chaetopteryx stankovici Marinkovic, 1966 (17)
Chaetopteryx maxima Kumanski, 1968 (20)
Chaetopteryx schmidi Botosaneanu, 1957 (20)
Chaetopteryx villosa Fabricius, 1798 (38)
Crunoecia bosniaca Marinkovic-Gospodnetic, 1970 (20)
Crunoecia kempnyi Morton, 1901 (20)

GOERIDAE

Goera pilosa Fabricius, 1775 (2, 17, 23, 27, 26, 29, 38, 40)
Lithax obscurus Hagen, 1859 (17, 27, 42, 46)
Lithax nigra Hagen, 1859 (22, 27)
Silo nigricornis Pictet, 1834 (2)
Silo pallipes Fabricius, 1781 (2, 17, 35, 38, 40, 42, 46)
Silo piceus Brauer, 1857 (17, 40, 46)

UENOIDAE

Thremma anomalum Mac Lachlan, 1877 (17)

LEPIDOSTOMATIDAE

Lasiocephala basalis Kolenati, 1848 (17, 38)
Lepidostoma hirtum Fabricius, 1775 (27, 50)

LEPTOCERIDAE

- Athripsodes albifrons* Linne, 1758 (*A. interjectus*) (7)
Athripsodes aterrimus Stephens, 1836 (17)
Athripsodes bilineatus Linne, 1758 (17, 50)
Athripsodes commutatus Rostock, 1874 (17)
Mystacides azurea Linne, 1761 (17, 42, 46)
Mystacides nigra Linne, 1758 (17)
Mystacides longicornis Linne, 1758 (*M. concolor*, *monochroa*, *leucoptera*) (40, 46)
Ylodes simulans Tjeder, 1929 (*Trienodes forsslundi*) (17)
Trienodes kawraiskii Martynov, 1909 (17)
Setodes hungaricus Ulmer, 1908 (17)
Leptocerus interruptus Fabricius, 1775 (17)
Ceracela annulicornis Stephens, 1836 (*Leptocerus futilis*, *lugens*, *recurvatus*, *Mystacides perfusus*) (42)
Adicella balcanica Botosaneanu et Novak, 1965 (17)
Adicella filicornis Pictet, 1834 (17)
Adicella syriaca Ulmer, 1907 (17)

SERICOSTOMATIDAE

- Oecismus monedula* Hagen, 1859 (17)
Sericostoma flavicorne Schneider, 1845 (*S. timidum*, *turbatum*, *pyrenaicum*, *schneiderii*) (17)
Sericostoma personatum Spence, 1826 (*S. pedemontanum*, *memorable*)
(7, 22, 26, 27, 28, 35, 38, 40, 46, 50)
Notidobia ciliaris Linnaeus, 1761 (20, 38, 40)

BERAIDAE

- Beraea pullata* Curtis, 1834 (20, 27)
Beraea maura Curtis, 1834 (27)
Beraeodes minutus Linne, 1761 (17, 46)
Beraemyia schmidi Botosaneanu, 1960 (17)
Ernodes articularis Pictet, 1834 (*E. martynovi*) (17)

HELICOPSYCHIDAE

- Helicopsyche bacescui* Orghidan et Botosaneanu, 1957 (17)
Helicopsyche sperata Mac Lachlan (27)

ODONTOCERIDAE

- Odontocerum albicorne* Scopoli, 1763 (7, 17, 27, 35, 38, 40, 46, 50)
Odontocerum hellenicum Malicky, 1972 (20)

References

- ARSOV, G. (1991). Taxonomico-bioecoenological analsis and altitude distribution on trichoptera larva fauna in the Zrnovska river. M.S.thesis. Faculty of Sciences, Skopje, 1–148. (in Macedonian)
BARACKOV, Z. (1973). Ecological investigations on fauna of the bottom of the Grosnicka River. Faculty of Sciences in Kragujevac, M.S. thesis, 1–100. (in Serbian)
BOTOSANEANU, L. (1960). Trichopteres de Yugoslavie recueillis en 1955 per le Dr. F. Schmid, D.E.Z. (N.F.), 7, 261–293.
HABDJIA, I. (1979). Larvae of Trichoptera as indicators of ecological conditions in benthos of karst waters. Proceedings, II congress of ecologists of Yugoslavia, Zadar, 1433–1446. (in Croatian)
HOLZENTHAL, R.W. AND BLAHNIK, R.J. (2000). Trichoptera Caddisflies.
<http://phylogeny.arizona.edu/tree/eukaryotes/animals/arthropoda/hexapoda/trichoptera/trichoptera.html>

- ILLIES, J. (1978). *Limnofauna Europaea* – Gustav Fischer Verlag, Stuttgart, 532.
- KONTA, P. S. (1997). The analysis of the influence of ecological factors on the macrozoobenthos of the Lomnicka Reka River. Biological Faculty in Belgrade, M.S. thesis. 1–99. (in Serbian)
- KRUSNIK, C. (1987). Trichoptera (Insecta). CANU, separate issue, book 21, Department of sciences, 13: The Fauna of Durmitor, notebook 2, 201–224. (in Serbian)
- KUMANSKI, K. P. (1985). Fauna in Balcia, Trichoptera – Annulipalpia. BAH, Sofia, 243. (in Bulgarian)
- KUMANSKI, K. P. (1988). Fauna na Balcia, Trichoptera – Integripalpia. BAH, Sofia, 353. (in Bulgarian)
- MALICKY, H. (1982). Zwei neues Köcherfliegen (Trichoptera) aus dem Mediterranean Gebirgen. *Entomologische Zeitschrift*, 92 (12), 161–163.
- MALICKY, H. (1983). *Atlas of European Trichoptera*. Serias Entomological, Vol. 24. The Hague–Boston–London, Junk, 1–298.
- MARINKOVIC-GOSPODNETIC, M. (1966). New species of Trichoptera from Yugoslavia. *Bulletin Scientifique*, Sec. A-Tome 11, No. 4–6.
- MARINKOVIC-GOSPODNETIC, M. (1970). Descriptions of some species of Trichoptera from Yugoslavia. *Annual of the Biological Institute of the University in Sarajevo*, vol. XXIII, 77–84.
- MARINKOVIC-GOSPODNETIC, M. (1970a). Fauna of Trichoptera of the Maglic, Volujak and Zelengor regions. *Bull. Mus. Nat. Hist. in BiH in Sarajevo*. Sciences, issue IX, 107–120. (in Serbian)
- MARINKOVIC-GOSPODNETIC, M. (1971). The species of the genus *Drusus* in Yugoslavia. *Annual of the Biological Institute of the University in Sarajevo*, vol. XXIV, 105–109.
- MARINKOVIC-GOSPODNETIC, M. (1975). Fauna Trichoptera SR Serbia. *Book of abstracts on entomofauna in Serbia 1*, 219–236. (in Serbian)
- MARINKOVIC-GOSPODNETIC, M. (1978). The Caddis-Flies (Trichoptera, Insecta) of Hercegovina (Yugoslavia). *Annual of the Biological Institute of the University in Sarajevo*, vol. XXXI, 115–131.
- MARINKOVIC-GOSPODNETIC, M. (1979). The Trichoptera (Insecta) of the large krastic Spring in the Dinarides. *Proceedings, II congress of ecologists of Yugoslavia, Zadar, 1837–1849*. (in Serbian)
- MARINKOVIC-GOSPODNETIC, M. (1980). Fauna Trichoptera SR Serbia. *Book of abstracts on fauna in Serbia 1*, 71–84. (in Serbian)
- MARINKOVIC-GOSPODNETIC, M. (1981). Trichoptera of the Morava and Plavnica Rivers drainages (in *The Biota and limnology of lake Skadar*). Titograd, 307–309.
- MARKOVIC, Z. (1995). The Detinja river, macrozoobenthos in evaluation of the quality of water – Ministry for the protection of the environment in the Republic of Serbia and the Center for Scientific Explorations U`ice, 1–131. (in Serbian)
- MARKOVIC, Z. and MILJANOVIC, B. (1995). Macrozoobenthos of the Kriveljska Reka river. *Book of abstracts "Nasa ekoloska istina"*, 221–225. (in Serbian)
- MARKOVIC, Z., MILJANOVIC, B., MITROVIC-TUTUNDIC, V. (1997). Saprobiological evaluation of the Banja Reka river and its tributary the Pocibrava by the use of macrozoobenthos as indicator. *Annual proceedings of the Yugoslav society for water protection*, 350–354. (in Serbian)
- MARKOVIC, Z., MITROVIC-TUTUNDZIC, V., SAVIC, I., RANDJELOVIC, N. (1997a) Analysis of spring to stream fauna transition by example of the river Banja source (Serbia, Yugoslavia). *Ekologija* 32 (2): 57–63.
- MARKOVIC, Z., MITROVIC-TUTUNDZIC, V., MILJANOVIC, B. (1997b). Effect of pollution on the macrozoobenthos diversity and structure in the river Obnica river (Serbia, Yugoslavia). *Ekologija* 32 (2): 37–46.
- MARKOVIC, Z. (1998). Springs in mountainous regions of Serbia: Ecological study of the macrozoobenthos. *Faculty of Biology, University of Belgrade*, 1–318. (in Serbian)
- MARKOVIC, Z., MILJANOVIC, B., MITROVIC-TUTUNDZIC, V. (1998). Macrozoobenthos as a water quality parameter in the Jablanica River. *Annual proceedings of the Yugoslav society for water protection*, 369–372. (in Serbian)
- MARKOVIC, Z., MILJANOVIC, B., MITROVIC-TUTUNDZIC, V. (1999). Macrozoobenthos as an indicator of the Kolubara Reka river water quality. *Annual proceedings of the Yugoslav society for water protection*, 261–266. (in Serbian)
- MARTINOVIC-VITANOVIC, V., KALAFATIC, V., JAKOVCEV, D., PAUNOVIC, M., I MARTINOVIC, M. J. (1998). The Veternica river – composition and structure of biocenosis, saprobiological status and quality of water. *Ecologica* 5, 3, 22–28. (in Serbian)
- MATONICKIN, I. (1959). Trihopter fauna and its relation to the speed of water in seder pools and their rapids. *Biological archives*, 12, 3–4, 97–104. (in Croatian)
- MATONICKIN, I., PAVLETIC, Z., ZUNJIC, K., HABDIJA, I. (1975). Ecosystem of the Piva river and biological valorization of its waters. *Archives of Republ. Institute for Natural Protection – Museum of Natural Protection Titograd*, br. 8, 61–79. (in Croatian)

- MATONICKIN, I. (1987). Material for the limnofauna of karst the running waters of Croatia. *Biosystematics*, Vol. 13, No. 1, 25–35. (in Croatian)
- MILJANOVIC, B. (2000). Biodiversity of macrozoobenthos of the upper course of the Kolubare. M.Sc.thesis. Institute of biology, Faculty of Sciences, University in Novi Sad, 1–156. (in Serbian)
- PAUNOVIC, M. (2001). Spatial and temporal dynamics of the macrozoobenthos of the Vlasina River. Biological Faculty in Belgrade, M.S. thesis. 1–200. (in Serbian)
- PAVLETIC, Z. and MATONICKIN, I. (1972). Biocenological structure in the Korana river as a reflection of the quality of water. *Ecologica*, Vol. 7, No. 1–2, 59–79. (in Croatian)
- PAVLOVSKI, T. (1991). Composition, dynamics and distribution of larvae fauna of Trichoptera in the Babuna river. Special issue., *Inst. Biol., Faculty of Sciences, Skopje* 7–82. (in Macedonian)
- RADOVANOVIC, M. (1931). The results of exploration of Triholoptera in the Balkans. Annual proceedings of the Yugoslav entomological society, year V–VI, notebook 1–2, 159–192. (in Serbian)
- RADOVANOVIC, M. (1933). Triholoptera in Slovenia. *Prirodoslovne Razprave*, 2, 112–121. (in Serbian)
- RADOVANOVIC, M. (1935). Trichoptera in Yugoslavia. *Bull. Mus. Nat. Hist. in BiH in Sarajevo*. Vol. XLVII, 73–84. (in Serbian)
- RADOVANOVIC, M. (1943). Triholoptera of Lake Ohrid and adjacent region. SANU, Special issue, book CXXXV, biological and mathematical papers, book 32, Ohrid proceedings, book 1, 3–46. (in Serbian)
- RADOVANOVIC, M. (1953). Contribution to the study of Trichoptera in the Balkan peninsula, primarily in caves and mountain lakes. *Bull. SAN CCX* (7): 11–38. (in Serbian)
- ROSS, H. (1967). The Evolution and Past Dispersal of the Trichoptera. *Ann. Rev. Ent.* 12, 169–202.
- SCHMERA, D. (2001). Check List of the Hungarian Trichoptera/Magyar Trichoptera fauna. <http://julianki.hu/trichoptera/species.htm>
- SHUKRIU, A. (1979). Ecological causes and zonal distribution of macrozoobenthos in Prizrenka Bistrica River, 6–124, Zagreb. (in Serbian)
- SIMIC, V. (1993). Saprobilogic evaluation of the Svrlijski and Trgoviski Timok rivers on the basis of the composition of macrozoobenthos. Biological Faculty in Belgrade, M.S. thesis. 1–263. (in Serbian)
- SIMIC, V. and OSTOJIC, A. (1994). The macrozoobenthos of Lake Vlasina. River. *Annul proceedings of the Yugoslav society for water protection*, 116–121. (in Serbian)
- STRAHINIC, I. (2000). Faunistical and ecological study of the macrozoobenthos of the Pusta Reka river. Biological Faculty in Belgrade, M.Sc.thesis. 1–185. (in Serbian)
- ŽIVIĆ, I., MARKOVIĆ, Z. and BRAJKOVIĆ, M. (2001). Contribution to the knowledge of qualitative composition of macrozoobenthos and quality of water of the Crvena Reka river. *Annul proceedings of the Yugoslav society for water protection*, 175–182. (in Serbian)
- ŽIVIĆ, I., MARKOVIĆ, Z. and BRAJKOVIĆ, M. (2001a). A contribution to the study of the Trichoptera (Insecta) fauna in the Toplica river. *Acta entomologica serbica*, 5 (1/2): 35–46. (in Serbian)
- ŽIVIĆ, I., MARKOVIĆ, Z. and BRAJKOVIĆ, M. (2001b). Fauna of the bottom of the Kudoski stream. *Protection of nature*, 53 (1): 79–87. (in Serbian)
- ŽIVIĆ, I., MARKOVIĆ, Z. and BRAJKOVIĆ, M. (2000a). The change of the structure of macrozoobenthos in the Jelenacki stream under the influence of pollution. *Ekologija*, Vol.35 (2): 105–114.

Ivana ŽIVIĆ, and Miloje BRAJKOVIĆ
 Institute of Zoology, Faculty of Biology,
 Studentski Trg 16,
 YU-11000 Belgrade, Yugoslavia:
 tel.+381-11-187266,
 e-mail: ivanas@bf.bio.bg.ac.yu

Zoran MARKOVIĆ
 Institute of Cattle Breeding, Faculty of Agriculture,
 Nemanjina 6,
 YU-11000 Belgrade, Yugoslavia:
 tel.+381-11-615315, lok.290