

FEKETE-SZEGÖ FUNCTIONAL FOR NON-BAZILEVIČ FUNCTIONS

NIKOLA TUNESKI AND MASLINA DARUS

ABSTRACT. Let $f(z) = z + a_2z^2 + a_3z^3 + \dots$ be an analytic function in the unit disk $\mathcal{U}$ and let the class of non-Bazilevič functions, for $0 < \lambda < 1$, be described with $\operatorname{Re} \left\{ f'(z) (z/f(z))^{1+\lambda} \right\} > 0, z \in \mathcal{U}$. In this paper we obtain sharp upper bound of $|a_2|$ and of the Fekete-Szegő functional $|a_3 - \mu a_2^2|$ for the class of non-Bazilevič functions and for some of its subclasses.

1. INTRODUCTION AND PRELIMINARIES

Let $\mathcal{A}$ denote the class of analytic functions in the unit disk $\mathcal{U} = \{z : |z| < 1\}$ normalized such that $f(0) = f'(0) - 1 = 0$, i.e., of type $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$.

A function $f \in \mathcal{A}$ is said to be of *Bazilevič type* if for a starlike function g ($g \in \mathcal{A}$ is starlike if and only if $\operatorname{Re} \{zg'(z)/g(z)\} > 0, z \in \mathcal{U}$) we have

$$\operatorname{Re} \{f'(z)(f(z)/z)^{\alpha+i\gamma-1}(g(z)/z)^{-\alpha}\} > 0,$$

$z \in \mathcal{U}$ (see more in [1]). This class and its subclasses were widely studied in the past decades. Specially, in [4] sharp upper bound of the Fekete-Szegő functional $|a_3 - \mu a_2^2|$ is obtained for all real μ when $\gamma = 0$. That result was partially extended in [2] to a wider subclass satisfying

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f^{1-\alpha}(z)g^\alpha(z)} \right\} > \beta, \quad z \in \mathcal{U},$$

where $\alpha > 0$ and $0 \leq \beta < 1$.

In [5], Obradović introduced a class of functions $f \in \mathcal{A}$ that for $0 < \lambda < 1$ is defined by

$$\operatorname{Re} \{f'(z)(z/f(z))^{1+\lambda}\} > 0, \quad z \in \mathcal{U}.$$

Recently, in his talk at the Conference ‘Computational Methods and Function Theory 2001’, he called this functions to be of *non-Bazilevič type*. By now, this class was studied in a direction of finding necessary conditions over λ that embeds this class into the class of univalent function or its subclasses, which is still an open problem. Here we will find sharp upper bound of $|a_2|$ and of the Fekete-Szegő functional $|a_3 - \mu a_2^2|$ for the class of non-Bazilevič functions and for some its subclasses. In that purpose we will need the following lemma.

Lemma 1. ([6], p.166, formula (10)) ([3], p.41) *Let $p \in \mathcal{P}$, that is, p be analytic in $\mathcal{U}$, be given by $p(z) = 1 + \sum_{n=1}^{\infty} p_n z^n$ and $\operatorname{Re} p(z) > 0$ for $z \in \mathcal{U}$. Then*

$$|p_2 - p_1^2/2| \leq 2 - |p_1|^2/2$$

and $|p_n| \leq 2$ for all $n \in \mathbb{N}$.

2000 *Mathematics Subject Classification.* 30C50, 30C45.

Key words and phrases. non-Bazilevič function, Fekete-Szegő functional, sharp upper bound.

2. MAIN RESULTS

Theorem 1. Let $f \in \mathcal{A}$, $0 < \lambda < 1$ and $0 \leq \alpha < 1$. If

$$(1) \quad \operatorname{Re} \{f'(z)(z/f(z))^{1+\lambda}\} > \alpha, \quad z \in \mathcal{U},$$

then $|a_2| \leq 2(1-\alpha)/(1-\lambda)$ and for all $\mu \in \mathbb{C}$ the following bound is sharp

$$|a_3 - \mu a_2^2| \leq \frac{2(1-\alpha)}{2-\lambda} \max \left\{ 1, \left| 1 + \frac{2(1+\lambda-\mu)(2-\lambda)(1-\alpha)}{(1-\lambda)^2} \right| \right\}.$$

Proof. Condition (1) is equivalent to

$$f'(z) = (f(z)/z)^{1+\lambda}[(1-\alpha)p(z) + \alpha], \quad z \in \mathcal{U},$$

for some $p \in \mathcal{P}$. Equating coefficients we obtain $a_2 = p_1(1-\alpha)/(1-\lambda)$,

$$a_3 = \frac{1-\alpha}{2-\lambda}p_2 + \frac{(1-\alpha)^2(1+\lambda)}{2(1-\lambda)^2}p_1^2$$

and further

$$\begin{aligned} a_3 - \mu a_2^2 &= \frac{1-\alpha}{2-\lambda} \left(p_2 - \frac{1}{2}p_1^2 \right) + \\ &+ \frac{(1-\alpha)(1-\lambda)^2 + (1-\alpha)^2(1+\lambda-2\mu)(2-\lambda)}{2(2-\lambda)(1-\lambda)^2} p_1^2. \end{aligned}$$

Now, using Lemma 1 we receive $|a_3 - \mu a_2^2| \leq H(x) = A + ABx^2/4$ where $x = |p_1| \leq 2$, $A = 2(1-\alpha)/(2-\lambda) > 0$, $B = (|C| - (1-\lambda)^2)/(1-\lambda)^2$ and $C = (1-\lambda)^2 + (1-\alpha)(1+\lambda-2\mu)(2-\lambda)$. So, we have

$$|a_3 - \mu a_2^2| \leq \begin{cases} H(0) = A, & |C| \leq (1-\lambda)^2 \\ H(2) = A|C|/(1-\lambda)^2, & |C| \geq (1-\lambda)^2 \end{cases}.$$

Equality is attained for functions given by

$$f'(z) \left(\frac{z}{f(z)} \right)^{1+\lambda} = \frac{1+z^2(1-2\alpha)}{1-z^2}$$

and

$$f'(z) \left(\frac{z}{f(z)} \right)^{1+\lambda} = \frac{1+z(1-2\alpha)}{1-z}$$

respectively. □

For $\alpha = 0$ we have the following corollary.

Corollary 1. Let $f \in \mathcal{A}$ and $0 < \lambda < 1$. If

$$\operatorname{Re} \{f'(z)(z/f(z))^{1+\lambda}\} > 0, \quad z \in \mathcal{U},$$

then $|a_2| \leq 2/(1-\lambda)$ and for all $\mu \in \mathbb{C}$ the following bound is sharp

$$|a_3 - \mu a_2^2| \leq \frac{2}{2-\lambda} \max \left\{ 1, \left| 1 + \frac{(1+\lambda-2\mu)(2-\lambda)}{(1-\lambda)^2} \right| \right\}.$$

Now we will consider one subclass of the class of non-Bazilevič function.

Theorem 2. Let $f \in \mathcal{A}$, $0 < \lambda < 1$ and $0 < k \leq 1$. If

$$(2) \quad |f'(z)(z/f(z))^{1+\lambda} - 1| < k, \quad z \in \mathcal{U},$$

then $|a_2| \leq k/(1-\lambda)$ and for all $\mu \in \mathbb{C}$ the following bound is sharp

$$|a_3 - \mu a_2^2| \leq \frac{k}{(2-\lambda)} \max \left\{ 1, \frac{k(2-\lambda)}{(1-\lambda)^2} \left| \frac{1+\lambda}{2} - \mu \right| \right\}.$$

Proof. Similarly as in the proof of Theorem 1, condition (2) implies that there exists a function $p \in \mathcal{P}$ such that for all $z \in \mathcal{U}$

$$f'(z) = (f(z)/z)^{1+\lambda} (2k/(1+p(z)) + 1 - k).$$

Equating the coefficients we obtain $a_2 = -kp_1/(2(1-\lambda))$,

$$a_3 = \frac{k^2}{8} \frac{1+\lambda}{(1-\lambda)^2} p_1^2 - \frac{k}{2(2-\lambda)} \left(p_2 - \frac{p_1^2}{2} \right)$$

and

$$a_3 - \mu a_2^2 = -\frac{k}{2(2-\lambda)} \left(p_2 - \frac{p_1^2}{2} \right) + \frac{k^2 p_1^2}{4(1-\lambda)^2} \left(\frac{1+\lambda}{2} - \mu \right).$$

So, $|a_3 - \mu a_2^2| \leq H(x) = A + Bx^2/4$ where $x = |p_1| \leq 2$, $A = k/(2-\lambda) > 0$, $B = k^2|C|/(1-\lambda)^2 - k/(2-\lambda)$ and $C = (1+\lambda)/2 - \mu$. Therefore

$$|a_3 - \mu a_2^2| \leq \begin{cases} H(0) = A, & |C| \leq (1-\lambda)^2/(k(2-\lambda)) \\ H(2) = Ak(2-\lambda)|C|/(1-\lambda)^2, & |C| \geq (1-\lambda)^2/(k(2-\lambda)) \end{cases}.$$

Here equality is attained for the functions given by $f'(z)(z/f(z))^{1+\lambda} = 1 - kz^2$ and $f'(z)(z/f(z))^{1+\lambda} = 1 - kz$, respectively. $\square$

For $k = 1$ we receive the following corollary.

Corollary 2. *Let $f \in \mathcal{A}$ and $0 < \lambda < 1$. If*

$$|f'(z)(z/f(z))^{1+\lambda} - 1| < 1, \quad z \in \mathcal{U},$$

then $|a_2| \leq 1/(1-\lambda)$ and for all $\mu \in \mathbb{C}$ the following bound is sharp

$$|a_3 - \mu a_2^2| \leq \frac{1}{(2-\lambda)^2} \max \left\{ 1, \frac{2-\lambda}{(1-\lambda)^2} \left| \frac{1+\lambda}{2} - \mu \right| \right\}.$$

REFERENCES

- [1] I.E. Bazilevič. On a case of integrability in quadratures of the Loewner-Kufarev equation. *Mat. Sb.*, 37:471–476, 1955. Russian.
- [2] M. Darus. Coefficient problem for Bazilevič functions. *J. Inst. Math. Comput. Sci. Math. Ser.*, 2, 2000.
- [3] P.L. Duren. *Univalent functions*. Springer-Verlag, 1983.
- [4] P.J. Eenigenburg and E.M. Silvia. A coefficient inequality for Bazilevič functions. *Annales Univ. Mariae Curie-Skłodowska*, 27:5–12, 1973.
- [5] M. Obradović. A class of univalent functions. *Hokkaido Math. J.*, 2:329–335, 1988.
- [6] C. Pommerenke. *Univalent Functions*. Studia Mathematica Mathematische Lehrbücher. Vandenhoeck & Ruprecht, 1975. With a chapter on quadratic differentials by Gerd Jensen.

Received March 25, 2002.

FACULTY OF MECHANICAL ENGINEERING, KARPOŠ II B.B.,
1000 SKOPJE, REPUBLIC OF MACEDONIA
E-mail address: nikolat@ereb1.mf.ukim.edu.mk

SCHOOL OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCES AND TECHNOLOGY,
UNIVERSITY KEBANGSAAN MALAYSIA, BANGI 43600 SELANGOR, MALAYSIA
E-mail address: maslina@pkrisc.cc.ukm.my