

A STUDY ON THE GENERALIZATION OF JANOWSKI FUNCTIONS IN THE UNIT DISC

Y. POLATOĞLU, M. BOLCAL, A. ŞEN AND E. YAVUZ

ABSTRACT. Let Ω be the class of functions $w(z)$, $w(0) = 0$, $|w(z)| < 1$ regular in the unit disc $D = \{z : |z| < 1\}$. For arbitrarily fixed numbers $A \in (-1, 1]$, $B \in [-1, A]$, $0 \leq \alpha < 1$ let $P(A, B, \alpha)$ be the class of regular functions $p(z)$ in D such that $p(0) = 1$, and which is $p(z) \in P(A, B, \alpha)$ if and only if $p(z) = \frac{1 + [(1-\alpha)A + \alpha B]w(z)}{1 + Bw(z)}$ for some function $w(z) \in \Omega$ and every $z \in D$.

In the present paper we apply the principle of subordination ([1], [3], [4], [5]) to give new proofs for some classical results concerning the class $S^*(A, B, \alpha)$ of functions $f(z)$ with $f(0) = 0$, $f'(0) = 1$, which are regular in D satisfying the condition: $f(z) \in S^*(A, B, \alpha)$ if and only if $z \frac{f'(z)}{f(z)} = p(z)$ for some $p(z) \in P(A, B, \alpha)$ and for all z in D .

1. INTRODUCTION

Let Ω be the family of functions $w(z)$ regular in the unit disc D and satisfying the conditions $w(0) = 0$, $|w(z)| < 1$, for $z \in D$.

For arbitrary fixed numbers A, B, α , $-1 \leq B < A \leq 1$, $0 \leq \alpha < 1$, let $P(A, B, \alpha)$ denote the family of functions

$$(1) \quad p(z) = 1 + p_1 z + p_2 z^2 + \cdots + p_n z^n + \cdots$$

regular in D and such that $p(z)$ is in $P(A, B, \alpha)$ if and only if

$$(2) \quad p(z) \prec \frac{1 + [(1-\alpha)A + \alpha B]z}{1 + Bz} \Leftrightarrow p(z) = \frac{1 + [(1-\alpha)A + \alpha B]w(z)}{1 + Bw(z)}$$

for some function $w(z) \in \Omega$ and every $z \in D$.

Furthermore, let $S^*(A, B, \alpha)$ denote the family of functions

$$(3) \quad f(z) = z + a_2 z^2 + a_3 z^3 + \cdots$$

regular in D and such that $f(z)$ is in $S^*(A, B, \alpha)$ if and only if

$$(4) \quad z \frac{f'(z)}{f(z)} = p(z)$$

for some $p(z)$ in $P(A, B, \alpha)$ and for all z in D .

2000 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Starlike functions, distortion theorem, representation theorem, radius of starlikeness.

2. NEW RESULTS ON THE CLASS $S^*(A, B, \alpha)$

In this section we shall give representation theorems, distortion theorems and establish the radius of starlikeness for the class $S^*(A, B, \alpha)$. Our proofs are based on I.S. Jack's Lemma [2].

Lemma 1. *Let $w(z)$ be a non-constant and analytic function in the unit disc D with $w(0) = 0$. If $|w(z)|$ attains its maximum value on the circle $|z| = r$ at the point z_1 , then $z_1 w'(z_1) = k w(z_1)$ and $k \geq 1$.*

From the definition of the class $P(1, -1, 0)$ called the Caratheodory class and $P(A, B, \alpha)$ we easily obtain the following lemma.

Lemma 2. *If $p(z) \in P(A, B, \alpha)$ if and only if*

$$(5) \quad p(z) = \frac{[1 + (1 - \alpha)A + \alpha B]q(z) + [(1 - \alpha)(A + B)]}{[1 + B]q(z) + [1 - B]}$$

for some $q(z) \in P(1, -1, 0)$.

Let ζ be an arbitrary fixed point of D . We consider the functional

$$(6) \quad F(p) = p(\zeta), p(z) \in P(A, B, \alpha).$$

Lemma 3. *The set of the values of the functional (6) is the closed disc with centered at $C(r)$ and having the radius $\rho(r)$, where*

$$\begin{cases} C(r) = \left(\frac{1-B[(1-\alpha)A+\alpha B]r^2}{1-B^2r^2}, 0 \right), & \rho(r) = \frac{(1-\alpha)(A-B)r}{1-B^2r^2}, \quad B \neq 0, \\ C(r) = (1, 0), & \rho(r) = (1 - \alpha) |A| r, \quad B = 0. \end{cases}$$

Proof. Every boundary function $p_0(z)$ of $P(A, B, \alpha)$ with respect to the functional (6) can be written in the form (5), where

$$q(z) = \frac{1 + \varepsilon z}{1 - \varepsilon z}, |\varepsilon| = 1.$$

Hence

$$(7) \quad p_0(z) = \frac{1 + [(1 - \alpha)A + \alpha B]z}{1 + Bz}.$$

Since $z = re^{i\theta}$, $0 \leq \theta \leq 2\pi$,

$$(8) \quad \begin{aligned} p_0(z) &= C(r) + \rho\eta, \\ \eta &= \varepsilon e^{i\theta} \frac{1 + Br\bar{\varepsilon}e^{-i\theta}}{1 + Br\varepsilon e^{i\theta}}, \end{aligned}$$

which completes the proof. □

Lemma 4. *The function*

$$w = w(z) = \begin{cases} \frac{(1-\alpha)(A-B)z}{1+Bz}, & B \neq 0, \\ (1 - \alpha)Az, & B = 0, \end{cases}$$

maps $|z| = r$ onto the disc centered at $C(r)$, and having the radius $\rho(r)$

$$\begin{cases} C(r) = \left(-\frac{B(1-\alpha)(A-B)r^2}{1-B^2r^2}, 0 \right), & \rho(r) = \frac{(1-\alpha)(A-B)r}{1-B^2r^2}, \quad B \neq 0, \\ C(r) = (0, 0), & \rho(r) = (1 - \alpha) |A| r, \quad B = 0. \end{cases}$$

Proof. This is immediate from

$$(9) \quad \begin{aligned} w &= \frac{(1-\alpha)(A-B)z}{1+Bz} \Rightarrow \\ u^2 + v^2 + \frac{2B(1-\alpha)(A-B)r^2}{1-B^2r^2}u - \frac{(1-\alpha)^2(A-B)^2r^2}{1-B^2r^2} &= 0, B \neq 0, \\ w &= (1-\alpha)Az \Rightarrow u^2 + v^2 - (1-\alpha)^2A^2r^2 = 0, B = 0. \end{aligned}$$

□

Theorem 1. Let $f(z) = z + a_2z^2 + \dots$ be an analytic function in the unit disc D . If $f(z)$ satisfying

$$(10) \quad \left(z \frac{f'(z)}{f(z)} - 1 \right) \prec \begin{cases} \frac{(1-\alpha)(A-B)z}{1+Bz} = F_1(z), & B \neq 0, \\ (1-\alpha)Az = F_2(z), & B = 0, \end{cases}$$

then $f(z) \in S^*(A, B, \alpha)$ and this result is as sharp as the function

$$\left(\frac{1 + [(1-\alpha)A + B\alpha]z}{1+Bz} \right).$$

Proof. We define the function $w(z)$ by

$$(11) \quad \frac{f(z)}{z} = \begin{cases} (1+Bw(z))^{\frac{(1-\alpha)(A-B)}{B}}, & B \neq 0, \\ e^{(1-\alpha)Aw(z)}, & B = 0, \end{cases}$$

where $(1+Bw(z))^{\frac{(1-\alpha)(A-B)}{B}}$ and $e^{(1-\alpha)Aw(z)}$ have the value 1 at the origin. Then $w(z)$ is analytic in D and $w(0) = 0$. If we take the logarithmic derivate of equality (11), simple calculations yield

$$(12) \quad \left(z \frac{f'(z)}{f(z)} - 1 \right) = \begin{cases} \frac{(1-\alpha)(A-B)zw'(z)}{1+Bw(z)}, & B \neq 0, \\ (1-\alpha)Aw'(z), & B = 0. \end{cases}$$

Now it is easy to realize that the subordination (10) is equivalent to $|w(z)| < 1$ for all $z \in D$ indeed assume the contrary. There exist $z_1 \in D$ such that $|w(z_1)| = 1$. Then by I.S. Jack's Lemma $z_1w'(z_1) = kw(z_1)$ and $k \geq 1$, for such $z_1 \in D$ and using Lemma 4 we have

$$(13) \quad \left(z_1 \frac{f'(z_1)}{f(z_1)} - 1 \right) = \begin{cases} \frac{(1-\alpha)(A-B)kw(z_1)}{1+Bw(z_1)} = F_1(w(z_1)) \notin F_1(D), & B \neq 0, \\ (1-\alpha)Aw'(z_1) = F_2(w(z_1)) \notin F_2(D), & B = 0, \end{cases}$$

because $|w(z_1)| = 1$ and $k \geq 1$. But this contradicts condition (10) of this theorem and so $|w(z)| < 1$ for all $z \in D$. By using condition (10) we get

$$z \frac{f'(z)}{f(z)} = \begin{cases} \frac{1 + [(1-\alpha)A + \alpha B]w(z)}{1+Bw(z)}, & B \neq 0, \\ 1 + (1-\alpha)Aw(z), & B = 0, \end{cases}$$

which ends the proof. □

Corollary 1. Let $f(z) \in S^*(A, B, \alpha)$. Then $f(z)$ can be written in the form

$$f(z) = \begin{cases} z(1+Bw(z))^{\frac{(1-\alpha)(A-B)}{B}}, & B \neq 0, \\ ze^{(1-\alpha)Aw(z)}, & B = 0. \end{cases}$$

Theorem 2. If $f(z) \in S^*(A, B, \alpha)$, then

$$(14) \quad \begin{cases} r(1-Br)^{\frac{(1-\alpha)(A-B)}{B}} \leq |f(z)| \leq r(1+Br)^{\frac{(1-\alpha)(A-B)}{B}}, & B \neq 0, \\ re^{-(1-\alpha)|A|r} \leq |f(z)| \leq re^{(1-\alpha)|A|r}, & B = 0. \end{cases}$$

These bounds are sharp with the extremal function

$$(15) \quad f_*(z) = \begin{cases} z(1+Bz)^{\frac{(1-\alpha)(A-B)}{B}}, & B \neq 0, \\ ze^{(1-\alpha)Az}, & B = 0. \end{cases}$$

Proof. The set of the values of $\left(z \frac{f'(z)}{f(z)}\right)$ is the closed disc with centered at $C(r) = \frac{1-B[A(1-\alpha)+B\alpha]r^2}{1-B^2r^2}$ and having the radius $\rho(r) = \frac{(1-\alpha)(A-B)r}{1-B^2r^2}$ by using Lemma 3, that is

$$(16) \quad \left| z \frac{f'(z)}{f(z)} - \frac{1-B[(1-\alpha)A+\alpha B]r^2}{1-B^2r^2} \right| \leq \frac{(1-\alpha)(A-B)r}{1-B^2r^2}.$$

After simple calculations from (16) we get

$$(17) \quad \begin{cases} \frac{1-(1-\alpha)(A-B)r-B[(1-\alpha)A+\alpha B]r^2}{1-B^2r^2} \leq \operatorname{Re} \left(z \frac{f'(z)}{f(z)} \right) \\ \leq \frac{1+(1-\alpha)(A-B)r-B[(1-\alpha)A+\alpha B]r^2}{1-B^2r^2}, & B \neq 0, \\ 1-(1-\alpha)|A|r \leq \operatorname{Re} \left(z \frac{f'(z)}{f(z)} \right) \leq 1+(1-\alpha)|A|r, & B = 0. \end{cases}$$

On the other hand we have

$$(18) \quad \operatorname{Re} \left(z \frac{f'(z)}{f(z)} \right) = r \frac{\partial}{\partial r} \log |f(z)|, |z| = r.$$

If we substitute (18) into the (17) we get

$$(19) \quad \begin{cases} \frac{1}{r} - \frac{(1-\alpha)(A-B)}{1-Br} \leq \frac{\partial}{\partial r} \log |f(z)| \leq \frac{1}{r} + \frac{(1-\alpha)(A-B)}{1+Br}, & B \neq 0, \\ \frac{1}{r} - (1-\alpha)|A| \leq \frac{\partial}{\partial r} \log |f(z)| \leq \frac{1}{r} + (1-\alpha)|A|, & B = 0. \end{cases}$$

Integrating both sides (19) we obtain (14). $\square$

Corollary 2. The radius of starlikeness of the class $S^*(A, B, \alpha)$ is

$$(20) \quad r_s = \frac{2}{(1-\alpha)(A-B) + \sqrt{(1-\alpha)^2(A-B)^2 + 4B[(1-\alpha)A + \alpha B]}}.$$

This radius is sharp because the extremal function is given in (15).

Proof. From (17) we have

$$(21) \quad \operatorname{Re} \left(z \frac{f'(z)}{f(z)} \right) \geq \frac{1-(1-\alpha)(A-B)r-B[(1-\alpha)A+\alpha B]r^2}{1-B^2r^2}.$$

Hence for $r < r_s$ the first hand side of the preceding inequality is positive this implies that

$$r_s = \frac{2}{(1-\alpha)(A-B) + \sqrt{(1-\alpha)^2(A-B)^2 + 4B[(1-\alpha)A + \alpha B]}}.$$

Also note that the inequality (20) becomes an equality for the function which is given in (15). It follows that

$$r_s = \frac{2}{(1-\alpha)(A-B) + \sqrt{(1-\alpha)^2(A-B)^2 + 4B[(1-\alpha)A + \alpha B]}}.$$

and the proof is complete. $\square$

Acknowledgements. The authors thank the referee for their useful suggestions.

REFERENCES

- [1] A. W. Goodman. *Univalent functions*. Mariner Publishing Co. Inc., Tampa, FL, 1983.
- [2] I. S. Jack. Functions starlike and convex of order α . *J. London Math. Soc. (2)*, 3:469–474, 1971.
- [3] W. Janowski. Some extremal problems for certain families of analytic functions. I. *Ann. Polon. Math.*, 28:297–326, 1973.
- [4] S. S. Miller and P. T. Mocanu. Second-order differential inequalities in the complex plane. *J. Math. Anal. Appl.*, 65(2):289–305, 1978.
- [5] C. Pommerenke. *Univalent functions*. Vandenhoeck & Ruprecht, Göttingen, 1975.

Received February 8, 2005; revised November 14, 2005.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES,
FACULTY OF SCIENCE AND ARTS,
İSTANBUL KÜLTÜR UNIVERSITY,
İSTANBUL, TURKEY

E-mail address: y.polatoglu@iku.edu.tr

E-mail address: m.bolcal@iku.edu.tr

E-mail address: a.sen@iku.edu.tr

E-mail address: e.yavuz@iku.edu.tr