

CHARACTERIZATION OF FINITE GROUPS BY THEIR COMMUTING GRAPH

A. IRANMANESH AND A. JAFARZADEH

ABSTRACT. The commuting graph of a group G , denoted by $\Gamma(G)$, is a simple graph whose vertices are all non-central elements of G and two distinct vertices x, y are adjacent if $xy = yx$. In [1] it is conjectured that if M is a simple group and G is a group satisfying $\Gamma(G) \cong \Gamma(M)$, then $G \cong M$. In this paper we prove this conjecture for many simple groups.

1. INTRODUCTION

We denote by $\pi(n)$ the set of all prime divisors of n and if G is a finite group, then $\pi(G)$ is defined to be $\pi(|G|)$.

In this paper we consider simple graphs which are undirected, with no loops or multiple edges. The following definitions are standard and you can find them for example in [10].

For any graph Γ , we denote the set of vertices of Γ by $V(\Gamma)$. The *degree* $d_\Gamma(v)$ of a vertex v in Γ , is the number of edges incident to v and if the graph is understood, then we denote $d_\Gamma(v)$ simply by $d(v)$. A graph is called *regular* if the degrees of its vertices are the same. Two distinct vertices in Γ are called to be *adjacent*, if they are joined by an edge in Γ . A *path* P is a sequence of distinct vertices $v_0v_1 \dots v_k$ such that for all i ($0 \leq i \leq k-1$), v_i and v_{i+1} are adjacent vertices. A graph Γ is a *connected* graph, if there is path between each distinct pair of its vertices; otherwise Γ is *disconnected*. A maximal connected subgraph of a graph Γ is called a *component* of Γ . The complement G' of a simple graph G is a simple graph with the same vertex set as G , two vertices being adjacent in G' if and only if they are not adjacent in G .

We construct the commuting graph, the non-commuting graph and the prime graph of G as follows:

The *commuting graph* of G , denoted by $\Gamma(G)$, is a graph whose vertex set is $G \setminus Z(G)$, and two distinct vertices x and y are adjacent whenever $xy = yx$ and

2000 *Mathematics Subject Classification.* 20D05, 20D60, 05C25.

Key words and phrases. Simple group, Commuting graph, prime graph, order components.

the *non-commuting graph* of a group G , denoted by $\nabla(G)$, is the complement of $\Gamma(G)$, i.e. the graph with $G \setminus Z(G)$ as its vertex set and two vertices x and y are adjacent, if $xy \neq yx$ (see [1, 32]).

Finally, the *prime graph* of G , denoted by $\Gamma_1(G)$, is a graph whose vertex set is $\pi(G)$, and two distinct vertices p and q are adjacent if and only if G contains an element of order pq (see [30, 34]).

Denote the number of components of the prime graph of a group G with $t(G)$, and let $\pi_1, \pi_2, \dots, \pi_{t(G)}$ be the vertex set of the components of $\Gamma_1(G)$ and $T(G) = \{\pi_i(G) | i = 1, 2, \dots, t(G)\}$. If $2 \in \pi(G)$, then we always suppose $2 \in \pi_1$. Therefore,

$$\pi(G) = \bigcup_{i=1}^{t(G)} \pi_i.$$

Now, $|G|$ can be expressed as a product of coprime positive integers $m_i, i = 1, 2, \dots, t(G)$ where $\pi(m_i) = \pi_i$. These integers are called the *order components* of G and the set of order components of G is denoted by $OC(G)$:

$$OC(G) = \{m_i | i = 1, 2, \dots, t(G)\}.$$

If $|G|$ is even, then m_1 is called the *even order component* and $m_2, m_3, \dots, m_{t(G)}$ are called the *odd order components* of G .

In 1996, Chen posed the following question:

Question 1.1. Let M be a finite simple group. If G is a group such that $OC(G) = OC(M)$, do we have $G \cong M$?

Although, the answer to this question is “No” in general, a positive answer has been given for many groups. A simple group M is said to be *characterizable by its order components*, if $M \cong G$ for each group G such that $OC(G) = OC(M)$.

Remark 1.2. Suppose M is a finite group. In [1, 32], the authors conjectured that if G is a finite group such that $\nabla(M) \cong \nabla(G)$, then $|M| = |G|$. For every group H , $\nabla(H)$ and $\Gamma(H)$ are complement graphs, therefore for the groups H and K we have $\nabla(H) \cong \nabla(K)$ if and only if $\Gamma(H) \cong \Gamma(K)$. Hence the above conjecture is equivalent to say if G is a finite group such that $\Gamma(M) \cong \Gamma(G)$, then $|M| = |G|$. Although, they proved the statement for the groups S_n, A_n, D_{2n} , all sporadic simple groups and all simple groups of Lie type with disconnected prime graph, recently in [31], the author found some counterexamples to this conjecture. Here we state his counterexamples briefly:

For a prime p and an integer $r > 1$, there exists a non-abelian p -group P of order p^{2r} such that:

- (1) $|Z(P)| = p^r$;
- (2) $P/Z(P)$ is an elementary abelian p -group;
- (3) for every non-central element x of P , $C_P(x) = Z(P)\langle x \rangle$;
- (4) the non-commuting graph of P is regular.

According to this statement, there exists a 2-group P of order 2^{10} such that $P/Z(P)$ is an elementary abelian 2-group of order 2^5 and $\nabla(P)$ is a regular graph. Let A be an abelian group such that $|A| = a$ and consider the product $G = P \times A$. Then $\nabla(G)$ is a $(2^5)a(30)$ -regular graph, i.e., a graph with $d(v) = (2^5)a(30)$ for every vertex v of $\nabla(G)$ and has $(2^5)a(31)$ vertices. With a similar discussion, there exists a 5-group Q of order 5^6 such that $Q/Z(Q)$ is an elementary abelian 5-group of order 5^3 and $\nabla(Q)$ is a regular graph and if B is any abelian group such that $|B| = b$ and $H = Q \times B$, then $\nabla(H)$ is a $4b(5^3)(30)$ -regular graph with $4b(5^3)(31)$ vertices. Now, we must choose A and B so that $a2^3 = b5^3$. If we do that, both non-commuting graphs $\nabla(G)$ and $\nabla(H)$ are $(2^5)a(30)$ -regular graphs with the same number of vertices, and in fact they are isomorphic. However, the corresponding groups G and H have different orders.

In [1], the authors put forward another conjecture for the non-commuting graph of a group G . We rewrite this conjecture for the commuting graph of groups as follows:

Conjecture 1.3. Let M be a finite simple group. If G is any finite group such that $\Gamma(M) \cong \Gamma(G)$, then we have $M \cong G$.

In this paper, we will find the relation between the commuting graph and the prime graph of finite groups and then give a positive answer to Conjecture 1.3 for the groups pointed in Remark 1.2, using their characterization by their prime graph. Note that this conjecture is not true if we suppose M is an arbitrary finite group. In particular, the dihedral group and quaternion group of order 8 are not isomorphic while $\Gamma(D_8) \cong \Gamma(Q_8)$.

We will also prove the following theorem that gives a special characterization for all finite non-abelian simple groups:

Theorem 1.4. *Let G and M be two finite simple non-abelian groups. If $\Gamma(G) \cong \Gamma(M)$, then $G \cong M$.*

For a group G , let $N(G) = \{n \mid G \text{ has a conjugacy class of size } n\}$.

Lemma 1.5. *Let G_1 and G_2 be finite groups satisfying $|G_1| = |G_2|$ and $N(G_1) = N(G_2)$. Then $t(G_1) = t(G_2)$ and $OC(G_1) = OC(G_2)$.*

This is an immediate consequence of Lemma 1.5 in [9]:

Lemma 1.6. *Let G_1 and G_2 be finite groups satisfying $|G_1| = |G_2|$ and $N(G_1) = N(G_2)$. Then $t(G_1) = t(G_2)$ and $T(G_1) = T(G_2)$.*

The following basic theorem makes a relation between commuting graph of groups and their order components:

Theorem 1.7. *Let G_1 and G_2 be finite groups such that $|G_1| = |G_2|$ and $\Gamma(G_1) \cong \Gamma(G_2)$. Then $N(G_1) = N(G_2)$ and $OC(G_1) = OC(G_2)$.*

Proof. Set $|G_1| = |G_2| = n$. Since $\Gamma(G_1) \cong \Gamma(G_2)$, we have $|G_1 \setminus Z(G_1)| = |G_2 \setminus Z(G_2)|$ and therefore $|Z(G_1)| = |Z(G_2)|$. Also there exists a bijection $\varphi : V(\Gamma(G_1)) \rightarrow V(\Gamma(G_2))$ such that for all vertices $a, b \in V(\Gamma(G_1))$, a and b are adjacent if and only if $\varphi(a)$ and $\varphi(b)$ are adjacent. Set $|Z(G_1)| = |Z(G_2)| = z$ and suppose $1 \neq k \in N(G_1)$. Thus, there exists an element $x \in G_1$ with the conjugacy class $cl_{G_1}(x)$ in G_1 including x of size k . Therefore, $|C_{G_1}(x)| = n/k$ and thus $d_{\Gamma(G_1)}(x) = n/k - z - 1$. Obviously we have $d_{\Gamma(G_2)}(\varphi(x)) = n/k - z - 1$ and thus $|C_{G_2}(\varphi(x))| = n/k$. Therefore, $|cl_{G_2}(\varphi(x))| = k$ and thus, $k \in N(G_2)$. Hence, we have $N(G_1) \subseteq N(G_2)$ and with a similar reason we have $N(G_2) \subseteq N(G_1)$ and therefore the first statement is proved. The second statement follows immediately from Lemma 1.5. $\square$

2. CHARACTERIZATION OF SOME FINITE GROUPS BY THEIR COMMUTING GRAPH

First, we present the proof of Theorem 1.4:

Proof of Theorem 1.4. Since, $\Gamma(G) \cong \Gamma(M)$, they must have the same number of vertices, so $|G \setminus Z(G)| = |M \setminus Z(M)|$. On the other hand, $|Z(G)| = |Z(M)| = 1$, therefore $|G| = |M|$. Now, it is known that the only pairs of simple groups of the same order are

$$(A_8, PSL(3, 4)) \text{ and } (O(2n + 1, q) = B_n(q), PSp(2n, q) = C_n(q)),$$

where $n \geq 3$ and q is odd (see [29] and [33]). Thus, if $G \not\cong M$, then $G \cong A_8$ and $M \cong PSL(3, 4)$ or $G \cong B_n(q)$ and $M \cong C_n(q)$. For the first case, the element $a = (1\ 2)(3\ 4) \in A_8$ is included in a conjugacy class of length 210 and since $|A_8| = 20160$, we have $d_{\Gamma(A_8)}(a) = |C_{A_8}(a)| - 2 = 94$, while $N(PSL(3, 4)) = \{1, 315, 1260, 2240, 2880, 4032\}$ and thus $94 \notin \{d_{\Gamma(PSL(3, 4))}(x) | x \in PSL(3, 4) \setminus \{1\}\}$. Therefore, we have $\Gamma(A_8) \not\cong \Gamma(PSL(3, 4))$, because they have different degrees. For the second case, it has been proved that $N(B_n(q)) \neq N(C_n(q))$ (see [3]). Therefore, $\Gamma(B_n(q))$ and $\Gamma(C_n(q))$ have different sets of degrees and hence $\Gamma(B_n(q)) \not\cong \Gamma(C_n(q))$. Therefore, we must have $G \cong M$. $\square$

Now, we characterize some groups by their commuting graph:

Corollary 2.1. *Let $M = PSL(p, q)$, where p is a prime number and q is a prime power. If G is any group such that $\Gamma(M) \cong \Gamma(G)$, then $M \cong G$.*

Proof. Using Remark 1.2 and since $t(M) \geq 2$, we get $|M| = |G|$. Then by Theorem 1.7, we have $OC(M) = OC(G)$. Finally, by [4, 11, 12, 22, 23] we get the result. $\square$

Corollary 2.2. *Let $M = PSU(p, q)$ where p is an odd prime and q is a prime power. If G is any group such that $\Gamma(M) \cong \Gamma(G)$, then $M \cong G$.*

Proof. Similar to the proof of Corollary 2.1 and since $t(M) \geq 2$, we have $OC(M) = OC(G)$. Thus, by [13, 18, 19, 20, 28], we get the result. $\square$

Now, we prove that the groups $B_n(q)$ and $C_n(q)$ where $n = 2^m \geq 2$, are characterized by their commuting graphs, although we know that they cannot be characterized by their order components (see [24]):

Corollary 2.3. *Let M be a simple group of type $B_n(q)$ or $C_n(q)$ where $n = 2^m \geq 4$ or $n = 2$ and $q > 5$. If G is any group such that $\Gamma(M) \cong \Gamma(G)$, then $M \cong G$.*

Proof. Suppose $M \cong B_n(q)$ where $n = 2^m \geq 4$ or $n = 2$ and $q > 5$ and suppose $\Gamma(M) \cong \Gamma(G)$. Similar to the proof of Corollary 2.1 and since $t(M) = 2$, we have $OC(M) = OC(G)$. Thus, by [24], if q is even, then $G \cong M$, and if q is odd, then we have $G \cong M$ or $G \cong C_n(q)$. But, if q is odd and $G \cong C_n(q)$, then $\Gamma(C_n(q)) \cong \Gamma(G) \cong \Gamma(B_n(q))$ and this is a contradiction by Theorem 1.4. Thus, $G \cong M$. The proof for the case $M \cong C_n(q)$ is the same. $\square$

Corollary 2.4. *Let M be a simple group of one of the following types:*

- (a): $E_6(q)$ or $E_8(q)$;
- (b): $F_4(q)$ where $q > 2$;
- (c): ${}^2D_n(q)$ where $n = 2^m \geq 4$;
- (d): ${}^2D_p(3)$ where $p = 2^n + 1 \geq 5$;
- (e): ${}^2E_6(q)$ where $q > 2$;
- (f): ${}^3D_4(q)$;
- (g): A Suzuki–Ree group, i.e. a group of type ${}^2B_2(q)$, ${}^2F_4(q)$ or ${}^2G_2(q)$;
- (h): A sporadic simple group.

If G is any group such that $\Gamma(M) \cong \Gamma(G)$, then $M \cong G$.

Proof. Similar to the proof of Corollary 2.1 and since $t(M) \geq 2$, we have $OC(M) = OC(G)$. Thus, by [2, 26, 16, 17, 25, 14, 27, 7, 21, 6, 5], we get the result. $\square$

Professor J. G. Thompson has conjectured that:

Conjecture 2.5. *If G is a finite group with $Z(G) = 1$ and M a finite non-abelian simple group such that $N(G) = N(M)$, then $M \cong G$.*

Lemma 2.6. *Suppose M is a finite non-abelian simple group with $t(M) \geq 2$ which satisfies Thompson’s Conjecture. If G is a group such that $\Gamma(M) \cong \Gamma(G)$, then $M \cong G$.*

Proof. Since $\Gamma(M) \cong \Gamma(G)$, we have $|M \setminus Z(M)| = |G \setminus Z(G)|$ and $Z(M) = 1$. On the other hand, $|M| = |G|$ by Remark 1.2. Hence, $Z(G) = 1$ and by Theorem 1.7 we have $N(G) = N(M)$. Therefore, $M \cong G$. $\square$

Therefore we have proved the following corollary:

Corollary 2.7. *Let M be a simple group of type $G_2(q)$ where $q > 2$. If G is any group such that $\Gamma(M) \cong \Gamma(G)$, then $M \cong G$.*

Proof. By [8], M satisfies Thompson’s Conjecture. Now, the result follows from Lemma 2.6. $\square$

REFERENCES

- [1] A. Abdollahi, S. Akbari and H. R. Maimani, Non-commuting graph of a group, *J. Algebra*, **298** (2006), 468–492.
- [2] G. Y. Chen, A new characterization of $E_8(q)$, *J. Southwest China Normal Univ.*, **21** (1996), No. 3, 215–217.
- [3] G. Y. Chen, A new characterization of finite simple groups, *Chin. Sci. Bull.*, **40** (1995), No. 6, 446–450.
- [4] G. Y. Chen, A new characterization of $PSL(2, q)$, *Southeast Asian Bull. Math.*, **22** (1998), 257–263.
- [5] G. Y. Chen, A new characterization of sporadic simple groups, *Algebra Colloq.*, **3** (1996), No. 1, 49–58.
- [6] G. Y. Chen, A new characterization of Suzuki–Ree groups, *Sci. China (Ser. A)*, **27** (1997), No. 5, 430–433.
- [7] G. Y. Chen, Characterization of ${}^3D_4(q)$, *Southeast Asian Bull. Math.*, **25** (2001), 389–401.
- [8] G. Y. Chen, Further reflections on thompson’s conjecture, *J. Algebra*, **218** (1999), 276–285.
- [9] G. Y. Chen, On thompson’s conjecture, *J. Algebra*, **185** (1996), 184–193.
- [10] R. Diestel, *Graph theory*, 3rd edition, Grad. Texts in Math., Vol. 173, Springer-Verlag, New York, 2005.
- [11] A. Iranmanesh and S. H. Alavi, A characterization of simple groups $PSL(5, q)$, *Bull. Austral. Math. Soc.*, **65** (2002), 211–222.
- [12] A. Iranmanesh, S. H. Alavi and B. Khosravi, A characterization of $PSL(3, q)$ where $q = 2^n$, *Acta Math. Sinica, English Ser.*, **18** (2002), No. 3, 463–472.
- [13] A. Iranmanesh, S. H. Alavi and B. Khosravi, A characterization of $PSU(3, q)$ where $q > 5$, *Southeast Asian Bull. Math.*, **26** (2002), No. 1, 33–44.
- [14] A. Iranmanesh and B. Khosravi, A characterization of ${}^2D_p(3)$ where $p = 2^n + 1 (n \geq 2)$, *Hadronic J. Supp.*, **18** (2003), No. 4, 465–478.
- [15] A. Iranmanesh and B. Khosravi, A characterization of $C_2(q)$ where $q > 5$, *Comment. Math. Univ. Carolinae*, **44** (2002), No. 1, 9–21.
- [16] A. Iranmanesh and B. Khosravi, A characterization of $F_4(q)$ where $q = 2^n (n > 1)$, *Far East J. Math. Sci.*, **2** (2000), No. 6, 853–859.
- [17] A. Iranmanesh and B. Khosravi, A characterization of $F_4(q)$ where q is an odd prime power, *London Math. Society Lecture notes series*, **304** (2001), No. 1, 277–283.
- [18] A. Iranmanesh and B. Khosravi, A characterization of $PSU(5, q)$, *Int. Math. J.*, **3** (2003), No. 2, 1329–141.
- [19] A. Iranmanesh and B. Khosravi, A characterization of $PSU(7, q)$, *Int. J. appl. Math.*, **15** (2004), No. 4, 329–340.
- [20] A. Iranmanesh and B. Khosravi, A characterization of $PSU(11, q)$, *Can. Math. Bull.*, **47** (2004), No. 4, 530–539.
- [21] A. Iranmanesh and B. Khosravi, Characterization of ${}^3D_4(q)$ where $q > 2$, *Far East J. Math. Sci.*, **4** (2002), No. 2, 251–271.
- [22] A. Iranmanesh, B. Khosravi and S. H. Alavi, A characterization of $PSL(3, q)$ where q is an odd prime power, *J. Pure Appl. Algebra*, **170** (2002), No. 2–3, 243–254.
- [23] A. Khosravi and B. Khosravi, A new characterization of $PSL(p, q)$, *Comm. Algebra*, **32** (2004), No. 6, 2325–2339.
- [24] A. Khosravi and B. Khosravi, r –Recognizability of $B_n(q)$ and $C_n(q)$ where $n = 2^m \geq 4$, *J. pure Appl. Algebra*, **199** (2005), 149–165.
- [25] B. Khosravi, A characterization of ${}^2D_n(q)$, *Int. J. Math. Game Theory and Algebra*, **13** (2003), No. 3, 253–265.

- [26] B. Khosravi and B. Khosravi, A characterization of $E_6(q)$, *Algebras, Groups and Geometries*, **19** (2002), No. 2, 225–243.
- [27] B. Khosravi and B. Khosravi, A characterization of ${}^2E_6(q)$, *Kumamoto J. Math.*, **16** (2003), 1–11.
- [28] B. Khosravi, B. Khosravi and B. Khosravi, A new characterization of $PSU(p, q)$, *Acta Math. Hungarica*, **107** (2005), No. 3, 235–252.
- [29] W. Kimmerle, R. Lyons, R. Sandling and D. N. Teague, Composition factors from the group rings and Artin’s theorem on orders of simple groups, *Proc. London Math. Soc.*, **60** (1990), No. 3, 89–122.
- [30] A. S. Kondrat’ev, Prime graph components of finite groups, *Math. USSR-sb*, **67** (1990), No. 1, 235–247.
- [31] A. R. Moghaddamfar, About noncommuting graphs, *Sib. Math. J.*, **47** (2006), No. 5, 911–914.
- [32] A. R. Moghaddamfar, W. J. Shi, W. Zhou and A. R. Zokayi, On the noncommuting graph associated with a finite group, *Sib. Math. J.*, **46** (2005), No. 2, 325–332.
- [33] W. J. Shi, on the orders of finite simple groups, *Chin. Sci. Bull.*, **38** (1993), No. 4, 296–298. (Chinese)
- [34] J. S. Williams, Prime graph components of finite groups, *J. Algebra*, **69** (1981), No. 2, 487–513.

Received 17 October, 2006.

DEPARTMENT OF MATHEMATICS,
 TARBIAT MODARES UNIVERSITY,
 P.O.Box: 14115-137, TEHRAN,
 IRAN
E-mail address: iranmana@modares.ac.ir