

ON A CLASS OF LIE p -ALGEBRAS

CAMELIA CIOBANU

ABSTRACT. In this paper we study the finite dimensional Lie p -algebras, $\mathcal{L}$ splitting on its abelian p -socle, the sum of its minimal abelian p -ideals. In addition, some properties of the Frattini p -subalgebra of $\mathcal{L}$ are pointed out.

1. INTRODUCTION

In this section, we recall some notions and properties necessary in the paper.

Definition 1.1. A Lie p -algebra is a Lie algebra $\mathcal{L}$ with a p -map $a \rightarrow a^p$, such that:

$$\begin{aligned}(\alpha x)^p &= \alpha^p x^p, \text{ for all } \alpha \in \mathbb{K}, x \in \mathcal{L}, \\ x(ady^p) &= x(ad y)^p, \text{ for all } x, y \in \mathcal{L}, \\ (x + y)^p &= x^p + y^p + \sum_{i=1}^{p-1} s_i(x, y) \text{ for all } x, y \in \mathcal{L},\end{aligned}$$

where $is_i(x, y)$ is the coefficient of X^{i-1} in the expansion of $x(ad(Xu + y))^{p-1}$.

A subalgebra (respectively, ideal) of $\mathcal{L}$ is p -subalgebra (respectively, p -ideal) if it is closed under the p -map.

The notions of *maximal p -subalgebra* respectively *maximal p -ideal* of $\mathcal{L}$ are defined as usual. The intersection of p -subalgebras (respectively p -ideals) is a p -subalgebra (respectively a p -ideal) of $\mathcal{L}$.

We denote by $\Phi_p(\mathcal{L})$ the p -subsubalgebra of $\mathcal{L}$ obtained by intersecting all maximal p -subalgebras of $\mathcal{L}$ and we call it *the Frattini p -subalgebra* of $\mathcal{L}$.

The largest p -ideal of $\mathcal{L}$ included into $\Phi_p(\mathcal{L})$ is called *the Frattini p -ideal* and is denoted by $\mathcal{F}_p(\mathcal{L})$.

2000 *Mathematics Subject Classification.* 17B60, 17B66, 17B20.

Key words and phrases. Finite dimensional Lie p -algebra, p -ideal, p -subalgebra.

This paper was supported by the GAR 12/2007 - Contract 103/2008.

These are the corresponding notions to the Frattini subalgebra $\Phi(\mathcal{L})$ and the Frattini ideal $\mathcal{F}(\mathcal{L})$ for a Lie algebra.

We shall use the following notations:

- $[x, y]$ is the product of x, y in $\mathcal{L}$;
- $\mathcal{L}^{(1)}$ the derived algebra of $\mathcal{L}$;
- $\mathcal{L}^{(n)} = (\mathcal{L}^{(n-1)})^{(1)}$, for all $n \geq 2$;
- $(\mathcal{A})$ is the subalgebra generated by the subset $\mathcal{A}$ of $\mathcal{L}$;
- $(\mathcal{A})_p = (\{x^{p^n} | x \in (\mathcal{A}), p \in \mathbb{N}\})$, where $x^{p^n} = x^{(p^n)}$;
- $\mathcal{A}^p = (\{x^p | x \in \mathcal{A}\})$, where $\mathcal{A}$ is a subalgebra of $\mathcal{L}$;
- $\mathcal{A}^{p^n} = (\mathcal{A}^{p^{n-1}})^p$;
- $\mathcal{L}_1 = \bigcap_{i=1}^{\infty} \mathcal{L}^{p^i}$;
- $\mathcal{L}_0 = \{x \in \mathcal{L} | x^{p^n} = 0 \text{ for some } n\}$;
- $Z(\mathcal{L})$ is the center of $\mathcal{L}$;
- $\mathcal{N}(\mathcal{L})$ is the nilradical of $\mathcal{L}$.

Note that, if $\mathcal{L}$ is a p -algebra (finite dimensional), then $Z(\mathcal{L})$ is closed as p -ideal of $\mathcal{L}$.

2. LIE p -ALGEBRAS WHICH ARE $\mathcal{F}_p$ -FREE

In [8], Stitzinger has proved the following

Proposition 2.1. *If $\mathcal{L}$ is a finite dimensional Lie algebra over a field $\mathbb{K}$, then*

$$\mathcal{L}^{(1)} \cap Z(\mathcal{L}) \subseteq \mathcal{F}(\mathcal{L}).$$

We may prove an analogue of this proposition for a Lie p -algebra.

Lemma 2.2. *If $\mathcal{L}$ is a finite dimensional Lie p -algebra over a field $\mathbb{K}$, then we have*

$$(\mathcal{L}^{(1)})_p \cap Z(\mathcal{L}) \subseteq \mathcal{F}_p(\mathcal{L}).$$

Proof. Let $\mathcal{M}$ be a maximal p -subalgebra of $\mathcal{L}$ and suppose that $Z(\mathcal{L}) \not\subseteq \mathcal{M}$. Then $\mathcal{L} = \mathcal{M} + Z(\mathcal{L})$, so $\mathcal{L}^{(1)} = \mathcal{M}^{(1)} \subseteq \mathcal{M}$ and hence

$$(\mathcal{L}^{(1)})_p \subseteq (\mathcal{M})_p \subseteq \mathcal{M}.$$

□

The *abelian socle* $\text{Sa}(\mathcal{L})$ is the sum of all minimal ideals of $\mathcal{L}$.

We may define the *abelian p -socle* of the finite dimensional Lie p -algebra $\mathcal{L}$ as being the sum of all minimal abelian p -ideals of $\mathcal{L}$ and we denote it by $\text{Sap}(\mathcal{L})$.

The abelian socle (respectively, the abelian p -socle) of a finite dimensional Lie (p)-algebra is an ideal (a p -ideal) of $\mathcal{L}$, as one can show easily.

Definition 2.3. Let $\mathcal{L}$ be a finite dimensional Lie p -algebra and I be a p -ideal of $\mathcal{L}$. We say that $\mathcal{L}$ p -splits over I if there exists a p -subalgebra B of $\mathcal{L}$ such that $\mathcal{L} = I + B$.

B is called a p -complement of the p -ideal I .

Theorem 2.4. Let $\mathcal{L}$ be a finite dimensional Lie p -algebra such that $\mathcal{L}^{(1)} \neq 0$ and $\mathcal{L}^{(1)}$ is nilpotent. Then the following statements are equivalent:

- (i) $\mathcal{F}_p(\mathcal{L}) = 0$.
- (ii) $\text{Sap}(\mathcal{L}) = \mathcal{N}(\mathcal{L})$, and $\mathcal{L}$ p -splits over $\mathcal{N}(\mathcal{L})$.
- (iii) $\mathcal{L}^{(1)}$ is abelian, $(\mathcal{L}^{(1)})^p = 0$, $\mathcal{L}$ p -splits over $\mathcal{L}^{(1)} \oplus Z(\mathcal{L})$, and

$$\text{Sap}(\mathcal{L}) = \mathcal{L}^{(1)} \oplus Z(\mathcal{L}).$$

In the same hypotheses, the Cartan subalgebra of $\mathcal{L}$ are exactly those subalgebras which have $\mathcal{L}^{(1)}$ as a p -complement.

Proof. (i) $\Rightarrow$ (ii): These implications are immediate from Theorems 4.1, 4.2 of [5].

(iii) $\Rightarrow$ (i): This also follows from Theorem 4.1 of [5].

(i) $\Rightarrow$ (iii): Suppose that $\mathcal{F}_p(\mathcal{L}) = 0$. Then $\mathcal{F}(\mathcal{L}) = 0$, and $\mathcal{L}^{(1)}$ is abelian. Now $(\mathcal{L}^{(1)})^p \subseteq Z(\mathcal{L})$ by Lemma 2.1 [6], and so

$$(\mathcal{L}^{(1)})^p \subseteq (\mathcal{L}^{(1)})^p \cap Z(\mathcal{L}) \subseteq (\mathcal{L}^{(1)})^p \cap Z(\mathcal{L}) \subseteq \mathcal{F}_p(\mathcal{L}) = 0,$$

by Lemma 2.2. Clearly $\mathcal{L}^{(1)} \oplus Z(\mathcal{L}) \subseteq \mathcal{N}(\mathcal{L}) = \text{Sap}(\mathcal{L})$.

Now let $\underline{m}$ be a minimal (and hence abelian) p -ideal of $\mathcal{L}$. Then $[\mathcal{L}, \underline{m}] = \underline{m}$ is an ideal of $\mathcal{L}$ and

$$[\mathcal{L}, \underline{m}]^p \subseteq (\mathcal{L}^{(1)})^p \cap \underline{m}^p \subseteq (\mathcal{L}^{(1)})^p \cap Z(\mathcal{L}) = 0$$

by Lemma 2.1 of [6] and by Lemma 2.2. Hence $[\mathcal{L}, \underline{m}]$ is p -closed, therefore $[\mathcal{L}, \underline{m}] = \underline{m}$ or $[\mathcal{L}, \underline{m}] = 0$.

The former implies that $\underline{m} \subseteq \mathcal{L}^{(1)}$, and the latter that $\underline{m} \subseteq Z(\mathcal{L})$ hence $\text{Sap}(\mathcal{L}) = \mathcal{L}^{(1)} \oplus Z(\mathcal{L})$ and (iii) follows.

The last part of the theorem precises that the Cartan subalgebras are exactly those subalgebras having $\mathcal{L}^{(1)}$ as a p -complement. This follows from Proposition 1 of [8], or from Theorem 4.4.1.1. of [10] and from the fact that Cartan subalgebras are p -closed. $\square$

Corollary 2.5. If $\mathcal{L}$ is a finite dimensional Lie p -algebra over $\mathbb{K}$ with $\mathcal{L}^{(1)}$ nilpotent, and nonzero $\mathcal{F}_p(\mathcal{L}) = 0$ and $\mathbb{K}$ is perfect, then the maximal toral subalgebras are precisely those having as p -complement $\mathcal{L}^{(1)} \oplus Z(\mathcal{L})$.

Proof. Take $\mathcal{L} = (\mathcal{L}^{(1)} \oplus Z(\mathcal{L})) + B$, with B p -closed and $B^{(1)} = 0$ and let $B = B_0 \oplus B_1$ be the Fitting decomposition of B relatively to the p -map. Then $\mathcal{L}^{(1)} \oplus Z(\mathcal{L}) = \text{Sap}(\mathcal{L}) = \mathcal{N}(\mathcal{L})$ from Theorem 2.4, (ii), (iii). But $\mathcal{L}^{(1)} \oplus Z(\mathcal{L}) + B_0$

is a nilpotent ideal of $\mathcal{L}$ and so $B_0 \subseteq \mathcal{N}(\mathcal{L}) \cap B = 0$. Hence $B = B_1$ is toral. It is clear that $B_1 + Z(\mathcal{L})_1$ is a maximal toral subalgebra of $\mathcal{L}$.

Finally, let T be any maximal torus of $\mathcal{L}$, and let $\mathcal{C} = Z_{\mathcal{L}}(T)$. Then $\mathcal{C}$ is a Cartan subalgebra of $\mathcal{L}$, (by Theorem 4.5.17 of [10]) and $\mathcal{L} = \mathcal{L}^{(1)} + \mathcal{C}$ as above. Clearly we can write $\mathcal{C} = \mathcal{C}_0 \oplus T$. But now $\mathcal{L}^{(1)} + \mathcal{C}_0$ is a nilpotent ideal of $\mathcal{L}$, and so $\mathcal{C}_0 \subseteq \mathcal{N}(\mathcal{L}) \cap \mathcal{C} = Z(\mathcal{L})$, making T a p -complement of $\mathcal{L}^{(1)} \oplus Z(\mathcal{L})_0$. $\square$

The condition “ $\text{Sap}(\mathcal{L}) = \mathcal{L}^{(1)} \oplus Z(\mathcal{L})$ ” in (iii) Theorem 2.4. cannot be weakened to “ $Z(\mathcal{L}) \subseteq \text{Sap}(\mathcal{L})$ ”, as the following example proves.

Example 1. We know which are the Lie algebras of dimension 2 over $\mathbb{K}$ and we take $\mathcal{L} = I + V$, where

$$I = \mathbb{K}a + \mathbb{K}b, \quad V = \mathbb{K}v_1 + \mathbb{K}v_2,$$

$$v_1^p = v_2^p = b^p = 0, \quad a^p = 0,$$

$$[V, V] = 0, \quad [a, b] = 0, \quad [a, v_1] = v_1, \quad [a, v_2] = v_2, \quad [b, v_1] = v_2, \quad [b, v_2] = 0.$$

Then $\mathcal{L}^{(1)} = V$ is abelian, $(\mathcal{L}^{(1)})^p = 0$, $Z(\mathcal{L}) = 0$. Now

$$\mathcal{N}(\mathcal{L}) = \mathbb{K}b + \mathbb{K}v_1 + \mathbb{K}v_2.$$

Also $\mathbb{K}v_2$ is a maximal p -ideal. Let J be a minimal p -ideal contained in $\mathcal{N}(\mathcal{L})$. Since $[\mathcal{N}(\mathcal{L}), \mathcal{N}(\mathcal{L})] = \mathbb{K}v_2$, either $J = \mathbb{K}v_2$ or $[\mathcal{N}(\mathcal{L}), J] = 0$. Suppose that $J \neq \mathbb{K}v_2$. Then $[b, J] = 0$ so $J \subseteq \mathbb{K}b + \mathbb{K}v_2$, and $[v_1, J] = 0$ so $J \subseteq \mathbb{K}v_1 + \mathbb{K}v_2$. Thus $J \subseteq \mathbb{K}v_2$, a contradiction. Hence $\mathcal{N}(\mathcal{L}) \neq \text{Sap}(\mathcal{L})$.

E. L. Stitzinger has shown that, for any Lie algebra $\mathcal{L}$ over the arbitrary field $\mathbb{K}$, such that $\mathcal{L}^{(1)}$ is nilpotent, $\mathcal{L}$ is $\mathcal{F}$ -free (that is $\mathcal{F}(\mathcal{L}) = 0$) if and only if each subalgebra of $\mathcal{L}$ is $\mathcal{F}$ -free.

The complete analogue of this result does not hold if $\mathcal{F}(\mathcal{L})$ is replaced by $\mathcal{F}_p(\mathcal{L})$, as the following example proves.

Example 2. Let $\mathcal{L} = \mathbb{K}a + \mathbb{K}b + \mathbb{K}v_1 + \mathbb{K}v_2$ with $\mathbb{K} = Z_2$,

$$a^2 = a, b^2 = a + b, [a, v_1] = v_1, [a, v_2] = v_2, [b, v_1] = v_2, [b, v_2] = v_1 + v_2,$$

$$[a, b] = [v_1, v_2] = 0, v_1^2 = v_2^2 = 0,$$

and $I = \mathbb{K}a + \mathbb{K}b$. We get $\mathcal{F}_p(\mathcal{L}) = 0$ where as $\mathcal{F}_p(I) = \mathbb{K}a$.

However some partial results can be obtained.

Theorem 2.6. *Let $\mathcal{L}$ be a finite-dimensional p -Lie algebra. Then the following statements are equivalent:*

(i) $\mathcal{L}^{(1)}$ is nilpotent and $\mathcal{F}_p(\mathcal{L}) = 0$.

(ii) $\mathcal{L} = I + B$ where B is an abelian subalgebra, I is an abelian p -ideal, the (adjoint) action of B on I is faithful and completely reducible, $Z(\mathcal{L})$ is completely reducible under the p -map, and the p -map is trivial on $[B, I]$.

Proof. (i) $\Rightarrow$ (ii) By Theorem 2.4, $\mathcal{L} = I + B$, where

$$I = \text{Sap}(\mathcal{L}) = I_1 \oplus \cdots \oplus I_n,$$

with I_i is a minimal p -ideal of $\mathcal{L}$, for $i = 1, 2, \dots, n$, and B is a p -subalgebra of $\mathcal{L}$. Now $Z_B(I) = \{x \in B \mid [x, B] = 0\}$ is an ideal in the solvable Lie algebra $\mathcal{L}$. If $Z_B(I) \neq 0$, it follows that

$$0 \neq Z_B(I) \cap \text{Sap}(\mathcal{L}) \subseteq B \cap I = 0,$$

which is a contradiction. Hence $Z_B(I) = 0$ and the action of B is faithful.

Now suppose that $I_i \not\subseteq Z(\mathcal{L})$. Then $I_i \cap Z(\mathcal{L}) \subset I_i$ and so, as $I_i \cap Z(\mathcal{L})$ is a p -ideal, $I_i \cap Z(\mathcal{L}) = 0$. If $a \in I_i$ then $(ada)^p = 0$, and so $ada^p = 0$, hence $a^p \in Z(\mathcal{L})$. Thus, $a^p \in I_i \cap Z(\mathcal{L}) = 0$, and the minimality of I_i implies that I_i is an irreducible B -module but, of course, if $I_i \subseteq Z(\mathcal{L})$ then I_i is a completely reducible B -module, so $I = I_1 \oplus \cdots \oplus I_n$ is a completely reducible B -module.

Now $\mathcal{L}^{(1)}$ is nilpotent, therefore $\text{ad } x$ is nilpotent, for every $x \in B^{(1)}$. It follows from Engel's Theorem that $[B^{(1)}, I_i] \subset I_i$ for every $i = 1, 2, \dots, n$. If $I_i \not\subseteq Z(\mathcal{L})$, this implies that $[B^{(1)}, I_i] = 0$, since I_i is an irreducible B -module. If $I_i \subseteq Z(\mathcal{L})$ then, clearly, $[B^{(1)}, I_i] = 0$ also. Thus $[B^{(1)}, I_i] = 0$, and so $B^{(1)} = 0$, as $Z_B(I) = 0$. Moreover, $Z(\mathcal{L}) \subseteq I$, since $Z_B(I) = 0$. If $a \in Z(\mathcal{L})$ and $a = a_1 + \cdots + a_n$, with $a_i \in I_i$, then $[x, a_1] + \cdots + [x, a_n] = 0$, for all $x \in \mathcal{L}$, so each $a_i \in Z(\mathcal{L})$. Hence $Z(\mathcal{L}) = \sum I_i$, where the sum is over all I_i contained in $Z(\mathcal{L})$. Since each $I_i \subseteq Z(\mathcal{L})$ is a minimal p -ideal, $Z(\mathcal{L})$ must be irreducible under the p -map.

(ii) $\Rightarrow$ (i). In view of Theorem 4.1. of [5], it suffices to show that $I = \text{Sap}(\mathcal{L})$. Now we have $I = [B, I] \oplus Z(\mathcal{L})$, $[B, I]$ is a direct sum of irreducible B -modules (each of which is a minimal p -ideal) and $Z(\mathcal{L})$ is a direct sum of irreducible subspaces for the p -map (each of which is a minimal p -ideal). Thus, $I \subseteq \text{Sap}(\mathcal{L})$. But, as B acts faithfully on $\mathcal{L}$, I is a maximal abelian ideal. Hence $I = \text{Sap } \mathcal{L}$, as required. $\square$

Corollary 2.7. *Let $\mathcal{L}$ be a finite dimensional Lie p -algebra with $\mathcal{L}^{(1)}$ nilpotent and $\mathcal{F}_p(\mathcal{L}) = 0$. Let P be a p -subalgebra of $\mathcal{L}$ containing $\text{Sap}(\mathcal{L})$. Then $\mathcal{F}_p(P) = 0$.*

Proof. Write $\mathcal{L} = I + B$ as in Theorem 2.4 (ii). Then $P = I + (B \cap P)$ since $I = \text{Sap}(\mathcal{L}) \subseteq P$. Now B acts completely reducibly on $[B, I]$, and hence so does $B \cap P$. It follows that $B \cap P$ acts completely reducibly on $[B \cap P, I]$. Moreover, $Z(P) = Z(\mathcal{L}) \oplus Z_{[B, I]}(B \cap P)$ and the p -map is trivial on $[B, I]$, so that $Z(P)$ is completely reducible under the p -map. The result now follows from Theorem 2.4. $\square$

Corollary 2.8. *Let $\mathcal{L}$ be a finite dimensional Lie p -algebra such that $\mathcal{L}^{(1)}$ is nilpotent and $\mathcal{F}_p(\mathcal{L}) = 0$. If J is an ideal of $\mathcal{L}$, then $\text{Sap}(J) = 0$.*

Proof. It suffices to show this for maximal ideals. By Corollary 2.5, we may assume that $I_1 \not\subseteq J$, where $\text{Sap}(\mathcal{L}) = I_1 \oplus \cdots \oplus I_n$, with $I_1, \dots, I_n$ minimal abelian p -ideals. Then $\mathcal{L} = J + I_1$, since J is maximal, and $J \cap I_1 = 0$. Thus $\mathcal{L} = J \oplus I_1$, $J \cong \mathcal{L}/I_1 \cong B + (I_2 \oplus \cdots \oplus I_n)$, and $I_1 \subseteq Z(\mathcal{L})$. Hence $Z_B(I_2 \oplus \cdots \oplus I_n) = Z_B(I) = 0$, and it is clear that all of the conditions of Theorem 2.4 (ii) hold. $\square$

Corollary 2.9. *If $\mathcal{L}$ is an abelian finite dimensional Lie p -algebra, then $\mathcal{F}_p(\mathcal{L}) = 0$, if and only if $\mathcal{L}$ is completely reducible under the p -map.*

Proof. This statement can be proved by using Theorem 2.4 and the fact $B = 0$ and $\mathcal{L} = Z(\mathcal{L})$. $\square$

Corollary 2.10. *Let $\mathcal{L}$ be a finite dimensional Lie p -algebra such that $\mathcal{L} = \text{Sap}(\mathcal{L}) + B$ and that the conditions of Theorem 2.4 (ii) are satisfied. Assume in addition that B is completely reducible under the p -map; that is $\text{Sap}(B) = B$. Then if P is any p -subalgebra of $\mathcal{L}$, $P = \text{Sap } P + B'$, the conditions of Theorem 2.4. (ii) are satisfied and B' is completely reducible under the p -map.*

Proof. If $\text{Sap}(\mathcal{L}) \subseteq P$, then $\text{Sap}(P) = \text{Sap}(\mathcal{L})$, and taking $B' = B \cap P$, we get the result.

It suffices to prove the Corollary for maximal p -subalgebras. So assume that P is maximal and that $I_1 \not\subseteq P$, where $\text{Sap}(\mathcal{L}) = I_1 \oplus \cdots \oplus I_n$, with $I_1, \dots, I_n$ minimal abelian p -ideals. Then $\mathcal{L} = I_1 + P$, with $P \cap I_1 = 0$. Hence $P \cong B + (I_2 \oplus \cdots \oplus I_n)$. As B is completely reducible under the p -map, we have

$$B = B' \oplus Z_B(I_2 \oplus \cdots \oplus I_n).$$

Then $\text{Sap}(P) = Z_B(I_2 \oplus \cdots \oplus I_n) \oplus I_2 \oplus \cdots \oplus I_n$, $P = \text{Sap}(P) + B'$, the conditions of Theorem 2.4 (ii) are satisfied and B' is completely reducible under the p -map. $\square$

Definition 2.11. A finite dimensional Lie p -algebra $\mathcal{L}$ is called *p -elementary*, if $\mathcal{F}_p(P) = 0$ for every p -subalgebra P of $\mathcal{L}$.

Corollary 2.12. *Assume $\mathcal{L}^{(1)}$ is a finite dimensional Lie p -algebra with nilpotent $\mathcal{L}^{(1)}$ and $\mathcal{F}_p(\mathcal{L}) = 0$. Let $\mathcal{L} = \text{Sap}(\mathcal{L}) + B$ as in Theorem 2.4 (ii). Then $\mathcal{L}$ is p -elementary, if and only if $B = \text{Sap}(B)$*

Proof. As $\mathcal{F}_p(\mathcal{L}) = 0$ and $\mathcal{L} = \text{Sap}(\mathcal{L}) + B$ (Theorem 2.4. (ii)), then B has a faithful completely reducible representation on $\text{Sap}(\mathcal{L})$. This is equivalent to the fact that B has a non-zero nilideals as in [7]. Since B is abelian, this is equivalent to the injectivity of the p -map. Since $\mathbb{K}$ is algebraically closed, this is equivalent to $\text{Sap}(B) = B$ as in [4]. It follows from Corollary 2.14 that $\mathcal{L}$ is p -elementary. The converse is immediate from the definition. $\square$

The result above cannot be extended to the case when $\mathbb{K}$ is a perfect field. Let us see the following example.

Example 3. Let $\mathcal{L}$ be any abelian Lie p -algebra for which the p -map is injective but $\mathcal{L}$ is not completely reducible under the p -map. Then $\mathcal{L}$ has a faithful completely reducible module B . Make B into an abelian Lie p -algebra with trivial p -map. Then $\mathcal{F}_p(B + \mathcal{L}) = 0$, but $\mathcal{F}_p(\mathcal{L}) \neq 0$.

Now, if $\mathbb{K}$ is not perfect, let $\lambda \in \mathbb{K} \setminus \mathbb{K}^p$ and take $\mathcal{L} = \mathbb{K}a + \mathbb{K}b$, with $a^p = a$, $b^p = \lambda a$. If $\lambda \in \mathbb{K}$ and $\mu^p - \mu + \lambda = 0$ has no solution in $\mathbb{K}$, take $\mathcal{L} = \mathbb{K}a + \mathbb{K}b$ with $a^p = a$, $b^p = b + \lambda a$. Here we may take B to be p -dimensional with a represented by the identity matrix and b represented by the matrix

$$\begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \\ -\lambda & 1 & 0 & \dots & 0 \end{pmatrix}$$

(the companion matrix of $\mu^p - \mu + \lambda$). If $\mathbb{K} = \mathbb{Z}_p$ we may take $\lambda = -1$.

Putting $p = 2$, we get the example 2.7.

REFERENCES

- [1] D. W. Barnes. Lie algebras. Lecture Notes, University of Tübingen.
- [2] C. Ciobanu. *Cohomological study of Lie p -algebras*. PhD thesis, Ovidius University, Constantza, 2001.
- [3] C. Ciobanu and M. Stefanescu. *Introducere in studiul algebrelor Lie*. Mircea cel Batran, Naval Academy Publishing House, Constanta, 2000.
- [4] N. Jacobson. *Lie algebras*. Interscience Tracts in Pure and Applied Mathematics, No. 10. Interscience Publishers (a division of John Wiley & Sons), New York-London, 1962.
- [5] M. Lincoln and D. Towers. Frattini theory for restricted Lie algebras. *Arch. Math. (Basel)*, 45(5):451–457, 1985.
- [6] M. Lincoln and D. Towers. The Frattini p -subalgebra of a solvable Lie p -algebra. *Proc. Edinburgh Math. Soc. (2)*, 40(1):31–40, 1997.
- [7] J. R. Schue. Cartan decompositions for Lie algebras of prime characteristic. *J. Algebra*, 11:25–52, 1969.
- [8] E. L. Stitzinger. Frattini subalgebras of a class of solvable Lie algebras. *Pacific J. Math.*, 34:177–182, 1970.
- [9] D. A. Towers. A Frattini theory for algebras. *Proc. London Math. Soc. (3)*, 27:440–462, 1973.
- [10] D. J. Winter. *Abstract Lie algebras*. The M.I.T. Press, Cambridge, Mass.-London, 1972.

“MIRCEA CEL BATRAN” NAVAL ACADEMY,
1, FULGERULUI SREET, 900218, CONSTANTZA,
ROMANIA
E-mail address: cciobanu@anmb.ro