

UNIFORM CONVEXITY OF KÖTHE–BOCHNER FUNCTION SPACES

M. KHANDAQJI, F. AWAWDEH

ABSTRACT. Of concern are the Köthe–Bochner function spaces $E(X)$, where X is a real Banach space. Thus, of concern is the uniform convexity on the Köthe–Bochner function space $E(X)$. We show that $E(X)$ is uniformly convex if and only if both spaces E and X are uniformly convex. It is the uniform convexity that is the focal point.

1. BASIC CONCEPTS

Throughout this paper $(T, \sum, \mu)$ denote a δ -finite complete measure space and $L^\circ = L^\circ(T)$ denotes the space of all (equivalence classes) of $\sum$ -measurable real valued functions. For $f, g \in L^\circ$, $f \leq g$ means that $f(t) \leq g(t)$ μ -almost every where $t \in T$.

A Banach space is said to be a Köthe space if:

- (1) For any $f, g \in L^\circ$, $|f| \leq |g|$, $g \in E$ imply $f \in E$ and $\|f\|_E \leq \|g\|_E$.
- (2) For each $A \in \sum$, if $\mu(A)$ is finite then $\chi_A \in E$. See [12, p. 28].

Let E be a Köthe space on the measure space $(T, \sum, \mu)$ and $(X, \|\cdot\|_X)$ be a real Banach space. Then $E(X)$ is the space (of all equivalence classes of) strongly measurable functions $f : T \rightarrow X$ such that $\|f(\cdot)\|_X \in E$ equipped with the norm

$$\|f\| = \|\|f(\cdot)\|_X\|_E.$$

The space $(E(X), \|\cdot\|)$ is a Banach space called the Köthe–Bochner function space [11, p. 147]. The most important class of Köthe–Bochner function spaces $E(X)$ are the Lebesgue–Bochner spaces $L^p(X)$, $(1 \leq p < \infty)$ and their generalization the Orlicz–Bochner spaces $L^\phi(X)$. They have been studied by many authors [1], [2], [6], [7]. The geometric properties of the Köthe–Bochner function spaces have been studied by many authors, (e.g. [1], [5], [10]).

2000 *Mathematics Subject Classification.* 46E40, 35E10.

Key words and phrases. Köthe–Bochner function space, uniform convexity.

A Banach space is uniformly convex if and only if for every $\epsilon > 0$, there is a unique $\delta > 0$, such that for all x, y in X , the conditions $\|x\| = \|y\| = 1$ and $\|x - y\| > \epsilon$ imply $\|\frac{x+y}{2}\| < 1 - \delta$. Moreover this definition is equivalent to the following, [3, p. 127], for every pair of sequences (x_n) and (y_n) in X with $\|x_n\| \leq 1, \|y_n\| \leq 1$ and $\|x_n + y_n\| \leq 2$, it follows $\|x_n - y_n\| \rightarrow 0$.

For a Banach space X , we denote by δ_X the modulus of convexity

$$\delta_X(\epsilon) = \inf \left\{ 1 - \frac{1}{2} \|x + y\| : x, y \in X, \|x\| = \|y\| = 1, \|x - y\| > \epsilon \right\}$$

for any $\epsilon \in [0, 2]$. Note that X is uniformly convex if and only if $\delta_X(\epsilon) > 0$ whenever $\epsilon > 0$. If X is uniformly convex, we define the characteristic of convexity of by

$$\epsilon_r(X) = \sup \{ \epsilon \in [0, 2] : \delta_X(\epsilon) \leq r \}.$$

The uniform convexity of both the Lebesgue–Bochner spaces $L^p(X)$, $(1 \leq p < \infty)$ and the Orlicz–Bochner spaces $L^\phi(X)$ have been studied by many authors (e.g. see [4], [5], [8], [9], [14]).

In [14], M. Smith and B. Truett showed that many properties akin to uniform convexity lift from X the Lebesgue–Bochner spaces $L^p(X)$, $(1 \leq p < \infty)$. A survey of rotundity notions in the Lebesgue–Bochner spaces $L^p(X)$, $(1 \leq p < \infty)$ and sequences spaces can be found in [13].

In [10], A. Kaminska and B. Truett showed that many properties akin to uniform convexity lift from X to $E(X)$. The approach used in their paper is different than that we used.

This paper is devoted to the study of uniform convexity of the Köthe–Bochner function space $E(X)$, where X is a real Banach space.

2. MAIN RESULTS

Let us prove some preliminary results which will allow us in Theorem 4 to obtain a characterization of the uniform convexity of the Köthe–Bochner function space $E(X)$.

Lemma 1. *If (f_n) and (g_n) are sequences in the Köthe–Bochner function space $E(X)$ with $\|f_n\| = \|g_n\| = 1$ and $\|f_n + g_n\| \rightarrow 2$ then*

$$\|f_n(\cdot)\|_X + \|g_n(\cdot)\|_X \rightarrow 2.$$

Proof. Using the following inequalities we get the required result

$$\begin{aligned} \|f_n + g_n\| &= \|f_n(\cdot) + g_n(\cdot)\|_X \\ &\leq \|f_n(\cdot)\|_X + \|g_n(\cdot)\|_X \\ &\leq \|f_n\| + \|g_n\| = 2. \end{aligned}$$

□

Lemma 2. *Let X be a real Banach space and E be a Köthe space. If $(E, \|\cdot\|_E)$ is uniformly convex Köthe space and $(f_n), (g_n)$ are sequences in the Köthe-Bochner function space $E(X)$ with $\|f_n\| = \|g_n\| = 1$ and $\|f_n + g_n\| \rightarrow 2$ then,*

- (i) $\|f_n(\cdot)\|_X - \|g_n(\cdot)\|_X\|_E \rightarrow 0.$
- (ii) $\|f_n(\cdot)\|_X + \|g_n(\cdot)\|_X - \|f_n(\cdot) + g_n(\cdot)\|_X\|_E \rightarrow 0.$

Proof. Note that $(\|f_n(\cdot)\|_X)$ and $(\|g_n(\cdot)\|_X)$ are sequences in E with

$$\|f_n(\cdot)\|_X\|_E = \|g_n(\cdot)\|_X\|_E = 1.$$

Using Lemma 1 and the fact that E is uniformly convex Köthe space, it is straightforward to get (i). To prove (ii) we note first the inequalities

$$\begin{aligned} 2\|f_n + g_n\| &= 2\|f_n(\cdot) + g_n(\cdot)\|_X\|_E \\ &\leq \|f_n(\cdot)\|_X + \|g_n(\cdot)\|_X + \|f_n(\cdot) + g_n(\cdot)\|_X\|_E \\ &\leq 2\|f_n(\cdot)\|_X + \|g_n(\cdot)\|_X\|_E \\ &\leq 2(\|f_n\| + \|g_n\|) = 4. \end{aligned}$$

In this line, we get

$$\left\| \frac{f_n(\cdot) + g_n(\cdot)}{2} \right\|_X + \frac{\|f_n(\cdot)\|_X}{2} + \frac{\|g_n(\cdot)\|_X}{2} \Big\|_E \rightarrow 2.$$

Due to the uniform convexity of E and the facts that $\left\| \frac{f_n(\cdot) + g_n(\cdot)}{2} \right\|_X\|_E \leq 1$ and $\left\| \frac{\|f_n(\cdot)\|_X}{2} + \frac{\|g_n(\cdot)\|_X}{2} \right\|_E \leq 1$, result (ii) is completely proved. $\square$

Lemma 3. *Let X be a uniformly convex Banach space and $E(X)$ be a Köthe-Bochner function space. If $f, g \in E(X)$ then*

$$\begin{aligned} \|f - g\| &= \epsilon_r(X)(\|f\| + \|g\|) + \frac{2}{\epsilon}(\|f(\cdot)\|_X - \|g(\cdot)\|_X\|_E + \\ &\quad \|f(\cdot)\|_X + \|g(\cdot)\|_X - \|f(\cdot) + g(\cdot)\|_X\|_E). \end{aligned}$$

Proof. Holding $t \in T$ fixed and letting $\epsilon > 0$, we have two inequalities

- (i) $|\|f(t)\|_X - \|g(t)\|_X| \leq \epsilon \max\{\|f(t)\|_X, \|g(t)\|_X\}.$
- (ii) $\|f(t)\|_X + \|g(t)\|_X - \|f(t) + g(t)\|_X \leq \epsilon \max\{\|f(t)\|_X, \|g(t)\|_X\}.$

We have three cases to be considered

Case 1. (i) and (ii) are true. Assuming that $\|f(t)\|_X \geq \|g(t)\|_X$ and letting $\tilde{f}(t) = \frac{f(t)}{\|f(t)\|_X}$, $\tilde{g}(t) = \frac{g(t)}{\|f(t)\|_X}$, we find that $\|\tilde{g}(t)\|_X \leq \|\tilde{f}(t)\|_X = 1$. Furthermore, it is straightforward to verify that

$$\begin{aligned} \|\tilde{g}(t) + \tilde{f}(t)\|_X &\geq \frac{\|f(t)\|_X + \|g(t)\|_X - \epsilon\|f(t)\|_X}{\|f(t)\|_X} \\ &= 1 + \frac{\|g(t)\|_X}{\|f(t)\|_X} - \epsilon \end{aligned}$$

$$\begin{aligned}
&\geq 1 + \frac{\|f(t)\|_X - \epsilon \|f(t)\|_X}{\|f(t)\|_X} - \epsilon \\
&= 2 - 2\epsilon
\end{aligned}$$

and, since X is uniformly convex and $\|f(t)\|_X, \|g(t)\|_X$ are elements of X for a fixed $t \in T$, find that

$$\left\| \tilde{f}(t) - \tilde{g}(t) \right\|_X \leq \epsilon_r(X).$$

Therefore we deduce that

$$(1) \quad \|f(t) - g(t)\|_X \leq \epsilon_r(X) \max \{ \|f(t)\|_X, \|g(t)\|_X \}.$$

Case 2. (i) is not true. We imply that if

$$|\|f(t)\|_X - \|g(t)\|_X| > \max \{ \|f(t)\|_X, \|g(t)\|_X \}.$$

And so

$$\begin{aligned}
(2) \quad &\|f(t) - g(t)\|_X \leq 2 \max \{ \|f(t)\|_X, \|g(t)\|_X \} \\
&< \frac{2}{\epsilon} |\|f(t)\|_X - \|g(t)\|_X|.
\end{aligned}$$

Case 3. (ii) is not true. We can get that

$$\begin{aligned}
(3) \quad &\|f(t) - g(t)\|_X \leq 2 \max \{ \|f(t)\|_X, \|g(t)\|_X \} \\
&< \frac{2}{\epsilon} (\|f(t)\|_X + \|g(t)\|_X - \|f(t) + g(t)\|_X).
\end{aligned}$$

Inequalities (1), (2), and (3) give the inequality

$$\begin{aligned}
\|f(t) - g(t)\|_X &\leq \epsilon_r(X)(\|f(t)\|_X + \|g(t)\|_X) + \frac{2}{\epsilon} (\|f(t)\|_X + \\
&\|g(t)\|_X + \|f(t)\|_X + \|g(t)\|_X - \|f(t) + g(t)\|_X)
\end{aligned}$$

and (ii) is completely proved. $\square$

In this line, we are able to introduce the following main theorem about the uniform convexity of the Köthe–Bochner function space $E(X)$.

Theorem 4. *Let X be a real Banach space and E be a Köthe space. Then $E(X)$ is a uniformly convex Köthe–Bochner space if and only if both X and E are uniformly convex.*

Proof. Suppose $E(X)$ is uniformly convex Köthe–Bochner function space. Since both spaces X and E are embedded isometrically into $E(X)$, and due to the fact that uniform convexity inherited by subspaces, we deduce that both spaces X and E are uniformly convex.

Conversely, suppose X and E are uniformly convex spaces. Let (f_n) and (g_n) be sequences in $E(X)$ with $\|f_n\| = \|g_n\| = 1$ and $\|f_n + g_n\| \rightarrow 2$, then by Lemma 2 it follows that

$$\| \|f_n(\cdot)\|_X - \|g_n(\cdot)\|_X \|_E \rightarrow 0$$

and

$$\| \|f_n(\cdot)\|_X + \|g_n(\cdot)\|_X - \|f_n(\cdot) + g_n(\cdot)\|_X \|_E \rightarrow 0.$$

Take $\epsilon^* > 0$ and choose $\delta > 0$ such that $\epsilon_r(X) \leq \frac{\epsilon^*}{4}$. Choose K sufficiently large so that for all $n > K$,

$$\| \|f_n(\cdot)\|_X + \|g_n(\cdot)\|_X \|_E + \| \|f_n(\cdot)\|_X + \|g_n(\cdot)\|_X - \|f_n(\cdot) + g_n(\cdot)\|_X \|_E < \frac{1}{4}\epsilon^*.$$

From Lemma 3 it follows that

$$\begin{aligned} \| \|f_n - g_n\| \| &= \epsilon_r(X)(\| \|f_n\| \| + \| \|g_n\| \|) + \frac{2}{\epsilon}(\| \|f_n(\cdot)\|_X - \| \|g_n(\cdot)\|_X \|_E + \\ &\quad \| \|f_n(\cdot)\|_X + \| \|g_n(\cdot)\|_X - \| \|f_n(\cdot) + g_n(\cdot)\|_X \|_E) \\ &\leq 2\epsilon_r(X) + \frac{2}{\epsilon}\left(\frac{1}{4}\epsilon\epsilon^*\right) \\ &\leq \frac{1}{2}\epsilon^* + \frac{1}{2}\epsilon^* = \epsilon^*, \quad \text{for all } n > K. \end{aligned}$$

Consequently $\| \|f_n - g_n\| \| \rightarrow 0$, which means that $E(X)$ is a uniformly convex Köthe-Bochner space, and thereby the theorem is completely proved. $\square$

REFERENCES

- [1] J. Cerdà, H. Hudzik, and M. Mastyło. Geometric properties of Köthe-Bochner spaces. *Math. Proc. Cambridge Philos. Soc.*, 120(3):521–533, 1996.
- [2] S. Chen and H. Hudzik. On some convexities of Orlicz and Orlicz-Bochner spaces. *Comment. Math. Univ. Carolin.*, 29(1):13–29, 1988.
- [3] J. Diestel. *Sequences and series in Banach spaces*, volume 92 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1984.
- [4] G. Emmanuele and A. Villani. Lifting of rotundity properties from E to $L^p(\mu, E)$. *Rocky Mountain J. Math.*, 17(3):617–627, 1987.
- [5] I. Halperin. Uniform convexity in function spaces. *Duke Math. J.*, 21:195–204, 1954.
- [6] H. Hudzik and T. R. Landes. Characteristic of convexity of Köthe function spaces. *Math. Ann.*, 294(1):117–124, 1992.
- [7] H. Hudzik and M. Mastyło. Strongly extreme points in Köthe-Bochner spaces. *Rocky Mountain J. Math.*, 23(3):899–909, 1993.
- [8] A. Kamińska and W. Kurc. Weak uniform rotundity in Orlicz spaces. *Comment. Math. Univ. Carolin.*, 27(4):651–664, 1986.
- [9] A. Kamińska and B. Turett. Uniformly non- $l^1(n)$ Orlicz-Bochner spaces. *Bull. Polish Acad. Sci. Math.*, 35(3-4):211–218, 1987.
- [10] A. Kamińska and B. Turett. Rotundity in Köthe spaces of vector-valued functions. *Canad. J. Math.*, 41(4):659–675, 1989.
- [11] P.-K. Lin. *Köthe-Bochner function spaces*. Birkhäuser Boston Inc., Boston, MA, 2004.
- [12] J. Lindenstrauss and L. Tzafriri. *Classical Banach spaces. II*, volume 97 of *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*. Springer-Verlag, Berlin, 1979. Function spaces.
- [13] M. A. Smith. Rotundity and extremity in $l^p(X_i)$ and $L^p(\mu, X)$. In *Geometry of normed linear spaces (Urbana-Champaign, Ill., 1983)*, volume 52 of *Contemp. Math.*, pages 143–162. Amer. Math. Soc., Providence, RI, 1986.
- [14] M. A. Smith and B. Turett. Rotundity in Lebesgue-Bochner function spaces. *Trans. Amer. Math. Soc.*, 257(1):105–118, 1980.

DEPARTMENT OF MATHEMATICS,
HASHEMITE UNIVERSITY,
JORDAN
E-mail address: mkhan@hu.edu.jo
E-mail address: awawdeh@hu.edu.jo