

MULTIPLY WARPED PRODUCT ON QUASI-EINSTEIN MANIFOLD WITH A SEMI-SYMMETRIC NON-METRIC CONNECTION

SAMPA PAHAN, BUDDHADEV PAL, AND ARINDAM BHATTACHARYYA

ABSTRACT. In this paper, we have studied warped products and multiply warped product on quasi-Einstein manifold with semi-symmetric non-metric connection. Then we have applied our results to generalized Robertson-Walker space times with a semi-symmetric non-metric connection.

1. INTRODUCTION

Let (M^n, g) , $(n > 2)$ be a Riemannian manifold and $U_S = \{x \in M : S \neq \frac{r}{n}g \text{ at } x\}$, then the manifold (M^n, g) is said to be quasi-Einstein manifold [4, 6] if on $U_S \subset M$, we have

$$(1.1) \quad S - \alpha g = \beta A \otimes A,$$

where A is a 1-form on U_S and α and β some functions on U_S . It is clear that the 1-form A as well as the function β are nonzero at every point on U_S . The scalars α, β are known as the associated scalars of the manifold. Also, the 1-form A is called the associated 1-form of the manifold defined by $g(X, \rho) = A(X)$ for any vector field X , ρ being a unit vector field, called the generator of the manifold. Such an n -dimensional quasi-Einstein manifold is denoted by $(QE)_n$.

Let (B, g_B) and (F, g_F) be two Riemannian manifolds and $f > 0$ is a differential function on B . Consider the product manifold $B \times F$ with its projections $\pi: B \times F \rightarrow B$ and $\sigma: B \times F \rightarrow F$. The warped product $B \times_f F$ is the manifold $B \times F$ with the Riemannian structure such that $\|X\|^2 = \|\pi^*(X)\|^2 + f^2(\pi(p))\|\sigma^*(X)\|^2$, for any vector field X on M . Thus we have $g_M = g_B + f^2 g_F$ holds on M . Here B is called the base of M and F the fiber. The function f is called the warping function of the warped product [9].

2010 *Mathematics Subject Classification.* 53C25, 53C15.

Key words and phrases. Quasi-Einstein manifold, warped product, multiply warped product, semi-symmetric non-metric connection.

The first author is supported by UGC JRF of India, Ref. No: 23/06/2013(i)EU-V..

The concept of warped products was first introduced by Bishop and O'Neil [3] to construct examples of Riemannian manifold with negative curvature.

Now, we can generalize warped products to multiply warped products. A multiply warped product is the product manifold $M = B \times_{b_1} F_1 \times_{b_2} F_2 \dots \times_{b_m} F_m$ with the metric $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \oplus b_3^2 g_{F_3} \dots \oplus b_m^2 g_{F_m}$, where each $i \in \{1, 2, \dots, m\}$, $b_i: B \rightarrow (0, \infty)$ is smooth and (F_i, g_{F_i}) is a pseudo-Riemannian manifold. In particular, when $B = (c, d)$, the metric $g_B = -dt^2$ is negative and (F_i, g_{F_i}) is a Riemannian manifold. We call M as the multiply generalized Robertson-Walker space-time.

A multiply twisted product (M, g) is a product manifold of the form $M = B \times_{b_1} F_1 \times_{b_2} F_2 \dots \times_{b_m} F_m$ with the metric $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \oplus b_3^2 g_{F_3} \dots \oplus b_m^2 g_{F_m}$, where each $i \in \{1, 2, \dots, m\}$, $b_i: B \times F_i \rightarrow (0, \infty)$ is smooth.

In 1924, Friedmann and Schouten was introduced the notion of a semi-symmetric linear connection on a differential manifold [5]. The idea of metric connection with torsion on Riemannian manifold has given by Hayden (1932) in [7]. In 1970, Yano [15] was introduced a systematic study of semi-symmetric metric connection on Riemannian manifold. Later K. S. Amur and S. S. Pujar [1], C. S. Bagewadi [2], Sharafuddin and Hussian (1976) [11], S. Sular, C. Özgür [12], M. Tripathi [13] have also studied semi-symmetric metric connection on Riemannian manifold. In [10], S. Sular and C. Özgür has studied warped product on semi-symmetric non-metric connection. Y. Wang has considered multiply warped product with a semi-symmetric non-metric connection, then applied the results to generalized Robertson-Walker space-time in [14].

In this paper, we have considered quasi-Einstein warped product manifolds endowed with semi-symmetric metric non-connection. First we have obtained the necessary and sufficient conditions of quasi-Einstein warped product manifold with semi-symmetric non-metric connection. Next we have established that under certain conditions Robertson-Walker space times would be converted to quasi-Einstein manifold with the above connection. Later we have shown that $(\bar{n}-1)$ -dimensional base is isometric to a $(\bar{n}-1)$ -dimensional sphere of a particular radius with respect to semi-symmetric non-metric connection. In the last section we have studied special multiply warped product with semi symmetric non-metric connection.

2. PRELIMINARIES

Let (M^n, g) be a Riemannian manifold with the Levi-civita connection ∇ . A linear connection $\tilde{\nabla}$ on (M^n, g) is said to be semi-symmetric if its torsion tensor T can be written as

$$T(X, Y) = \tilde{\nabla}_X Y - \tilde{\nabla}_Y X - [X, Y],$$

satisfies the condition

$$T(X, Y) = \pi(Y)X - \pi(X)Y,$$

where π is an 1-form on M^n with the associated vector field P defined by $\pi(X) = g(X, P)$, for all vector fields $X \in \chi(M^n)$.

A connection $\tilde{\nabla}$ is called semi-symmetric non-metric connection if $\tilde{\nabla}g \neq 0$.

The relation between semi-symmetric non-metric connection $\tilde{\nabla}$ and the Levi-Civita connection ∇ of M^n and it is given by [14]

$$(2.1) \quad \tilde{\nabla}_X Y = \nabla_X Y + \pi(Y)X,$$

where $g(X, P) = \pi(X)$.

Further, a relation between the curvature tensors R and $\tilde{R}$ of type (1,3) of the connections ∇ and $\tilde{\nabla}$ respectively is given by [14],

$$(2.2) \quad \begin{aligned} \tilde{R}(X, Y)Z &= R(X, Y)Z + g(Z, \nabla_X P)Y - g(Z, \nabla_Y P)X \\ &\quad + \pi(Z)[\pi(Y)X - \pi(X)Y], \end{aligned}$$

for any vector field X, Y, Z on M^n .

3. GENERALIZED ROBERTSON-WALKER SPACE-TIMES WITH A SEMI-SYMMETRIC NON-METRIC CONNECTION

In this section we have considered quasi-Einstein warped product manifolds with respect to semi-symmetric non-metric connection. Now, we have proved the following theorem.

Theorem 3.1. *Let (M, g) be a warped product $I \times_f F$ where I is an open interval in $\mathbb{R}$, $\dim I = 1$ and $\dim F = \bar{n} - 1$, ($\bar{n} \geq 3$). Then (M, g) is a quasi-Einstein manifold with respect to semi-symmetric non-metric connection iff F is quasi-Einstein manifold for $P \in \chi(B)$ with respect to the Levi-Civita connection or the warping function f is a constant on I for $P \in \chi(F)$.*

Proof. Assume that $P \in \chi(B)$ and let g_I be the metric on I . Taking $f = e^{\frac{q}{2}}$ and by using the proposition use of [10] we get

$$(3.1) \quad \tilde{S}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = -\frac{\bar{n}-1}{4}[2q'' + (q')^2 - 4]g_I\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right),$$

$$(3.2) \quad \tilde{S}\left(\frac{\partial}{\partial t}, V\right) = 0,$$

$$(3.3) \quad \tilde{S}(V, W) = S^F(V, W) + e^q\left[\frac{\bar{n}-1}{2}q' + \frac{q''}{2} - \frac{\bar{n}-3}{4}(q')^2\right]g_F(V, W),$$

for any vector field V, W on F .

Since, M is a quasi-Einstein manifold with respect to the semi-symmetric non-metric connection, we have

$$\tilde{S}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = \alpha g\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) + \beta \eta\left(\frac{\partial}{\partial t}\right)\eta\left(\frac{\partial}{\partial t}\right),$$

and

$$\tilde{S}(V, W) = \alpha g(V, W) + \beta \eta(V)\eta(W),$$

Then the last equations reduce to

$$(3.4) \quad \tilde{S}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = \alpha g_I\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) + \beta \eta\left(\frac{\partial}{\partial t}\right) \eta\left(\frac{\partial}{\partial t}\right),$$

and

$$(3.5) \quad \tilde{S}(V, W) = \alpha e^q g_F(V, W) + \beta \eta(V) \eta(W).$$

Decomposing the vector field U uniquely into its components U_I and U_F on I and F , respectively, then we have $U = U_I + U_F$. Since $\dim I = 1$, we can take $U_I = v \frac{\partial}{\partial t}$ which gives $U = v \frac{\partial}{\partial t} + U_F$, where v is a function on M . Then we can write

$$(3.6) \quad \eta\left(\frac{\partial}{\partial t}\right) = g(U, \frac{\partial}{\partial t}) = v.$$

Using the equations (3.6), the equations (3.4), (3.5) reduce to

$$(3.7) \quad \tilde{S}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = \alpha + \beta v^2,$$

and

$$(3.8) \quad \tilde{S}(V, W) = \alpha e^q g_F(V, W) + \beta \eta(V) \eta(W).$$

Comparing the right hand sides of (3.1) and (3.7) we get,

$$(3.9) \quad \alpha + \beta v^2 = -\frac{\bar{n} - 1}{4} [2q'' + (q')^2 - 4].$$

Similarly comparing the right hand sides of (3.3) and (3.8) we obtain

$$(3.10) \quad S^F(V, W) = e^q \left[\alpha + \frac{\bar{n} - 3}{4} (q')^2 - \frac{(\bar{n} - 1)}{2} q' - \frac{q''}{2} \right] g_F(V, W) + \beta \eta(V) \eta(W),$$

which gives that F is a quasi-Einstein manifold with respect to the Levi-Civita connection for $P \in \chi(B)$.

Now taking $P \in \chi(F)$ and by use of [10] we get,

$$(3.11) \quad \tilde{S}\left(\frac{\partial}{\partial t}, V\right) = (\bar{n} - 1) \frac{q'}{2} \pi(V) g_I\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right)$$

and

$$(3.12) \quad \tilde{S}\left(V, \frac{\partial}{\partial t}\right) = (1 - \bar{n}) \frac{q'}{2} \pi(V) g_I\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right),$$

for any vector field $V \in \chi(F)$.

Since M is a quasi-Einstein manifold, we have

$$(3.13) \quad \tilde{S}\left(\frac{\partial}{\partial t}, V\right) = \tilde{S}\left(V, \frac{\partial}{\partial t}\right) = \alpha g\left(V, \frac{\partial}{\partial t}\right) + \beta \eta(V) \eta\left(\frac{\partial}{\partial t}\right).$$

Now $g(V, \frac{\partial}{\partial t}) = 0$ as $\frac{\partial}{\partial t} \in \chi(B)$ and $V \in \chi(F)$.

Hence from the last equation we get

$$(3.14) \quad \tilde{S}\left(\frac{\partial}{\partial t}, V\right) = \tilde{S}\left(V, \frac{\partial}{\partial t}\right) = \beta \eta(V) \eta\left(\frac{\partial}{\partial t}\right).$$

Therefore we have

$$(3.15) \quad \eta(V)\eta\left(\frac{\partial}{\partial t}\right) = (\bar{n} - 1)\frac{q'}{2}\pi(V)g_I\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right),$$

$$(3.16) \quad \eta(V)\eta\left(\frac{\partial}{\partial t}\right) = (1 - \bar{n})\frac{q'}{2}\pi(V)g_I\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right).$$

Comparing from (3.15), (3.16) we get

$$q' = 0.$$

Hence, q is constant. Therefore f is constant. $\square$

Now, we consider the warped product $M = B \times_f I$ with $\dim B = \bar{n} - 1$, $\dim I = 1$ ($\bar{n} \geq 3$). Under this assumption we have obtained the following theorem.

Theorem 3.2. *Let (M, g) be a warped product $B \times_f I$, where $\dim I = 1$ and $\dim B = \bar{n} - 1$ ($\bar{n} \geq 3$).*

- i) *If (M, g) is a quasi-Einstein manifold with scalars α, β respect to the semi-symmetric non-metric connection, $P \in \chi(B)$ is parallel on B with respect to the Levi-Civita connection on B and f is a constant on B , then $\alpha = 0$.*
- ii) *If (M, g) is a quasi-Einstein manifold with respect to the semi-symmetric non-metric connection for $P \in \chi(F)$, then f is a constant on B .*
- iii) *If f is a constant on B and B is a quasi-Einstein manifold with respect to the Levi-Civita connection for $P \in \chi(F)$, then M is an quasi-Einstein manifold with respect to the semi-symmetric non-metric connection.*

Proof. Assume that (M, g) is a quasi-Einstein manifold with respect to the semi-symmetric non-metric connection. Then we write

$$(3.17) \quad \tilde{S}(X, Y) = \alpha g(X, Y) + \beta \eta(X)\eta(Y).$$

Decomposing the vector field U uniquely into its components U_B and U_I on B and I , respectively, then we have

$$(3.18) \quad U = U_B + U_I.$$

Since $\dim I = 1$, we can take $U_I = v \frac{\partial}{\partial t}$ which gives $U = v \frac{\partial}{\partial t} + U_B$, where v is a function on M . From (3.17), (3.18) and from the proposition of [10], we have,

$$(3.19) \quad \tilde{S}^B(X, Y) = \alpha g_B(X, Y) + \beta g_B(X, U_B)g_B(Y, U_B) - \frac{H^f(X, Y)}{f} - g(Y, \nabla_X P) + \pi(X)\pi(Y).$$

By contraction over X and Y and we get

$$(3.20) \quad \tilde{r}^B = \alpha(\bar{n} - 1) + \beta g_B(U_B, U_B) - \frac{\Delta f}{f} + \pi(P) - \sum_{i=1}^{\bar{n}-1} g(e_i, \nabla_{e_i} P).$$

Also from (3.17) we have

$$(3.21) \quad \tilde{r}^M = \alpha \bar{n} + \beta g_B(U_B, U_B),$$

So, by the use of (3.21) in (3.20) we get

$$(3.22) \quad \tilde{r}^B = \tilde{r}^M - \alpha - \frac{\Delta f}{f} + \pi(P) - \sum_{i=1}^{\bar{n}-1} g(e_i, \nabla_{e_i} P)$$

Also from the proposition of [10] we get

$$\tilde{r}^M = \tilde{r}^B - 2 \frac{\Delta f}{f} - \pi(P) + \sum_{i=1}^{\bar{n}-1} g(e_i, \nabla_{e_i} P) + (\bar{n} - 1) \frac{Pf}{f}.$$

Therefore, from above two relation we get

$$\alpha + \frac{\Delta f}{f} - \pi(P) + \sum_{i=1}^{\bar{n}-1} g(e_i, \nabla_{e_i} P) = -2 \frac{\Delta f}{f} - \pi(P) + \sum_{i=1}^{\bar{n}-1} g(e_i, \nabla_{e_i} P) + (\bar{n} - 1) \frac{Pf}{f}.$$

Since $P \in \chi(B)$ is parallel and f is a constant on B , then we get $\alpha = 0$.

ii) Let $P \in \chi(F)$. By the use of the proposition of [10] we get,

$$\tilde{S}(X, V) = (\bar{n} - 1) \pi(V) \frac{Xf}{f}$$

and

$$\tilde{S}(V, X) = (1 - \bar{n}) \pi(V) \frac{Xf}{f},$$

for any vector field $X \in \chi(B)$ and $V \in \chi(F)$. Since $F = I$, then taking $V = P$ we have

$$(3.23) \quad \tilde{S}(X, P) = (\bar{n} - 1) \pi(P) \frac{Xf}{f},$$

and

$$(3.24) \quad \tilde{S}(P, X) = (1 - \bar{n}) \pi(P) \frac{Xf}{f}.$$

Since M is a quasi-Einstein manifold, we have

$$\tilde{S}(X, P) = \tilde{S}(P, X) = \alpha g(P, X) + \beta \eta(P) \eta(X).$$

Again we have $g(P, X) = 0$ for $X \in \chi(B)$ and $P \in \chi(F)$.

Hence, we have $Xf = 0$. This implies that f is constant.

iii) Assume that B is a quasi-Einstein manifold with respect to Levi-Civita connection. Then we have

$$(3.25) \quad S^B(X, Y) = \alpha g(X, Y) + \beta \eta(X) \eta(Y),$$

for any vector field X, Y tangent to B .

$$\tilde{S}^M(X, Y) = S^B(X, Y) + \frac{H^f(X, Y)}{f},$$

for any vector field $P \in \chi(F)$. Since f is a constant, then $H^f(X, Y) = 0$ for all $X, Y \in \chi(B)$.

The above equation reduces to

$$(3.26) \quad \tilde{S}^M(X, Y) = S^B(X, Y).$$

By the use of (3.25) in (3.26) we get

$$(3.27) \quad \tilde{S}^M(X, Y) = \alpha g(X, Y) + \beta \eta(X) \eta(Y).$$

which shows that M is a quasi-Einstein manifold with respect to the semi-symmetric non-metric connection. $\square$

Next, we consider generalized Robertson-Walker space time with a semi-symmetric non-metric connection. Now we prove the following theorem.

Theorem 3.3. *Let (M, g) be a warped product $I \times_f F$ with the metric tensor $-dt^2 + f(t)^2 g_F$, $P = \frac{\partial}{\partial t}$, $\dim F = l$. Then (M, g) is a quasi-Einstein manifold with respect to semi-symmetric non-metric connection $\tilde{\nabla}$ with constant associated scalars α and β , if and only if the following conditions are satisfied.*

- i) (F, g_F) is quasi-Einstein manifold with scalar α_F, β_F .
- ii) $l(1 - \frac{f''}{f}) = \alpha - v^2 \beta$,
- iii) $\alpha_F + (1 - l)f'^2 - \alpha f^2 + f''f + lf'f = 0$ and $\beta = \beta_F$.

Proof. By the proposition of [10] we have

$$\tilde{S}(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}) = l(\frac{f''}{f} - 1),$$

$$\tilde{S}(\frac{\partial}{\partial t}, V) = \tilde{S}(V, \frac{\partial}{\partial t}) = 0,$$

$$\tilde{S}(V, W) = S^F(V, W) + g_F(V, W)\{ff'' - (l - 1)f'^2 + lff'\}.$$

Then by the quasi-Einstein condition, we get the theorem 3.3. $\square$

From the theorem 3.3. Putting $\dim F = 1$ we get the following corollary.

Corollary 3.1. *Let (M, g) be a warped product $I \times_f F$ with the metric tensor $-dt^2 + f(t)^2 g_F$, $P = \frac{\partial}{\partial t}$, $\dim F = 1$. Then (M, g) is a quasi-Einstein manifold with respect to semi-symmetric non-metric connection if and only if*

$$f'' + (\alpha - v^2 \beta - 1)f = 0.$$

By the corollary 3.1. and elementary methods for ordinary differential equations we get

Theorem 3.4. *Let (M, g) be a warped product $I \times_f F$ with the metric tensor $-dt^2 + f(t)^2 g_F$, $P = \frac{\partial}{\partial t}$, $\dim F = 1$. Then (M, g) is a quasi-Einstein manifold with respect to semi-symmetric non-metric connection if and only if*

- i) when $\alpha - v^2 \beta < 1$, $f(t) = c_1 e^{(\sqrt{1 - (\alpha - v^2 \beta)})t} + c_2 e^{-(\sqrt{1 - (\alpha - v^2 \beta)})t}$,
- ii) when $\alpha - v^2 \beta = 1$, $f(t) = c_1 + c_2 t$,

- iii) when $\alpha - v^2\beta > 1$, we have that $f(t) = c_1 \cos((\sqrt{\alpha - v^2\beta - 1})t) + c_2 \sin((\sqrt{\alpha - v^2\beta - 1})t)$.

Next the following theorem shows when base of quasi-Einstein warped product manifold is isometric to a sphere of a particular radius.

Theorem 3.5. *Let (M, g) be a warped product $B \times_f I$ connected with $(\bar{n} - 1)$ -dimensional Riemannian manifold B where $\bar{n} \geq 3$ and one-dimensional Riemannian manifold I . If (M, g) is a quasi-Einstein manifold with constant associated scalars α and β , $U \in \chi(M)$ with respect to semi-symmetric non-metric connection, $P \in \chi(B)$ and the Hessian of f is proportional to metric tensor g_B , then (B, g_B) is a $(\bar{n} - 1)$ -dimensional sphere of radius $\rho = \frac{\bar{n}-1}{\sqrt{\tilde{r}^B + \alpha}}$.*

Proof. Let M be a warped product manifold. Then from the proposition of [10] we have

$$(3.28) \quad \tilde{S}^M(X, Y) = \tilde{S}^B(X, Y) + \left[\frac{H^f(X, Y)}{f} + g(\nabla_X P, Y) - \pi(X)\pi(Y) \right],$$

for any vector field X, Y on B . Since M is a quasi-Einstein manifold with respect to semi-symmetric non-metric connection, we have

$$(3.29) \quad \tilde{S}^M(X, Y) = \alpha g(X, Y) + \beta \eta(X)\eta(Y).$$

Decomposing the vector field U uniquely into its components U_B and U_I on B and I , respectively, then we have

$$(3.30) \quad U = U_B + U_I.$$

Since $\dim I = 1$, we can take $U_I = v \frac{\partial}{\partial t}$ which gives $U = v \frac{\partial}{\partial t} + U_B$, where v is a function on M . Putting the value of (3.29), (3.30) in (3.28) we get

$$(3.31) \quad \begin{aligned} \tilde{S}^B(X, Y) &= \alpha g_B(X, Y) + \beta g_B(X, U_B)g_B(Y, U_B) \\ &\quad - \left[\frac{H^f(X, Y)}{f} + g(\nabla_X P, Y) - \pi(X)\pi(Y) \right]. \end{aligned}$$

By contraction over X and Y we get,

$$(3.32) \quad \tilde{r}^B = \tilde{r}^M - \alpha - \frac{\Delta f}{f} + \pi(P) - \sum_{i=1}^{\bar{n}-1} g(e_i, \nabla_{e_i} P).$$

Again from the proposition of [10] we get

$$(3.33) \quad \frac{\tilde{r}^M}{\bar{n}} = (\bar{n} - 1) \frac{Pf}{f} - \frac{\Delta f}{f}.$$

From the last two equations we get

$$(3.34) \quad (\tilde{r}^B + \alpha)f = \bar{n}(\bar{n} - 1)Pf - (\bar{n} + 1)\Delta f + f\pi(P) - \sum_{i=1}^{\bar{n}-1} fg(e_i, \nabla_{e_i} P).$$

Hence we get

$$(3.35) \quad \frac{(\tilde{r}^B + \alpha)f}{\bar{n}(\bar{n} - 1)} = Pf - \frac{(\bar{n} + 1)\Delta f}{\bar{n}(\bar{n} - 1)} + \frac{f\pi(P)}{\bar{n}(\bar{n} - 1)} - \sum_{i=1}^{\bar{n}-1} \frac{fg(e_i, \nabla_{e_i}P)}{\bar{n}(\bar{n} - 1)}.$$

Since, the Hessian of f is proportional to metric tensor g_B , then we have

$$(3.36) \quad H^f(X, Y) = \frac{\bar{n}}{\bar{n} - 1} \left[-Pf + \frac{(\bar{n} + 1)\Delta f}{\bar{n}(\bar{n} - 1)} - \frac{f\pi(P)}{\bar{n}(\bar{n} - 1)} + \sum_{i=1}^{\bar{n}-1} \frac{fg(e_i, \nabla_{e_i}P)}{n(n - 1)} \right] g_B(X, Y).$$

Hence from the equations (3.35), (3.36) we get

$$(3.37) \quad H^f(X, Y) + \frac{\tilde{r}^B + \alpha}{(\bar{n} - 1)^2} fg_B(X, Y) = 0.$$

So, B is isometric to the $(\bar{n} - 1)$ -dimensional sphere of radius $\frac{\bar{n}-1}{\sqrt{\tilde{r}^B + \alpha}}$ [8]. Thus the theorem is proved. $\square$

4. SPECIAL MULTIPLY WARPED PRODUCT MANIFOLDS WITH SEMI-SYMMETRIC NON-METRIC CONNECTION

Let $M = B \times_{b_1} F_1 \times_{b_2} F_2 \dots \times_{b_m} F_m$ be a multiply warped product with the metric tensor $-dt^2 \oplus b_1^2 g_{F_1} \oplus \dots \oplus b_m^2 g_{F_m}$ and I is an open interval in $\mathbb{R}$ and $b_i \in C^\infty(I)$.

Now, we prove the following theorem for multiply generalized Robertson-Walker space time.

Theorem 4.1. *Let $M = I \times_{b_1} F_1 \times_{b_2} F_2 \dots \times_{b_m} F_m$ be a multiply warped product with the metric tensor $-dt^2 \oplus b_1^2 g_{F_1} \oplus \dots \oplus b_m^2 g_{F_m}$ and $P = \frac{\partial}{\partial t}$. Then (M, g) is a quasi-Einstein manifold with respect to semi-symmetric non-metric connection $\tilde{\nabla}$ with constant associated scalars α and β , if and only if the following conditions are satisfied.*

- i) (F_i, g_{F_i}) is quasi-Einstein manifold with scalar $\alpha_{F_i}, \beta_{F_i}, i \in \{1, 2, \dots, m\}$,
- ii) $\sum_{i=1}^m l_i (1 - \frac{b_i''}{b_i}) = \alpha - v^2 \beta$,
- iii) $\alpha b_i^2 - \alpha_{F_i} + b_i b_i'' + (l_i - 1) b_i'^2 + b_i b_i' \sum_{j \neq i} l_j \frac{b_j'}{b_j} - b_i^2 \sum_{j=1}^m l_j \frac{b_j'}{b_j} = 0$ and $\beta = \beta_{F_i}$.

Proof. By the proposition of [14] we have

$$(4.1) \quad \tilde{S}(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}) = - \sum_{i=1}^m l_i (1 - \frac{b_i''}{b_i}),$$

$$(4.2) \quad \tilde{S}(\frac{\partial}{\partial t}, V) = \tilde{S}(V, \frac{\partial}{\partial t}) = 0,$$

$$(4.3) \quad \tilde{S}(V, W) =$$

$$S^{F_i}(V, W) + g_{F_i}(V, W)\{-b_i b_i'' - (l_i - 1)b_i'^2 - b_i b_i' \sum_{j \neq i} l_j \frac{b_j'}{b_j} + b_i^2 \sum_{j=1}^m l_j \frac{b_j'}{b_j}\}.$$

Since, M is a quasi-Einstein manifold. So,

$$\tilde{S}(X, Y) = \alpha g(X, Y) + \beta \eta(X) \eta(Y).$$

Now,

$$\tilde{S}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = \alpha g\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) + \beta \eta\left(\frac{\partial}{\partial t}\right) \eta\left(\frac{\partial}{\partial t}\right).$$

Decomposing the vector field U uniquely into its components U_I and U_F on I and F , respectively, then we have $U = U_I + U_F$. Since $\dim I = 1$, we can take $U_I = v \frac{\partial}{\partial t}$ which gives $U = v \frac{\partial}{\partial t} + U_F$, where v is a function on M . Then we can write

$$(4.4) \quad \eta\left(\frac{\partial}{\partial t}\right) = g(U, \frac{\partial}{\partial t}) = v.$$

Hence, we get

$$\sum_{i=1}^m l_i \left(1 - \frac{b_i''}{b_i}\right) = \alpha - v^2 \beta.$$

Again, $\tilde{S}(V, W) = \alpha g(V, W) + \beta \eta(V) \eta(W)$.

From by the proposition of [14] and the equation (4.3) we get (F_i, g_{F_i}) is quasi-Einstein manifold.

Also, after some calculation we can show that

$$\alpha b_i^2 - \alpha_{F_i} + b_i b_i'' + (l_i - 1)b_i'^2 + b_i b_i' \sum_{j \neq i} l_j \frac{b_j'}{b_j} - b_i^2 \sum_{j=1}^m l_j \frac{b_j'}{b_j} = 0$$

and $\beta = \beta_{F_i}$. □

Next, we have obtained the following theorem with some condition of fibre and warping function with semi-symmetric non-metric connection.

Theorem 4.2. *Let $M = I \times_{b_1} F_1 \times_{b_2} F_2 \dots \times_{b_m} F_m$ be a multiply warped product with the metric tensor $-dt^2 \oplus b_1^2 g_{F_1} \oplus \dots \oplus b_m^2 g_{F_m}$ with $P \in \chi(F_r)$ and $g_{F_r}(P, P) = 1$ and $\bar{n} \geq 3$. Then (M, g) is a quasi-Einstein manifold with respect to semi-symmetric non-metric connection $\check{\nabla}$ with constant associated scalars α and β , if and only if the following conditions are satisfied.*

- i) (F_i, g_{F_i}) ($i \neq r$) is quasi-Einstein manifold with scalar $\alpha_{F_i}, \beta_{F_i}$, $i \in \{1, 2, \dots, m\}$.
- ii) $\sum_{i=1}^m l_i \frac{b_i''}{b_i} = -\alpha + v^2 \beta$.
- iii) $\alpha_{F_i} - b_i b_i'' - (l_i - 1)b_i'^2 - b_i b_i' \sum_{j \neq i} l_j \frac{b_j'}{b_j} - \alpha b_i^2 = 0$ and $\beta = \beta_{F_i}$.

iv)

$$S^{F_i}(V, W) - g_{F_i}(V, W)[b_i b_i'' + (l_i - 1)b_i'^2 + \alpha b_i^2 + b_i b_i' \sum_{j \neq i} l_j \frac{b_j'}{b_j'}] =$$

$$(\bar{n} - 1)[\pi(V)\pi(W) - \frac{g(W, \nabla_V P) + g(V, \nabla_W P)}{2}],$$

for $V, W \in \Gamma(TF_r)$, $r = i$.

Proof. By the proposition of [14] and $g_{F_r}(P, P) = 1$, we have that b_r is constant. So, we have

$$\tilde{S}(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}) = \sum_{i=1}^m l_i \frac{b_i''}{b_i} = -\alpha + v^2 \beta.$$

By variables separation, we have

$$\tilde{S}(V, W) = S^{F_i}(V, W) + b_i^2 g_{F_i}(V, W)[- \frac{b_i''}{b_i} - (l_i - 1) \frac{b_i'^2}{b_i^2} - \sum_{j \neq i} l_j \frac{b_i' b_j'}{b_i b_j}]$$

$$+ (\bar{n} - 1)[g(W, \nabla_V P) - \pi(V)\pi(W)].$$

When $i \neq r$, then $\pi(V) = \nabla_V P = \nabla_W P = 0$.

$$\tilde{S}(V, W) = S^{F_i}(V, W) + b_i^2 g_{F_i}(V, W)[- \frac{b_i''}{b_i} - (l_i - 1) \frac{b_i'^2}{b_i^2} - \sum_{j \neq i} l_j \frac{b_i' b_j'}{b_i b_j}]$$

$$= \alpha b_i^2 g_F(V, W) + \beta \eta(V) \eta(W).$$

By variables separation, we have (F_i, g_{F_i}) ($i \neq r$) is quasi-Einstein manifold with scalar $\alpha_{F_i}, \beta_{F_i}$, $i \in \{1, 2, \dots, m\}$.

When $i = r$, we get

$$S^{F_i}(V, W) - g_{F_i}(V, W)[b_i b_i'' + (l_i - 1)b_i'^2 + \alpha b_i^2 + b_i b_i' \sum_{j \neq i} l_j \frac{b_j'}{b_j'}] =$$

$$(\bar{n} - 1)[\pi(V)\pi(W) - \frac{g(W, \nabla_V P) + g(V, \nabla_W P)}{2}]. \quad \square$$

5. ACKNOWLEDGMENTS

The authors wish to express their sincere thanks and gratitude to the editors and referees.

REFERENCES

- [1] K. Amur and S. S. Pujar. On submanifolds of a Riemannian manifold admitting a metric semisymmetric connection. *Tensor (N.S.)*, 32(1):35–38, 1978.
- [2] C. S. Bagewadi. On totally real submanifolds of a Kählerian manifold admitting semisymmetric metric F -connection. *Indian J. Pure Appl. Math.*, 13(5):528–536, 1982.
- [3] R. L. Bishop and B. O'Neill. Manifolds of negative curvature. *Trans. Amer. Math. Soc.*, 145:1–49, 1969.

- [4] M. C. Chaki and R. K. Maity. On quasi Einstein manifolds. *Publ. Math. Debrecen*, 57(3-4):297–306, 2000.
- [5] A. Friedmann and J. A. Schouten. über die Geometrie der halbsymmetrischen übertragungen. *Math. Z.*, 21(1):211–223, 1924.
- [6] M. g. Gł ogowska. On quasi-Einstein Cartan type hypersurfaces. *J. Geom. Phys.*, 58(5):599–614, 2008.
- [7] H. A. Hayden. Sub-Spaces of a Space with Torsion. *Proc. London Math. Soc.*, S2-34(1):27.
- [8] M. Obata. Certain conditions for a Riemannian manifold to be isometric with a sphere. *J. Math. Soc. Japan*, 14:333–340, 1962.
- [9] B. O'Neill. *Semi-Riemannian geometry*, volume 103 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1983. With applications to relativity.
- [10] C. Özgür and S. Sular. Warped products with a semi-symmetric non-metric connection. *Arab. J. Sci. Eng.*, 36(3):461–473, 2011.
- [11] A. Sharfuddin and S. I. Husain. Semi-symmetric metric connexions in almost contact manifolds. *Tensor (N.S.)*, 30(2):133–139, 1976.
- [12] S. Sular and C. Özgür. Warped products with a semi-symmetric metric connection. *Taiwanese J. Math.*, 15(4):1701–1719, 2011.
- [13] M. M. Tripathi. A new connection in a Riemannian manifold. *Int. Electron. J. Geom.*, 1(1):15–24, 2008.
- [14] Y. Wang. Curvature of multiply warped products with an affine connection. *Bull. Korean Math. Soc.*, 50(5):1567–1586, 2013.
- [15] K. Yano. On semi-symmetric metric connection. *Rev. Roumaine Math. Pures Appl.*, 15:1579–1586, 1970.

Received February 4, 2016.

SAMPA PAHAN,
 DEPARTMENT OF MATHEMATICS,
 JADAVPUR UNIVERSITY,
 KOLKATA-700032,
 INDIA
E-mail address: sampapahan25@gmail.com

BUDDHADEV PAL,
 DEPARTMENT OF MATHEMATICS,
 INSTITUTE OF SCIENCE,
 BANARAS HINDU UNIVERSITY,
 VARANASI-221005,
 INDIA
E-mail address: pal.buddha@gmail.com

ARINDAM BHATTACHARYYA,
 DEPARTMENT OF MATHEMATICS,
 JADAVPUR UNIVERSITY,
 KOLKATA-700032,
 INDIA
E-mail address: bhattachar1968@yahoo.co.in