

t –BALANCING NUMBERS, PELL NUMBERS AND SQUARE TRIANGULAR NUMBERS

AHMET TEKCAN AND AZIZ YAZLA

ABSTRACT. Let $t \geq 2$ be an integer. In this work we get all integer solutions of the Diophantine equation $8r^2 + 8tr + 1 = y^2$ in order to determine the general terms of all t –balancing numbers for which $2t^2 - 1$ is prime. Later we obtain some formulas for the sums of Pell, Pell–Lucas, balancing and Lucas–balancing numbers in terms of t –balancing numbers and also we deduce the general terms of all t –balancing numbers in terms of square triangular numbers.

1. PRELIMINARIES.

A positive integer n is called a balancing number (see [1] and [3]) if the Diophantine equation

$$(1.1) \quad 1 + 2 + \cdots + (n - 1) = (n + 1) + (n + 2) + \cdots + (n + r)$$

holds for some positive integer r which is called cobalancing number (or balancer). If n is a balancing number with balancer r , then from (1.1) one has $\frac{(n-1)n}{2} = rn + \frac{r(r+1)}{2}$ and so

$$(1.2) \quad r = \frac{-(2n + 1) + \sqrt{8n^2 + 1}}{2} \quad \text{and} \quad n = \frac{2r + 1 + \sqrt{8r^2 + 8r + 1}}{2}.$$

Let B_n denote the n^{th} balancing number, and let b_n denote the n^{th} cobalancing number. Then $B_1 = 1, B_2 = 6, B_{n+1} = 6B_n - B_{n-1}$ and $b_1 = 0, b_2 = 2, b_{n+1} = 6b_n - b_{n-1} + 2$ for $n \geq 2$. The zeros of the characteristic equation $x^2 - 6x + 1 = 0$ for balancing numbers are $\alpha_1 = 3 + \sqrt{8}$ and $\beta_1 = 3 - \sqrt{8}$. Ray derived some nice results on balancing and cobalancing numbers in his Phd thesis (*Balancing and Cobalancing Numbers*, Department of Maths., National Institute of Technology, Rourkela, India, 2009). Since x is a balancing number if and only if $8x^2 + 1$ is a perfect square, he set $y^2 - 8x^2 = 1$ for some integer $y \neq 0$. The fundamental solution is $(y_1, x_1) = (3, 1)$. So $y_n + x_n\sqrt{8} = (3 + \sqrt{8})^n$

2010 *Mathematics Subject Classification.* 05A19, 11B37, 11B39.

Key words and phrases. Pell equation, balancing number, t –balancing number, square triangular number.

and similarly $y_n - x_n\sqrt{8} = (3 - \sqrt{8})^n$ for $n \geq 1$. Thus $x_n = \frac{(3+\sqrt{8})^n - (3-\sqrt{8})^n}{2\sqrt{8}}$ which is the Binet formula for balancing numbers and is denoted by B_n . Let $\alpha = 1 + \sqrt{2}$ and $\beta = 1 - \sqrt{2}$ be the roots of the characteristic equation for Pell (and also Pell–Lucas) numbers defined by $P_0 = 0, P_1 = 1, P_n = 2P_{n-1} + P_{n-2}$ (and $Q_0 = Q_1 = 2, Q_n = 2Q_{n-1} + Q_{n-2}$) for $n \geq 2$. Since $\alpha^2 = 3 + \sqrt{8}$ and $\beta^2 = 3 - \sqrt{8}$, the Binet formula for balancing numbers is $B_n = \frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}}$. Similarly $b_n = \frac{\alpha^{2n-1} - \beta^{2n-1}}{4\sqrt{2}} - \frac{1}{2}$.

From (1.2), we note that B_n is a balancing number if and only if $8B_n^2 + 1$ is a perfect square and b_n is a cobalancing number if and only if $8b_n^2 + 8b_n + 1$ is a perfect square. Thus

$$(1.3) \quad C_n = \sqrt{8B_n^2 + 1} \text{ and } c_n = \sqrt{8b_n^2 + 8b_n + 1}$$

are integers called the n^{th} Lucas–balancing and n^{th} Lucas–cobalancing number. Their Binet formulas are $C_n = \frac{\alpha^{2n} + \beta^{2n}}{2}$ and $c_n = \frac{\alpha^{2n-1} + \beta^{2n-1}}{2}$ (for further details see [7, 8, 9, 10]).

Balancing numbers and their generalizations have been investigated by several authors from many aspects (see [4, 5, 6, 12]). Recently in [2], Dash, Ota and Dash considered the t –balancing numbers for an integer $t \geq 2$. A positive integer n is called a t –balancing number if

$$(1.4) \quad 1 + 2 + \cdots + n = (n + 1 + t) + (n + 2 + t) + \cdots + (n + r + t)$$

holds for some positive integer r which is called t –cobalancing (or t –balancer) number. For example

- 2, 14, 84, 492, 2870, $\dots$ are 0–balancing numbers with 0–balancers 1, 6, 35, 204, 1189, $\dots$;
- 5, 34, 203, 1188, 6929, $\dots$ are 1–balancing numbers with 1–balancers 2, 14, 84, 492, 2870, $\dots$;
- 3, 8, 25, 54, 153, $\dots$ are 2–balancing numbers with 2–balancers 1, 3, 10, 22, 63, $\dots$;
- 6, 11, 45, 74, 272, $\dots$ are 3–balancing numbers with 3–balancers 2, 4, 18, 30, 112, $\dots$.

(Here we note that 0–and 1–balancing numbers can be given in terms of balancing numbers, indeed, $B_n^0 = b_{n+1}, b_n^0 = B_n, C_n^0 = c_{n+1}, c_n^0 = C_n$ and $B_n^1 = B_{n+1} - 1, b_n^1 = b_{n+1}, C_n^1 = C_{n+1}, c_n^1 = c_{n+1}$, that is why it is assumed that $t \geq 2$).

From (1.4) we see that

$$(1.5) \quad r = \frac{-(2n + 2t + 1) + \sqrt{8n^2 + 8n(1 + t) + (2t + 1)^2}}{2} \text{ and}$$

$$n = \frac{(2r - 1) + \sqrt{8r^2 + 8tr + 1}}{2}.$$

Let B_n^t denote the n^{th} t -balancing number and let b_n^t denote the n^{th} t -cobalancing number. Then from (1.5), we see that B_n^t is a t -balancing number if and only if $8(B_n^t)^2 + 8B_n^t(1+t) + (2t+1)^2$ is a perfect square and b_n^t is a t -cobalancing number if and only if $8(b_n^t)^2 + 8tb_n^t + 1$ is a perfect square. So

$$(1.6) \quad C_n^t = \sqrt{8(B_n^t)^2 + 8B_n^t(1+t) + (2t+1)^2} \quad \text{and} \quad c_n^t = \sqrt{8(b_n^t)^2 + 8tb_n^t + 1}$$

are integers which are called the n^{th} Lucas t -balancing and n^{th} Lucas t -cobalancing number.

2. RESULTS.

In the present paper, we want to determine the general terms of all t -balancing numbers. But we first determine the set of all positive integer solutions of the Diophantine equation

$$(2.1) \quad 8r^2 + 8tr + 1 = y^2.$$

Let us explain why? We note that for giving any t -cobalancing number r , $8r^2 + 8tr + 1$ is a perfect square. So we let $8r^2 + 8tr + 1 = y^2$ for some integer $y \neq 0$. Thus from (2.1), we deduce that $2(2r+t)^2 - y^2 = 2t^2 - 1$. So putting $x = 2r + t$, we get the Pell equation

$$(2.2) \quad 2x^2 - y^2 = 2t^2 - 1.$$

Now let Δ be a positive non-square discriminant and let $O_\Delta = \{x + y\rho_\Delta : x, y \in \mathbb{Z}\}$, where $\rho_\Delta = \sqrt{\frac{\Delta}{4}}$ if $\Delta \equiv 0 \pmod{4}$, or $\frac{1+\sqrt{\Delta}}{2}$ if $\Delta \equiv 1 \pmod{4}$. So O_Δ is a subring of $\mathbb{Q}(\sqrt{\Delta}) = \{x + y\sqrt{\Delta} : x, y \in \mathbb{Q}\}$. Then the unit group O_Δ^* is defined to be the group of units of the ring O_Δ . For the quadratic form $F(x, y) = ax^2 + bxy + cy^2$ of discriminant $\Delta = b^2 - 4ac$, we can write $F(x, y) = \frac{(xa+y\frac{b+\sqrt{\Delta}}{2})(xa+y\frac{b-\sqrt{\Delta}}{2})}{a}$. So the module M_F of F is the O_Δ -module $M_F = \{xa + y\frac{b+\sqrt{\Delta}}{2} : x, y \in \mathbb{Z}\} \subset \mathbb{Q}(\sqrt{\Delta})$. Therefore we get $(u + v\rho_\Delta)(xa + y\frac{b+\sqrt{\Delta}}{2}) = x'a + y'\frac{b+\sqrt{\Delta}}{2}$, where

$$(2.3) \quad [x' \ y'] = \begin{cases} [x \ y] \begin{bmatrix} u - \frac{b}{2}v & av \\ -cv & u + \frac{b}{2}v \end{bmatrix} & \text{if } \Delta \equiv 0 \pmod{4} \\ [x \ y] \begin{bmatrix} u + \frac{1-b}{2}v & av \\ -cv & u + \frac{1+b}{2}v \end{bmatrix} & \text{if } \Delta \equiv 1 \pmod{4}. \end{cases}$$

So there is a bijection $\Psi : \{(x, y) : F(x, y) = m\} \rightarrow \{\gamma \in M_F : N(\gamma) = am\}$ for solving the Diophantine equation $F(x, y) = m$, that is, $ax^2 + bxy + cy^2 = m$. The action of $O_{\Delta,1}^* = \{\alpha \in O_\Delta^* : N(\alpha) = 1\}$ on the set $\Omega = \{(x, y) : F(x, y) = m\}$ of integral solutions of the equation $F(x, y) = m$ is most interesting when Δ is a positive non-square since $O_{\Delta,1}^*$ is infinite. Therefore the orbit of each solution will be infinite and so the set Ω is either empty or infinite. Since $O_{\Delta,1}^*$ can be explicitly determined, the set Ω is satisfactorily described by the representation of such a list, called a set of representatives of the orbits.

Let ε_Δ be the smallest unit of O_Δ that is greater than 1 and let $\tau_\Delta = \varepsilon_\Delta$ if $N(\varepsilon_\Delta) = 1$; or ε_Δ^2 if $N(\varepsilon_\Delta) = -1$. Then every $O_{\Delta,1}^*$ orbit of integral solutions of $F(x, y) = m$ contains a solution $(x, y) \in \mathbb{Z}^2$ such that $0 \leq y \leq U$, where $U = \left| \frac{am\tau_\Delta}{\Delta} \right|^{\frac{1}{2}} \left(1 - \frac{1}{\tau_\Delta}\right)$ if $am > 0$; or $U = \left| \frac{am\tau_\Delta}{\Delta} \right|^{\frac{1}{2}} \left(1 + \frac{1}{\tau_\Delta}\right)$ if $am < 0$. So for finding a set of representatives of the $O_{\Delta,1}^*$ orbits of integral solutions of $F(x, y) = m$, we must find for each integer y such that $0 \leq y \leq U$, all integers x that satisfy $F(x, y) = m$. If $F(x, y) = m$, then $\Delta y^2 + 4am = (2ax + by)^2$ and so $x = \frac{-by \pm \sqrt{\Delta y^2 + 4am}}{2a}$.

Here we notice that there are one, two, three (maybe or more) sets of representatives depending on t for the Pell equation $2x^2 - y^2 = 2t^2 - 1$. For example, for $t = 3$, the set of representatives is $\{[\pm 3, 1]\}$; for $t = 5$, the set of representatives is $\{[\pm 5, 1], [\pm 7, 7]\}$; for $t = 37$, the set of representatives is $\{[\pm 37, 1], [\pm 41, 25], [\pm 43, 31], [\pm 47, 41]\}$. To determine all (positive) integer solutions of $2x^2 - y^2 = 2t^2 - 1$ we have to put some restrictions on t . From now on, we assume that t is an integer such that $2t^2 - 1$ is a prime.

Theorem 2.1. *Let $2t^2 - 1$ be a prime for an integer $t \geq 2$. Then for the Pell equation $2x^2 - y^2 = 2t^2 - 1$, we have*

- (1) *The set of all positive integer solutions is $\Omega = \{(x_{2n}, y_{2n}), (x_{2n+1}, y_{2n+1})\}$, where*

$$\begin{aligned} [x_{2n+1} \ y_{2n+1}] &= [t \ 1]M^n \text{ for } n \geq 0 \\ [x_{2n} \ y_{2n}] &= [t \ -1]M^n \text{ for } n \geq 1, \end{aligned}$$

$$\text{and } M = \begin{bmatrix} 3 & 4 \\ 2 & 3 \end{bmatrix}.$$

- (2) *The n^{th} power of M is*

$$M^n = \begin{bmatrix} C_n & 4B_n \\ 2B_n & C_n \end{bmatrix}.$$

- (3) *The set of all positive integer solutions can be given in terms of balancing and Lucas-balancing numbers, that is,*

$$\begin{aligned} (x_{2n+1}, y_{2n+1}) &= (tC_n + 2B_n, 4tB_n + C_n) \text{ for } n \geq 0 \\ (x_{2n}, y_{2n}) &= (tC_n - 2B_n, 4tB_n - C_n) \text{ for } n \geq 1 \end{aligned}$$

or in terms of Pell numbers, that is,

$$\begin{aligned} (x_{2n+1}, y_{2n+1}) &= ((P_{2n} + P_{2n-1})t + P_{2n}, 2tP_{2n} + P_{2n} + P_{2n-1}) \\ (x_{2n}, y_{2n}) &= ((P_{2n} + P_{2n-1})t - P_{2n}, 2tP_{2n} - P_{2n} - P_{2n-1}) \end{aligned}$$

for $n \geq 1$.

Proof. (1) For the Pell equation $2x^2 - y^2 = 2t^2 - 1$, we get $\tau_8 = 3 + 2\sqrt{2}$ and $8(y^2 + 2t^2 - 1)$ is a square only for $y = 1$ in the range $0 \leq y \leq U$ and in this case $x = \pm t$. Therefore, we find that there is exactly one $O_{8,1}^*$ set of representative

of the orbits and that $\{\pm t - 1\}$ is a set of representatives. $[t - 1]M^n$ generates the solutions (x_{2n+1}, y_{2n+1}) for $n \geq 0$ and $[t - 1]M^n$ generates the solutions (x_{2n}, y_{2n}) for $n \geq 1$, where $M = \begin{bmatrix} 3 & 4 \\ 2 & 3 \end{bmatrix}$ by (2.3). So the set of all positive integer solutions of $2x^2 - y^2 = 2t^2 - 1$ is $\Omega = \{(x_{2n}, y_{2n}), (x_{2n+1}, y_{2n+1})\}$, where $[x_{2n+1} \ y_{2n+1}] = [t \ -1]M^n$ for $n \geq 0$ and $[x_{2n} \ y_{2n}] = [t \ -1]M^n$ for $n \geq 1$.

(2) We prove it by induction on n . Let us assume that $n = 1$. Then since $C_1 = 3$ and $B_1 = 1$, this relation is true. Let us assume that this relation is satisfied for $n - 1$, that is,

$$M^{n-1} = \begin{bmatrix} C_{n-1} & 4B_{n-1} \\ 2B_{n-1} & C_{n-1} \end{bmatrix}.$$

Then we get

$$(2.4) \quad M \cdot M^{n-1} = \begin{bmatrix} 3C_{n-1} + 8B_{n-1} & 12B_{n-1} + 4C_{n-1} \\ 2C_{n-1} + 6B_{n-1} & 8B_{n-1} + 3C_{n-1} \end{bmatrix}.$$

Since $3C_{n-1} + 8B_{n-1} = C_n$, $3B_{n-1} + C_{n-1} = B_n$, $C_{n-1} + 3B_{n-1} = B_n$ and $8B_{n-1} + 3C_{n-1} = C_n$, (2.4) becomes

$$M \cdot M^{n-1} = \begin{bmatrix} C_n & 4B_n \\ 2B_n & C_n \end{bmatrix} = M^n.$$

(3) From (1) and (2), it is easily seen that $(x_{2n+1}, y_{2n+1}) = (tC_n + 2B_n, 4tB_n + C_n)$ for $n \geq 0$ and $(x_{2n}, y_{2n}) = (tC_n - 2B_n, 4tB_n - C_n)$ for $n \geq 1$. The last assertion is obvious since $P_{2n} = 2B_n$ and $P_{2n} + P_{2n-1} = C_n$. $\square$

Hence we can give the following main result.

Theorem 2.2. (1) *For balancing numbers, we have*

$$\begin{aligned} 16B_nC_n + 32B_n^2 + 2C_n^2 + 2 &= (4b_{n+1} + 2)^2 \\ 24B_nC_n + 32B_n^2 + 4C_n^2 &= 2c_{n+1}(4b_{n+1} + 2) \\ 8B_n^2 + 2C_n^2 + 8B_nC_n - 1 &= c_{n+1}^2 \end{aligned}$$

for $n \geq 1$.

(2) *The general terms of all t -balancing numbers can be given in terms of balancing numbers as*

$$\begin{aligned} B_{2n-1}^t &= \frac{(4B_n + C_n - 1)t - (2B_n + C_n + 1)}{2} \\ b_{2n-1}^t &= \frac{(C_n - 1)t - 2B_n}{2} \\ C_{2n-1}^t &= (4b_{n+1} + 2)t - c_{n+1} \\ c_{2n-1}^t &= 4tB_n - C_n \\ B_{2n}^t &= \frac{(4B_n + C_n - 1)t + (2B_n + C_n - 1)}{2} \end{aligned}$$

$$\begin{aligned}
b_{2n}^t &= \frac{(C_n - 1)t + 2B_n}{2} \\
C_{2n}^t &= (4b_{n+1} + 2)t + c_{n+1} \\
c_{2n}^t &= 4tB_n + C_n
\end{aligned}$$

for $n \geq 1$.

- (3) *The general terms of balancing numbers can be given in terms of t -balancing numbers as*

$$\begin{aligned}
B_n &= \frac{b_{2n}^t - b_{2n-1}^t}{2} \quad \text{and} \quad C_n = \frac{c_{2n}^t - c_{2n-1}^t}{2} \quad \text{for } n \geq 1 \\
b_n &= \frac{C_{2n-2}^t + C_{2n-3}^t - 4t}{8t} \quad \text{and} \quad c_n = \frac{C_{2n-2}^t - C_{2n-3}^t}{2} \quad \text{for } n \geq 2.
\end{aligned}$$

- (4) *Binet formulas for t -balancing numbers are*

$$\begin{aligned}
B_{2n-1}^t &= t \left(\frac{\alpha^{2n+1} + \beta^{2n+1} - 2}{4} \right) - \frac{\alpha^{2n+1} - \beta^{2n+1} + 2\sqrt{2}}{4\sqrt{2}} \\
b_{2n-1}^t &= t \left(\frac{\alpha^{2n} + \beta^{2n} - 2}{4} \right) - \frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} \\
C_{2n-1}^t &= t \left(\frac{\alpha^{2n+1} - \beta^{2n+1}}{\sqrt{2}} \right) - \frac{\alpha^{2n+1} + \beta^{2n+1}}{2} \\
c_{2n-1}^t &= t \left(\frac{\alpha^{2n} - \beta^{2n}}{\sqrt{2}} \right) - \frac{\alpha^{2n} + \beta^{2n}}{2} \\
B_{2n}^t &= t \left(\frac{\alpha^{2n+1} + \beta^{2n+1} - 2}{4} \right) + \frac{\alpha^{2n+1} - \beta^{2n+1} - 2\sqrt{2}}{4\sqrt{2}} \\
b_{2n}^t &= t \left(\frac{\alpha^{2n} + \beta^{2n} - 2}{4} \right) + \frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} \\
C_{2n}^t &= t \left(\frac{\alpha^{2n+1} - \beta^{2n+1}}{\sqrt{2}} \right) + \frac{\alpha^{2n+1} + \beta^{2n+1}}{2} \\
c_{2n}^t &= t \left(\frac{\alpha^{2n} - \beta^{2n}}{\sqrt{2}} \right) + \frac{\alpha^{2n} + \beta^{2n}}{2}
\end{aligned}$$

for $n \geq 1$.

- (5) *The general terms of Pell and Pell-Lucas numbers can be given in terms of t -balancing numbers as*

$$\begin{aligned}
P_{2n} &= b_{2n}^t - b_{2n-1}^t, \quad P_{2n+1} = \frac{C_{2n}^t + C_{2n-1}^t}{4t}, \\
Q_{2n} &= \frac{b_{4n}^t - b_{4n-1}^t}{b_{2n}^t - b_{2n-1}^t} \quad \text{and} \quad Q_{2n+1} = C_{2n}^t - C_{2n-1}^t
\end{aligned}$$

for $n \geq 1$.

Proof. (1) Note that $B_n = \frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}}$, $C_n = \frac{\alpha^{2n} + \beta^{2n}}{2}$ and $b_n = \frac{\alpha^{2n-1} - \beta^{2n-1}}{4\sqrt{2}} - \frac{1}{2}$. So

$$\begin{aligned}
 & 16B_nC_n + 32B_n^2 + 2C_n^2 + 2 \\
 &= 16\left(\frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}}\right)\left(\frac{\alpha^{2n} + \beta^{2n}}{2}\right) + 32\left(\frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}}\right)^2 + 2\left(\frac{\alpha^{2n} + \beta^{2n}}{2}\right)^2 + 2 \\
 &= \frac{\alpha^{4n+2} - 2(\alpha\beta)^{2n+1} + \beta^{4n+2}}{2} \\
 &= 16\left(\frac{\alpha^{2n+1} - \beta^{2n+1}}{4\sqrt{2}} - \frac{1}{2}\right)^2 + 16\left(\frac{\alpha^{2n+1} - \beta^{2n+1}}{4\sqrt{2}}\right) - 4 \\
 &= 16\left(\frac{\alpha^{2n+1} - \beta^{2n+1}}{4\sqrt{2}} - \frac{1}{2}\right)^2 + 16\left(\frac{\alpha^{2n+1} - \beta^{2n+1}}{4\sqrt{2}} - \frac{1}{2}\right) + 4 \\
 &= 16b_{n+1}^2 + 16b_{n+1} + 4 = (4b_{n+1} + 2)^2.
 \end{aligned}$$

The others can be proved similarly.

(2) Since $x = 2r + t$, we get from (3) of Theorem 2.1 that

$$b_{2n-1}^t = \frac{(C_n - 1)t - 2B_n}{2}.$$

Thus from (1.6), we get

$$\begin{aligned}
 c_{2n-1}^t &= \sqrt{8(b_{2n-1}^t)^2 + 8tb_{2n-1}^t + 1} \\
 &= \sqrt{8\left(\frac{(C_n - 1)t - 2B_n}{2}\right)^2 + 8t\left(\frac{(C_n - 1)t - 2B_n}{2}\right) + 1} \\
 &= \sqrt{16t^2B_n^2 - 8tB_nC_n + C_n^2} \\
 &= 4tB_n - C_n
 \end{aligned}$$

since $8B_n^2 + 1 = C_n^2$. So from (1.5), we deduce that

$$B_{2n-1}^t = \frac{(4B_n + C_n - 1)t - (2B_n + C_n + 1)}{2}$$

and hence

$$\begin{aligned}
 c_{2n-1}^t &= \sqrt{8(B_{2n-1}^t)^2 + 8B_{2n-1}^t(1+t) + (2t+1)^2} \\
 &= \sqrt{8\left(\frac{(4B_n + C_n - 1)t - (2B_n + C_n + 1)}{2}\right)^2 + 8\left(\frac{(4B_n + C_n - 1)t - (2B_n + C_n + 1)}{2}\right)(1+t) + (2t+1)^2} \\
 &= \sqrt{t^2(16B_nC_n + 32B_n^2 + 2C_n^2 + 2) - t(24B_nC_n + 32B_n^2 + 4C_n^2) + (8B_n^2 + 2C_n^2 + 8B_nC_n - 1)} \\
 &= \sqrt{t^2(4b_{n+1} + 2)^2 - 2tc_{n+1}(4b_{n+1} + 2) + c_{n+1}^2}
 \end{aligned}$$

$$= (4b_{n+1} + 2)t - c_{n+1}$$

by (1). The others can be proved similarly.

(3) Since $b_{2n-1}^t = \frac{(C_n-1)t-2B_n}{2}$ and $b_{2n}^t = \frac{(C_n-1)t+2B_n}{2}$ by (2), we easily deduce that $\frac{b_{2n}^t - b_{2n-1}^t}{2} = B_n$. The others are similar.

(4) Recall that $B_n = \frac{\alpha^{2n}-\beta^{2n}}{4\sqrt{2}}$ and $C_n = \frac{\alpha^{2n}+\beta^{2n}}{2}$. So we get from (2) that

$$\begin{aligned} B_{2n-1}^t &= \frac{(4B_n + C_n - 1)t - (2B_n + C_n + 1)}{2} \\ &= \frac{t \left[4 \left(\frac{\alpha^{2n}-\beta^{2n}}{4\sqrt{2}} \right) + \frac{\alpha^{2n}+\beta^{2n}}{2} - 1 \right] - 2 \left(\frac{\alpha^{2n}-\beta^{2n}}{4\sqrt{2}} \right) - \frac{\alpha^{2n}+\beta^{2n}}{2} - 1}{2} \\ &= \frac{t \left(\frac{\alpha^{2n}(1+\sqrt{2})+\beta^{2n}(1-\sqrt{2})-2}{2} \right) - \frac{\alpha^{2n}(1+\sqrt{2})-\beta^{2n}(1-\sqrt{2})+2\sqrt{2}}{2\sqrt{2}}}{2} \\ &= t \left(\frac{\alpha^{2n+1} + \beta^{2n+1} - 2}{4} \right) - \frac{\alpha^{2n+1} - \beta^{2n+1} + 2\sqrt{2}}{4\sqrt{2}}. \end{aligned}$$

The others can be proved similarly.

(5) Note that $b_{2n-1}^t = \frac{(C_n-1)t-2B_n}{2}$ and $b_{2n}^t = \frac{(C_n-1)t+2B_n}{2}$ by (2). So

$$b_{2n}^t - b_{2n-1}^t = \frac{(C_n - 1)t + 2B_n}{2} - \frac{(C_n - 1)t - 2B_n}{2} = P_{2n}$$

since $B_n = \frac{P_{2n}}{2}$. The others can be proved similarly. $\square$

2.1. Sums. In this subsection, we consider the sums of numbers we mentioned above.

Theorem 2.3. (1) *For the sums of t -balancing numbers, we have*

$$\begin{aligned} \sum_{i=1}^n B_i^t &= (B_{\frac{n+2}{2}} + b_{\frac{n+2}{2}} - \frac{n+2}{2})t - \frac{n}{2} \\ \sum_{i=1}^n b_i^t &= (B_{\frac{n}{2}} + b_{\frac{n+2}{2}} - \frac{n}{2})t \\ \sum_{i=1}^n C_i^t &= (3B_{\frac{n+2}{2}} + B_{\frac{n}{2}} + 2b_{\frac{n+2}{2}} - 3)t \\ \sum_{i=1}^n c_i^t &= 4b_{\frac{n+2}{2}}t \end{aligned}$$

for even $n \geq 2$ or

$$\sum_{i=1}^n B_i^t = (B_{\frac{n+3}{2}} - B_{\frac{n+1}{2}} - \frac{n+3}{2})t - b_{\frac{n+3}{2}} - \frac{n+1}{2}$$

$$\begin{aligned}\sum_{i=1}^n b_i^t &= (2B_{\frac{n+1}{2}} - \frac{n+1}{2})t - B_{\frac{n+1}{2}} \\ \sum_{i=1}^n C_i^t &= (b_{\frac{n+5}{2}} - b_{\frac{n+1}{2}} - 4)t - 2B_{\frac{n+3}{2}} + 2b_{\frac{n+3}{2}} + 1 \\ \sum_{i=1}^n c_i^t &= 4(B_{\frac{n+1}{2}} + b_{\frac{n+1}{2}})t - 2B_{\frac{n+1}{2}} - 2b_{\frac{n+1}{2}} - 1\end{aligned}$$

for odd $n \geq 1$.

(2) For the sums of Pell numbers, we have

$$\begin{aligned}\sum_{i=1}^n P_{2i-1} &= \frac{b_{2n}^t - b_{2n-1}^t}{2} \\ \sum_{i=1}^n P_{2i} &= \frac{C_{2n}^t + C_{2n-1}^t - 4t}{8t} \\ \sum_{i=0}^{2n} P_{2i+1} &= \frac{(C_{2n}^t + C_{2n-1}^t)(C_{2n}^t - C_{2n-1}^t)}{8t} \\ \sum_{i=1}^{2n} P_{2i} &= \frac{(b_{2n}^t - b_{2n-1}^t)(C_{2n}^t - C_{2n-1}^t)}{2} \\ \sum_{i=0}^{2n} (P_{2i+1} + P_{2i+2}) &= \frac{(c_{2n+2}^t - c_{2n+1}^t)(C_{2n}^t - C_{2n-1}^t)}{4}.\end{aligned}$$

(3) For the sums of Pell–Lucas numbers, we have

$$\begin{aligned}\sum_{i=0}^{2n} Q_i &= \frac{2(b_{4n+2}^t - b_{4n+1}^t)}{C_{2n}^t - C_{2n-1}^t} \\ \sum_{i=1}^{2n} Q_{2i} &= \frac{(b_{2n}^t - b_{2n-1}^t)(C_{2n}^t + C_{2n-1}^t)}{t}.\end{aligned}$$

(4) For the sums of balancing numbers, we have

$$\begin{aligned}\sum_{i=1}^{2n} B_i &= \frac{(b_{2n}^t - b_{2n-1}^t)(C_{2n}^t - C_{2n-1}^t)}{4} \\ \sum_{i=1}^{2n} B_{2i} &= \frac{(b_{2n}^t - b_{2n-1}^t)(c_{2n}^t - c_{2n-1}^t)(b_{4n+2}^t - b_{4n+1}^t)}{4} \\ \sum_{i=1}^{2n} (B_i + B_{i+1}) &= 2(b_{2n}^t - b_{2n-1}^t)(b_{2n+2}^t - b_{2n+1}^t)\end{aligned}$$

$$\sum_{i=0}^{2n} B_{2i+1} = \frac{(C_{2n}^t + C_{2n-1}^t)(C_{2n}^t - C_{2n-1}^t)(b_{4n+2}^t - b_{4n+1}^t)}{16t}$$

$$\sum_{i=0}^{2n} (B_{2i+1} + B_{2i+2}) = \frac{(C_{4n+2}^t - C_{4n+1}^t)(b_{4n+2}^t - b_{4n+1}^t)}{4}.$$

(5) For the sums of Lucas–cobalancing numbers, we have

$$\sum_{i=1}^{2n+1} c_{i+1} = \frac{(C_{2n}^t - C_{2n-1}^t)(C_{2n+2}^t - C_{2n+1}^t)}{4}$$

$$\sum_{i=1}^{2n+1} c_{2i+1} = \frac{(C_{2n}^t - C_{2n-1}^t)(C_{2n}^t + C_{2n-1}^t)(C_{4n+4}^t - C_{4n+3}^t)}{16t}.$$

Proof. (1) Let n be even, say $n = 2k$ for an integer $k \geq 1$. Then from (4) of Theorem 2.2, we easily get

$$\begin{aligned} \sum_{i=1}^{2k} B_i^t &= B_1^t + B_2^t + \cdots + B_{2k}^t \\ &= \left[\left(\frac{\alpha^3 + \beta^3 - 2}{4} \right) t - \frac{\alpha^3 - \beta^3 + 2\sqrt{2}}{4\sqrt{2}} \right] \\ &\quad + \left[\left(\frac{\alpha^3 + \beta^3 - 2}{4} \right) t + \frac{\alpha^3 - \beta^3 - 2\sqrt{2}}{4\sqrt{2}} \right] \\ &\quad + \cdots + \left[\left(\frac{\alpha^{2k+1} + \beta^{2k+1} - 2}{4} \right) t - \frac{\alpha^{2k+1} - \beta^{2k+1} - 2\sqrt{2}}{4\sqrt{2}} \right] \\ &= \left(\frac{\alpha^3 + \alpha^5 + \cdots + \alpha^{2k+1} + \beta^3 + \beta^5 + \cdots + \beta^{2k+1}}{2} - k \right) t - k \\ &= \left(\frac{\alpha^{2k+2} - \beta^{2k+2}}{4\sqrt{2}} + \frac{\alpha^{2k+1} - \beta^{2k+1}}{4\sqrt{2}} - \frac{1}{2} - \frac{2k+2}{2} \right) t - k \\ &= \left(B_{\frac{n+2}{2}} + b_{\frac{n+2}{2}} - \frac{n+2}{2} \right) t - \frac{n}{2}. \end{aligned}$$

The others can be proved similarly. $\square$

In [11], Santana and Diaz–Barrero proved that the sum of first nonzero $4n+1$ terms of Pell numbers is a perfect square, that is,

$$(2.5) \quad \sum_{i=1}^{4n+1} P_i = \left(\sum_{i=0}^n \binom{2n+1}{2i} 2^i \right)^2.$$

Later in [13, Theorem 2.1], Tekcan and Tayat proved that the sum of first nonzero $2n+1$ terms of Pell numbers is a perfect square if n is even or half of

a perfect square if n is odd, that is,

$$\sum_{i=1}^{2n+1} P_i = \begin{cases} \left(\frac{\alpha^{n+1} + \beta^{n+1}}{2} \right)^2 & \text{for even } n \\ \left(\frac{\alpha^{n+1} - \beta^{n+1}}{\sqrt{2}} \right)^2 & \text{for odd } n. \end{cases}$$

They set $X_n = \frac{\alpha^{n+1} + \beta^{n+1}}{2}$ and $Y_n = \frac{\alpha^{n+1} - \beta^{n+1}}{\sqrt{2}}$ for $n \geq 0$ and proved that the right hand side of (2.5) is $(2X_n^2 - 2X_n Y_{n-1} + (-1)^{n+1})^2$. Similarly, we can give the following theorem.

Theorem 2.4. *Let P_n denote the n^{th} Pell number, let Q_n denote the n^{th} Pell-Lucas number and let B_n denote the n^{th} balancing number. Then*

- (1) *The sum of Pell numbers from 1 to $4n - 3$ is a perfect square and is*

$$\sum_{i=1}^{4n-3} P_i = \left(\frac{C_{2n-2}^t - C_{2n-3}^t}{2} \right)^2$$

for $n \geq 2$.

- (2) *The sum of Pell numbers from 1 to $4n - 1$ and adding 1 is a perfect square and is*

$$1 + \sum_{i=1}^{4n-1} P_i = \left(\frac{C_{2n}^t - C_{2n-1}^t}{2} \right)^2$$

for $n \geq 1$.

- (3) *The sum of Pell numbers from 1 to $2n - 1$ is a perfect square and is*

$$\sum_{i=1}^{2n-1} P_i = \left(\frac{C_{n-1}^t - C_{n-2}^t}{2} \right)^2$$

for odd $n \geq 3$, and the half of the sum of Pell numbers from 1 to $2n - 1$ is a perfect square and is

$$\frac{\sum_{i=1}^{2n-1} P_i}{2} = (b_n^t - b_{n-1}^t)^2$$

for even $n \geq 2$.

- (4) *The sum of $(2i - 1)^{\text{st}}$ Pell-Lucas numbers from 1 to $2n$ is a perfect square and is*

$$\sum_{i=1}^{2n} Q_{2i-1} = (2(b_{2n}^t - b_{2n-1}^t))^2$$

for $n \geq 1$.

- (5) *The half of the sum of $(2i+1)^{st}$ Pell-Lucas numbers from 0 to $2n$ is a perfect square and is*

$$\frac{\sum_{i=0}^{2n} Q_{2i+1}}{2} = \left(\frac{C_{2n}^t - C_{2n-1}^t}{2} \right)^2$$

for $n \geq 1$.

- (6) *The sum of $(2i-1)^{st}$ balancing numbers from 1 to $2n$ is a perfect square and is*

$$\sum_{i=1}^{2n} B_{2i-1} = \left(\frac{b_{4n}^t - b_{4n-1}^t}{2} \right)^2$$

for $n \geq 1$.

Proof. (1) Note that $P_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}$ and $\sum_{i=1}^n P_i = \frac{P_n + P_{n+1} - 1}{2}$. So

$$\begin{aligned} \sum_{i=1}^{4n-3} P_{2i-1} &= \frac{P_{4n-3} + P_{4n-2} - 1}{2} \\ &= \frac{\frac{\alpha^{4n-3} - \beta^{4n-3}}{2\sqrt{2}} + \frac{\alpha^{4n-2} - \beta^{4n-2}}{2\sqrt{2}} - 1}{2} \\ &= \frac{\alpha^{4n-2} + \beta^{4n-2} - 2}{4} \\ &= \left(\frac{(4b_n + 2)t + c_n - (4b_n + 2)t + c_n}{2} \right)^2 \\ &= \left(\frac{C_{2n-2}^t - C_{2n-3}^t}{2} \right)^2 \end{aligned}$$

by (2) of Theorem 2.2. The other cases can be proved similarly. $\square$

2.2. Relationship with Triangular Numbers. In this subsection, we consider the relationship between t -balancing numbers and triangular numbers which are the numbers of the form $T_n = \frac{n(n+1)}{2}$ for $n \geq 0$. There are infinitely many triangular numbers that are also square numbers which are called square triangular numbers and is denoted by S_n . For the n^{th} square triangular number S_n , we can write

$$S_n = s_n^2 = \frac{t_n(t_n + 1)}{2},$$

where s_n and t_n are the sides of the corresponding square and triangle.

In the following theorem, we can give the general terms of s_n, t_n and S_n in terms of t -balancing numbers and contrary, we can give the general terms of all t -balancing numbers in terms of squares and triangles.

Theorem 2.5. (1) *The general terms of s_n, t_n and S_n are*

$$s_n = \frac{b_{2n}^t - b_{2n-1}^t}{2}, \quad t_n = \frac{c_{2n}^t - c_{2n-1}^t - 2}{4}, \quad S_n = \left(\frac{C_{2n}^t - C_{2n-1}^t - b_{2n+2}^t + b_{2n+1}^t}{2} \right)^2$$

for $n \geq 1$.

(2) *The general terms of all t -balancing numbers are*

$$\begin{aligned} B_{2n-1}^t &= (t_n + 2s_n)t - (s_n + t_n + 1) \\ b_{2n-1}^t &= t_n t - s_n \\ C_{2n-1}^t &= (4s_n + 4t_n + 2)t - (s_n + s_{n+1}) \\ c_{2n-1}^t &= 4s_n t - (2t_n + 1) \\ B_{2n}^t &= (t_n + 2s_n)t + (s_n + t_n) \\ b_{2n}^t &= t_n t + s_n \\ C_{2n}^t &= (4s_n + 4t_n + 2)t + (s_n + s_{n+1}) \\ c_{2n}^t &= 4s_n t + (2t_n + 1) \end{aligned}$$

for $n \geq 1$.

Proof. (1) Since $s_n = \frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}}$, $t_n = \frac{\alpha^{2n} + \beta^{2n} - 2}{4}$ and $S_n = \left(\frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} \right)^2$, we deduce from (4) of Theorem 2.2 that

$$\begin{aligned} s_n &= \frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} \\ &= \frac{\left(t \left(\frac{\alpha^{2n} + \beta^{2n} - 2}{4} \right) + \frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} \right) - \left(t \left(\frac{\alpha^{2n} + \beta^{2n} - 2}{4} \right) - \frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} \right)}{2} \\ &= \frac{b_{2n}^t - b_{2n-1}^t}{2} \end{aligned}$$

and

$$\begin{aligned} t_n &= \frac{\alpha^{2n} + \beta^{2n} - 2}{4} \\ &= \frac{\left(t \left(\frac{\alpha^{2n} - \beta^{2n}}{\sqrt{2}} \right) + \frac{\alpha^{2n} + \beta^{2n}}{2} \right) - \left(t \left(\frac{\alpha^{2n} - \beta^{2n}}{\sqrt{2}} \right) - \frac{\alpha^{2n} + \beta^{2n}}{2} \right) - 2}{4} \\ &= \frac{c_{2n}^t - c_{2n-1}^t - 2}{4}. \end{aligned}$$

Similarly it can be showed that $S_n = \left(\frac{C_{2n}^t - C_{2n-1}^t - b_{2n+2}^t + b_{2n+1}^t}{2} \right)^2$.

(2) We get from (2) and (4) of Theorem 2.2 that

$$\begin{aligned} B_{2n-1}^t &= \frac{(4B_n + C_n - 1)t - (2B_n + C_n + 1)}{2} \\ &= \frac{\left[4 \left(\frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} \right) + \frac{\alpha^{2n} + \beta^{2n}}{2} - 1 \right] t - \left[2 \left(\frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} \right) + \frac{\alpha^{2n} + \beta^{2n}}{2} + 1 \right]}{2} \end{aligned}$$

$$\begin{aligned}
&= \left(\frac{\alpha^{2n+1} + \beta^{2n+1} - 2}{4} \right) t - \left(\frac{\alpha^{2n+1} - \beta^{2n+1} + 2\sqrt{2}}{4\sqrt{2}} \right) \\
&= \frac{(\alpha^{2n} + \beta^{2n} - 2 + \sqrt{2}\alpha^{2n} - \sqrt{2}\beta^{2n})t}{4} - \frac{\alpha^{2n} - \beta^{2n} + \sqrt{2}\alpha^{2n} + \sqrt{2}\beta^{2n} + 2\sqrt{2}}{4\sqrt{2}} \\
&= \left(\frac{\alpha^{2n} + \beta^{2n} - 2}{4} + \frac{\alpha^{2n} - \beta^{2n}}{2\sqrt{2}} \right) t - \left(\frac{\alpha^{2n} - \beta^{2n}}{4\sqrt{2}} + \frac{\alpha^{2n} + \beta^{2n} - 2}{4} + 1 \right) \\
&= (t_n + 2s_n)t - (s_n + t_n + 1).
\end{aligned}$$

The others can be proved similarly. $\square$

REFERENCES

- [1] A. Behera and G.K. Panda. *On the Square Roots of Triangular Numbers*. The Fibonacci Quarterly, **37**(2)(1999), 98–105.
- [2] K.K. Dash, R.S. Ota and S. Dash. *t-Balancing Numbers*. Int. J. Contemp. Math. Sciences, **7**(41)(2012), 1999–2012.
- [3] R. Finkelstein. *The House Problem*. Am. Math. Mon. **72**(1965), 1082–1088.
- [4] T. Kovacs, L. Liptai and P. Olajos. *On (a, b)-Balancing Numbers*. Publ. Math. Debrecen **77**/3–4(2010), 485–498.
- [5] K. Liptai. *Lucas Balancing Numbers*. Acta Math. Univ. Ostrav. **14**(2006), 43–47.
- [6] K. Liptai, F. Luca, Á. Pinter and L. Szalay. *Generalized Balancing Numbers*. Indag. Mathem., N.S. **20**(1)(2009), 87–100.
- [7] P. Olajos. *Properties of Balancing, Cobalancing and Generalized Balancing Numbers*. Annales Mathematicae et Informaticae **37**(2010), 125–138.
- [8] G.K. Panda. *Some Fascinating Properties of Balancing Numbers*. Proceedings of the Eleventh International Conference on Fibonacci Numbers and their Applications, Cong. Numer. **194**(2009), 185–189.
- [9] G.K. Panda and P.K. Ray. *Some Links of Balancing and Cobalancing Numbers with Pell and Associated Pell Numbers*. Bul. of Inst. of Math. Acad. Sinica **6**(1)(2011), 41–72.
- [10] G.K. Panda and P.K. Ray. *Cobalancing Numbers and Cobalancers*. Int. J. Math. Sci. **8**(2005), 1189–1200.
- [11] S.F. Santana and J.L. Diaz-Barrero. *Some Properties of Sums Involving Pell Numbers*. Missouri Journal of Mathematical Science **18**(1)(2006), 33–40.
- [12] L. Szalay. *On the Resolution of Simultaneous Pell Equations*. Ann. Math. Info. **34**(2007), 77–87.
- [13] A. Tekcan and M. Tayat. *Generalized Pell Numbers, Balancing Numbers and Binary Quadratic Forms*. Creative Mathematics and Inf. **23**(1)(2014), 115–122.

Received January 27, 2016.

ULUDAG UNIVERSITY
 FACULTY OF SCIENCE
 DEPARTMENT OF MATHEMATICS
 BURSA-TURKIYE
E-mail address: `tekcan@uludag.edu.tr`