

A NOTE ON GENERALIZED PSEUDO-RICCI SYMMETRIC MANIFOLD

KANAK KANTI BAISHYA

ABSTRACT. The object of the present paper is to introduce the notion of generalized pseudo-Ricci symmetric space with a non-trivial example. The beauty of such space is that it has the flavour of Ricci symmetric space, Ricci recurrent space, generalized Ricci recurrent space and pseudo-Ricci symmetric space. Furthermore, having found a faulty example in [14], the present paper attempts to construct a new example of pseudo-Ricci symmetric manifold.

1. INTRODUCTION

In the sense of Chaki, a non-flat n -dimensional Riemannian manifold (M^n, g) , ($n > 3$) is said to be a pseudo-Ricci symmetric manifold [9], if its Ricci tensor S of type $(0, 2)$ is not identically zero and satisfies the following equation

$$(1) \quad (\nabla_X S)(Y, U) = 2B(X)S(Y, U) + B(Y)S(X, U) + B(U)S(X, Y)$$

for all vector fields $X, Y, U \in \chi(M^n)$, where B is a nonzero 1-form defined by $B(X) = g(X, \varrho) \forall X$, where ϱ is called the associated vector field to the 1-form, $\chi(M^n)$ denotes the Lie algebra of all smooth vector fields over $C^\infty(M^n)$ on the manifold M^n and ∇ is the operator of the covariant differentiation with respect to the metric tensor g . The local expression of the above equation is

$$R_{ik,l} = 2B_l R_{ik} + B_i R_{kl} + B_k R_{il},$$

2020 *Mathematics Subject Classification.* 53C15, 53C25.

Key words and phrases. generalized pseudo Ricci symmetric manifolds, quasi-Einstein.

We have made all the calculations in Wolfram mathematica. The author gratefully acknowledge the financial support of UGC, ERO, Kolkata, India.

where B_l is non-zero co-vectors and comma followed by indices denotes the covariant differentiation with respect to the metric tensor g . An n -dimensional manifold of this kind is abbreviated by $(PRS)_n$.

Keeping the tune of Dubey [11], in the present paper, we attempt to introduce the notion of *generalized pseudo-Ricci symmetric manifold* which is abbreviated by $(GPRS)_n$ -manifold and defined as follows. A non-flat n -dimensional Riemannian manifold (M^n, g) ($n > 3$) is termed as generalized pseudo-Ricci symmetric manifold if its Ricci tensor S of type $(0, 2)$ is not identically zero and fulfills the identity

$$(2) \quad (\nabla_X S)(Y, U) = 2B(X)S(Y, U) + B(Y)S(X, U) + B(U)S(X, Y) \\ + 2C(X)g(Y, U) + C(Y)g(X, U) + C(U)g(X, Y),$$

where B and C are two non-zero 1-forms defined by $B(X) = g(X, \varrho)$ and $C(X) = g(X, \pi)$. The local expression of the above equation is

$$R_{ik,l} = 2B_l R_{ik} + B_i R_{lk} + B_k R_{il} \\ + 2C_l g_{ik} + C_i g_{lk} + C_k g_{il},$$

where B_l and C_l are two non-zero co-vectors. The beauty of such $(GPRS)_n$ -space is that it has the flavour of

- (1) Ricci symmetric space in the sense of Cartan (for $B = 0 = C$),
- (2) Ricci recurrent space by Patterson [4] (for $B(X) \neq 0$, $B(Y) = B(U) = 0$ and $C(V) = 0$, $\forall V$),
- (3) generalized Ricci recurrent space by De et al. [13] (for $B(X) \neq 0$, $C(X) \neq 0$ and $B(Y) = B(U) = 0 = C(Y) = C(U)$),
- (4) pseudo-Ricci symmetric space [8] (for $B \neq 0$ and $C(V) = 0$, $\forall V$).

We structured our paper as follows: Section 2 is concerned with *generalized pseudo-Ricci symmetric manifold* and obtained some interesting results of conformally flat $(GPRS)_n$ -manifold. In section 3, we cite an example of a manifold $(\mathbb{R}^4, g)$ which is a pseudo-Ricci symmetric for some choice of the 1-forms but fails to be a generalized pseudo-Ricci symmetric space. Finally, we observe that there exists a manifold $(\mathbb{R}^4, g)$ which is a generalized pseudo-Ricci symmetric space in some cases and a pseudo-Ricci symmetric space in some other cases depending on the choice of the 1-forms.

2. $(GPRS)_n$ -manifold

In this section, we assume a non-flat n -dimensional Riemannian manifold (M^n, g) , ($n > 3$) to be a generalized pseudo-Ricci symmetric manifold. Next, if the 1-form B is co-directional to C , that is,

$C(X) = \phi B(X) \forall X$, where ϕ is a constant, then the relation (2) turns into

$$(\nabla_X Z)(Y, U) = 2B(X)Z(Y, U) + B(Y)Z(X, U) + B(U)Z(X, U),$$

where $Z(X, Y) = S(X, Y) + \phi g(X, Y)$ is a well known Z -tensor introduced in ([2], [3]). This leads to the following statement.

Theorem 1. *Every $(GPRS)_n$ -manifold is a pseudo Z -symmetric manifold provided the vector fields associated to the 1-forms B and C are co-directional to each other.*

Definition 1. A non-flat Riemannian manifold (M^n, g) , $(n > 3)$ is said to be a quasi-Einstein manifold [10] if its Ricci tensor S of type $(0, 2)$ is not identically zero and satisfies the condition

$$S(X, Y) = \lambda g(X, Y) + \mu \psi(X)\psi(Y),$$

where $\lambda, \mu \in \mathbb{R}$ and ψ is a non-zero 1-form such that $g(X, U) = \psi(X)$, for all vector fields X , where U is a unit vector.

Now, contracting Y over U in (1) we obtain

$$dr(X) = 2rB(X) + 2\bar{B}(X) + (2n + 2)C(X),$$

where $\bar{B}(X) = S(X, \theta)$. Again, from (1), one can easily see

$$(3) \quad (\nabla_X S)(Y, U) - (\nabla_U S)(X, Y) = B(X)S(Y, U) - B(U)S(X, Y) \\ + C(X)g(Y, U) - C(U)g(X, Y)$$

after contraction, which leaves

$$(4) \quad dr(X) = 2rB(X) - 2\bar{B}(X) + 2(n - 1)C(X).$$

It is known ([7], p. 41) that a conformally flat (M^n, g) possesses the relation

$$(5) \quad (\nabla_X S)(Y, U) - (\nabla_U S)(X, Y) = \frac{[g(Y, U)dr(X) - g(X, Y)dr(U)]}{2(n - 1)}.$$

By virtue of (3), (4), and (5) we find

$$(6) \quad (n - 1)[B(X)S(Y, U) - B(U)S(X, Y)] \\ = [rB(X) - \bar{B}(X)]g(Y, U) - [rB(U) - \bar{B}(U)]g(X, Y),$$

which yields

$$(7) \quad B(X)\bar{B}(U) = B(U)\bar{B}(X)$$

for $Y = \varrho$. Assuming that the Ricci tensor of the manifold is codazzi type (in the sense of [12]) and then using (4), we obtain, from (7), that

$$(8) \quad B(X)C(U) = B(U)C(X), \forall X \text{ and } U.$$

This motivate us to state the following.

Proposition 1. *In a conformally flat $(GPRS)_4$ -manifold with codazzi type of Ricci tensor, the vector fields associated to the 1-forms B and C are co-directional.*

Again, for constant scalar curvature tensor (or codazzi type of Ricci tensor) by virtue of (4), (6), (8), we can easily get

$$(9) \quad S(Y, U) = -\frac{C(\varrho)}{B(\varrho)}g(Y, U) + \frac{1}{B(\varrho)}[rB(Y) + nC(Y)]B(U),$$

where $\frac{C(U)}{B(U)} = k, \forall U$. If the vector fields associated to the 1-forms B and C are co-directional, then (9) takes the following form

$$S(Y, U) = \alpha g(Y, U) + \beta B(Y)B(U).$$

This leads to the followings.

Theorem 2. *A conformally flat $(GPRS)_n$ -manifold with codazzi type of Ricci tensor is a quasi-Einstein manifold.*

Corollary 1. *A conformally flat generalized pseudo-Ricci symmetric manifold with constant scalar curvature is a space of quasi constant curvature.*

3. EXISTENCE OF $(PRS)_4$ -SPACE

In the example given in ([14], Example 3.1, p. 214-215) authors have calculated or assumed the value of the covariant derivatives corresponding to the vanishing components of the Ricci tensor R_{14} , R_{24} , and R_{34} (namely, $R_{14,1}$, $R_{24,2}$, and $R_{34,3}$) to be zero. But these values are found to be $R_{14,1} = R_{24,2} = R_{34,3} = -\frac{8}{9(x^4)^{5/3}}$ which is non-zero. Consequently, their choice of the 1-forms

$$B_i(x) = \begin{cases} -\frac{1}{x^4} & \text{for } i = 4, \\ 0 & \text{otherwise;} \end{cases}$$

and the relations

$$R_{14,1} = 2B_1R_{14} + B_1R_{14} + B_4R_{11}$$

$$R_{24,2} = 2B_2R_{24} + B_2R_{24} + B_4R_{22}$$

$$R_{34,3} = 2B_3R_{34} + B_3R_{34} + B_4R_{33}$$

do not stand as $R_{11} = R_{22} = R_{33} \neq 0$. Hence, the $(\mathbb{R}^4, g)$ under-considered metric ([14], p. 214) cannot be a pseudo-Ricci symmetric manifold.

Example 1. Let $(\mathbb{R}^4, g)$ be a 4-dimensional Riemannian space endowed with the Riemannian metric g given by

$$ds^2 = g_{ij}dx^i dx^j = (dx^2)^2 + 2e^{x^2}[dx^1 dx^2 + dx^3 dx^4]$$

for $i, j = 1, 2, 3, 4$. The non-zero components of Riemannian curvature tensor, Ricci tensors and scalar curvature are (up to symmetry and skew-symmetry)

$$R_{2324} = \frac{1}{4}e^{x^2}, \quad R_{22} = \frac{1}{2}, \quad r = 0.$$

Covariant derivatives of Ricci tensors is expressed as

$$R_{22,2} = -1.$$

For the 1-form

$$B_i = \begin{cases} -1 & \text{for } i = 2, \\ 0 & \text{otherwise} \end{cases}$$

one can easily get the followings

$$\begin{aligned} R_{12,k} &= 2B_k R_{12} + B_1 R_{k2} + B_2 R_{1k}, \\ R_{13,k} &= 2B_k R_{13} + B_1 R_{k3} + B_3 R_{1k}, \\ R_{14,k} &= 2B_k R_{14} + B_1 R_{k4} + B_4 R_{1k}, \\ R_{23,k} &= 2B_k R_{23} + B_2 R_{k3} + B_3 R_{2k}, \\ R_{24,k} &= 2B_k R_{24} + B_2 R_{k4} + B_4 R_{2k}, \\ R_{34,k} &= 2B_k R_{34} + B_3 R_{k4} + B_4 R_{3k}, \\ R_{11,k} &= 2B_k R_{11} + B_1 R_{k1} + B_1 R_{1k}, \\ R_{22,k} &= 2B_k R_{22} + B_2 R_{k2} + B_2 R_{2k}, \\ R_{33,k} &= 2B_k R_{33} + B_3 R_{k3} + B_3 R_{3k}, \\ R_{44,k} &= 2B_k R_{44} + B_4 R_{k4} + B_4 R_{4k}, \end{aligned}$$

for $k = 1, 2, 3, 4$.

Consequently, we can state the following.

Theorem 3. *There exists a manifold $(\mathbb{R}^4, g)$ which is pseudo-Ricci symmetric for the above mentioned choice of the 1-forms.*

It can be easily shown that the manifold $(\mathbb{R}^4, g)$ under consideration fails to be a generalized pseudo-Ricci symmetric space.

4. EXISTENCE OF A (GPRS)₄-SPACE

Example 2. Let $(\mathbb{R}^4, g)$ be a 4-dimensional Riemannian space endowed with the Riemannian metric g given by

$$ds^2 = g_{ij}dx^i dx^j = e^{-x^1}[(dx^1)^2 + (dx^2)^2 + 2dx^3 dx^4]$$

for $i, j = 1, 2, 3, 4$. The non-zero components of Riemannian curvature tensor, Ricci tensors and scalar curvature are (up to symmetry and skew-symmetry)

$$R_{2324} = -\frac{1}{4}e^{-x^1} = R_{3434}, \quad R_{22} = -\frac{1}{2} = R_{34}, \quad r = -\frac{3}{2}e^{x^1}.$$

Covariant derivatives of Ricci tensors (up to symmetry) are expressed as

$$R_{12,2} = R_{13,4} = R_{14,3} = -\frac{1}{4}, \quad R_{22,1} = R_{34,1} = -\frac{1}{2}.$$

For the following choice of the 1-forms

$$B_i = \begin{cases} \frac{1}{4} & \text{for } i = 4, \\ 0 & \text{otherwise;} \end{cases}$$

$$C_i = \begin{cases} -\frac{e^{x^1}}{8} & \text{for } i = 4, \\ 0 & \text{otherwise} \end{cases}$$

one can verify the followings

$$\begin{aligned} R_{12,k} &= (A_k + B_k) R_{12} + A_1 R_{k2} + A_2 R_{1k} + (C_k + D_k) g_{12} + C_1 g_{k2} + C_2 g_{1k}, \\ R_{13,k} &= (A_k + B_k) R_{13} + A_1 R_{k3} + A_3 R_{1k} + (C_k + D_k) g_{13} + C_1 g_{k3} + C_3 g_{1k}, \\ R_{14,k} &= (A_k + B_k) R_{14} + A_1 R_{k4} + A_4 R_{1k} + (C_k + D_k) g_{14} + C_1 g_{k4} + C_4 g_{1k}, \\ R_{23,k} &= (A_k + B_k) R_{23} + A_2 R_{k3} + A_3 R_{2k} + (C_k + D_k) g_{23} + C_2 g_{k3} + C_3 g_{2k}, \\ R_{24,k} &= (A_k + B_k) R_{24} + A_2 R_{k4} + A_4 R_{2k} + (C_k + D_k) g_{24} + C_2 g_{k4} + C_4 g_{2k}, \\ R_{34,k} &= (A_k + B_k) R_{34} + A_3 R_{k4} + A_4 R_{3k} + (C_k + D_k) g_{34} + C_3 g_{k4} + C_4 g_{3k}, \\ R_{11,k} &= (A_k + B_k) R_{11} + A_1 R_{k1} + A_1 R_{1k} + (C_k + D_k) g_{11} + C_1 g_{k1} + C_1 g_{1k}, \\ R_{22,k} &= (A_k + B_k) R_{22} + A_2 R_{k2} + A_2 R_{2k} + (C_k + D_k) g_{22} + C_2 g_{k2} + C_2 g_{2k}, \\ R_{33,k} &= (A_k + B_k) R_{33} + A_3 R_{k3} + A_3 R_{3k} + (C_k + D_k) g_{33} + C_3 g_{k3} + C_3 g_{3k}, \\ R_{44,k} &= (A_k + B_k) R_{44} + A_4 R_{k4} + A_4 R_{4k} + (C_k + D_k) g_{44} + C_4 g_{k4} + C_4 g_{4k}, \end{aligned}$$

where $k = 1, 2, 3, 4$.

This motivates us to state the following.

Theorem 4. *There exists a manifold $(\mathbb{R}^4, g)$ which is generalized pseudo-Ricci symmetric for the above choice of the 1-forms.*

However, for the 1-form

$$B_i = \begin{cases} \frac{1}{2} & \text{for } i = 1, \\ 0 & \text{otherwise} \end{cases}$$

one can easily obtain the followings

$$\begin{aligned} R_{12,k} &= 2B_k R_{12} + B_1 R_{k2} + B_2 R_{1k}, \\ R_{13,k} &= 2B_k R_{13} + B_1 R_{k3} + B_3 R_{1k}, \\ R_{14,k} &= 2B_k R_{14} + B_1 R_{k4} + B_4 R_{1k}, \\ R_{23,k} &= 2B_k R_{23} + B_2 R_{k3} + B_3 R_{2k}, \\ R_{24,k} &= 2B_k R_{24} + B_2 R_{k4} + B_4 R_{2k}, \\ R_{34,k} &= 2B_k R_{34} + B_3 R_{k4} + B_4 R_{3k}, \\ R_{11,k} &= 2B_k R_{11} + B_1 R_{k1} + B_1 R_{1k}, \\ R_{22,k} &= 2B_k R_{22} + B_2 R_{k2} + B_2 R_{2k}, \\ R_{33,k} &= 2B_k R_{33} + B_3 R_{k3} + B_3 R_{3k}, \\ R_{44,k} &= 2B_k R_{44} + B_4 R_{k4} + B_4 R_{4k}, \end{aligned}$$

for $k = 1, 2, 3, 4$. In consequence of the above, one can see the following.

Theorem 5. *There exists a manifold $(\mathbb{R}^4, g)$ which a generalized pseudo-Ricci symmetric in some cases and a pseudo-Ricci symmetric in some other cases depending on the choice of the 1-forms.*

It is obvious that the manifold under consideration cannot be Ricci symmetric, Ricci recurrent and generalized Ricci recurrent.

REFERENCES

- [1] B. Y. Chen, K. Yano, *Hypersurfaces of a conformally flat space*, Tensor (N.S.), 26 (1972), 318-322.
- [2] C. A. Mantica, L. G. Molinari, *Weakly Z symmetric manifolds*, Acta Math. Hungar., 135 (2012), 80-96.
- [3] C. A. Mantica, Y. J. Suh, *Pseudo Z symmetric Riemannian manifolds with harmonic curvature tensors*, Int. J. Geom. Meth. Mod. Phys., 9 (2012), No. 1250004, pp. 21.
- [4] E. M. Patterson, *Some theorems on Ricci recurrent spaces*, J. London. Math. Soc., 27 (1952), 287-295.
- [5] K. K. Baishya, *On generalized weakly symmetric manifolds*, Bull. Transilv. Univ. Braşov Ser. III. Math. Comput. Sci., 10(59)(1) (2017), 31-38.
- [6] K. K. Baishya, *Note on almost generalized pseudo Ricci symmetric manifolds*, Kyungpook Math. J., 57(3) (2017), 517-523.
- [7] K. Yano, M. Kon, *Structures on manifolds*, Series in Pure Mathematics, 3, World Scientific Publishing, Bucharest, 2008.

- [8] M. C. Chaki, *On pseudo Ricci symmetric manifolds*, Bulg. J. Physics, 15 (1988), 526-531.
- [9] M. C. Chaki, T. Kawaguchi, *On almost pseudo Ricci symmetric manifolds*. Tensor (N.S.), 68(1) (2007), 10-14.
- [10] R. Deszcz, M. Glogowska, M. Hotlos, Z. Senturk, *On certain quasi-Einstein semisymmetric hypersurfaces*, Ann. Univ. Sci. Budapest. Eötvös Sect. Math., 41 (1998), 151-164.
- [11] R. S. D. Dubey, *Generalized recurrent spaces*, Indian J. Pure Appl. Math., 10 (1979), 1508-1513.
- [12] S. Mukhopadhyay, B. Barua, *On a type of non-flat Riemannian Manifold*, Tensor (N.S.), 56 (1995), 227-231.
- [13] U. C. De, N. Guha, D. Kamilya, *On generalized Ricci recurrent manifolds*, Tensor (N.S.), 56 (1995), 312-317.
- [14] U. C. De, A. K. Gazi, *On Pseudo-Ricci Symmetric manifolds*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.), 58(1) (2012), 209-222.

Received November 22, 2016.

DEPARTMENT OF MATHEMATICS,
 KURSEONG COLLEGE, KURSEONG,
 DARJEELING-734203, WEST BENGAL, INDIA
Email address: kanakkanti.kc@gmail.com